ANAESTHETIC MANAGEMENT OF AN HEMIGLOSSECTOMY IN A “FULL STOMACH”

PATIENT CASE REPORT

Bruno Lima¹; Carolina Martins²; Sara Tomé²; Manuel Chedas²

1- Anaesthesiology Department, Hospital de Braga (HB). Braga, Portugal

2- Anaesthesiology Department, Hospitalar de Lisboa Ocidental. Lisboa, Portugal

INTRODUCTION

Anaesthetic management in a patient with a full stomach poses challenges to the anaesthesia provider: The prevention of gastric regurgitation and pulmonary aspiration while instituting an appropriate airway management. Trachea structures are innervated by the recurrent laryngeal nerve and sympathetic nerve fibers, derived mainly from the middle cervical ganglion. Cervical plexus block could be indicated for tracheostomy, providing analgesia or anesthesia of the neck. There is evidence that superficial cervical plexus blocks are at least equivalent to deep blocks, being significantly safer.

CASE REPORT

PATIENT

Male, 54 years old, ASA III with a malignant tumor of the tongue with cervical ganglion invasion

KNOWN CO-MORBIDITIES

Chronic venous insufficiency with bilateral intermittent claudication for 100 meters

SURGERY

Tracheostomy + Hemiglossectomy + Radical neck dissection

Early surgery could be potentially curative avoiding radical glossectomy and free flap surgery. Miscommunication led to a full stomach patient at the time of the surgery. Due to the urgency of the procedure both the patient and all the staff involved decided to go on with the surgery. At first the patient, with his consentment was submitted to surface based bilateral superficial cervical plexus block, enabling awake tracheostomy, followed by intravenous anaesthetic induction and manutention.

ANAESTHETIC TECHNIQUE

Combined Anaesthesia

- **Surface based, Bilateral Superficial Cervical Plexus Block**
- **Total Intravenous Anaesthesia**

Figure 2: Anatomical references for surface based superficial cervical plexus block; SCM- Sternocleidomastoid muscle

MONITORING

Standard ASA + monitoring bispectral index (BIS® to a target of 40 to 60) + Neuromuscular block monitoring (Train of four)

Bilateral Superficial Cervical Plexus Block (5mL 0,5% Ropivacaine on each side) followed by **Tracheostomy** with awake patient, before anaesthetic induction

INDUCTION

- Remifentanil TCI 5 ng/ml
- Propofol TCI 5ng/ml
- Cisatracurium besilate 0,1 mg/kg

MAINTENANCE

- Remifentanil TCI 3,5 - 5 ng/ml
- Propofol TCI 2 – 3,5 ng/ml
- Cisatracurium besilate 0,07 mg/kg bolus every 45-60 min
- O2/air mixture
- MAP 65-70 mmHg, HR 80-90 bpm, SpO2 95-100%
- EtCO₂ 28-33 mmHg

EVENTS

During radical neck dissection accidental ligation of the right jugular vein.

At the end of the surgery the patient was transferred ventilated and sedated to the ICU.

FOLLOW-UP

First post-op day

The patient was awoken, spontaneously breathing through the tracheostomy tube. No respiratory complications were reported.

Fifteenth post-op day

Severe dysphagia, deglutition therapy was proposed.

First post-op month

Tongue mobility diminished as well pharynx sensibility both responsible for dysphagia. Proposed a PEG dispositive.

DISCUSSION

Securing the airway in a awake patient with full stomach avoids loss of protective reflexes and the risk of failed tracheal intubation.

Bilateral cervical ganglion plexus block provided awake tracheostomy without any complication of a full stomach patient and allowed for peri-operative analgesia for performing radical neck dissection.

CONCLUSIONS

In the reported case, cervical plexus block was useful for performing awake tracheostomy in the patient at risk for pulmonary aspiration, proposed for urgent surgery, to maintain upper airway reflexes intact. These type of block should be performed by skilled professionals.

REFERENCES

- El-Boghdady K, et al. Phrenic Nerve Palsy and Regional Anesthesia for Shoulder Surgery: Anatomical, Physiologic, and McGoldrick KE. Day-case Shoulder Surgery: Anesthetic Challenges. Ambulatory Surgery, 2012; 17(3): 47-49.
- Mulier J. Anestesia libre de opioides: ¿un cambio de paradigma? Rev Esp Anesthesiol Reanim. 2017;64:427-430.
- Palhais N, et al. Extrafascial injection for interscalene brachial plexus block reduces respiratory complications compared with a conventional intrafascial injection: a randomized, controlled, double-blind trial. Br J Anaesth. 2016 Apr; 116(4): 531-537.
- Tran DQ, et al. Diaphragm-Sparing Clinical Considerations. Anesthesiology. 2017 Jul; 127(1):173-191.
- Nerve Blocks for Shoulder Surgery. Reg Anesth Pain Med. 2017 Jan/Feb; 42(1):32-38.