CENTRAL LINE ASSOCIATED BLOODSTREAM INFECTIONS IN PEDIATRIC UNITS IN GREECE

Kouni S¹, Dimolitsa C², Iosifidis E³, Kalampalikis P⁴, Kapetanakis I⁵, Karakosta P¹, Katsibardi K⁶, Kazantzi M⁴, Kourkouni E¹, Mexi P⁷, Nikaina E.⁸, Papadopoulou S⁹, Petrikkos L¹⁰, Siahandidou T⁸, Skordala M¹¹, Tragiannidis A¹², Tsolia M¹³, Zanikos K¹⁴, Zaoutis E.T¹

¹. Center for Clinical Epidemiology and Outcomes Research (CLEO), Non-Profit Civil Partenrship, Athens, Greece. ². Paediatric Intensive Care Unit, General Children's Hospital of Athens "P. & A. Kyriakou", Athens, Greece. ³. Paediatric Infectious Disease Specialist University Scholar, 3rd Pediatric Department, Aristotle University of Thessaloniki, Thessaloniki, Greece. ⁴. Paediatric Intensive Care Unit, General Children's Hospital of Athens "Agia Sofia", Athens, Greece. ⁵. Neonatal Intensive Care Unit, General Children's Hospital of Athens "P. & A. Kyriakou", Athens, Greece. ⁶. First Department of Paediatrics, National and Kapodistrian University of Athens, General Children's Hospital of Athens "Agia Sofia", Athens, Greece. ⁷. Neonatal Intensive Care Unit, University General Hospital of Athens «Attikon», Athens, Greece. ⁸. Neonatal Unit of the First Department of Paediatrics, National & Kapodistrian University of Athens, Athens, Greece. ⁹. Neonatal Intensive Care Unit, General Children's Hospital of Athens "Agia Sofia", Athens, Greece. ¹⁰. Department of Paediatric Hematology-Oncology, General Children's Hospital of Athens "Agia Sofia", Athens, Greece. ¹¹. Neonatal Department, Neonatal Intensive Care Unit, University General Hospital of Alexandroupolis, Greece. ¹². 2nd Hematology Oncology Unit, 2nd Paediatric Department, Aristotle University of Thessaloniki, AHEPA Hospital, Thessaloniki, Greece. ¹³. Second Department of Paediatrics, General Children's Hospital of Athens "P. & A. Kyriakou", University of Athens, Greece. ¹⁴. A Neonatal Intensive Care Unit, General Children's Hospital of Athens "Agia Sofia", Athens, Greece.

Table 1. Enrolled facilities contributing data used in this report, during the surveillance period June 2016 to February 2017

Type of acute care facility	No of locations	No of CLABSI	No of MBI* CLABSI	Total Central line days	Total patient days	Total antibiotic days	25%	median	75%
NICUs	14	55	0	9141	62368	23027	291	359	401
PICUs	4	21	0	3430	5059	2977	333	563	848
ONCs	6	35	15	12572	14881	8725	445	477	619
ALL UNITS	24	111	15	25143	82308	34729			

LOT=(no of calendar days in which antibiotics were received/ total number of patient days)*1000
*Mucosal Barrier Injury Laboratory

Results:

A total of 111 CLABSI episodes were recorded. Among pediatric oncology patients, a total of 35 CLABSI were identified during the study period, 15/35 (42.86%) were classified as MBI-LCBI (Table 1). The monthly CLABSI rates for all units across the whole surveillance period are presented in Figure 1. Stratified analysis by ward type (NICUs, PICUs and ONCs) revealed that the higher CLABSI rates were noticed in the PICUs and NICUs, and the higher central line utilization ratio was in ONCs (Table 2). The stratification of CLABSI rates among the units were 6.02 for NICUs, 6.12 for PICUs and 2.78 infections per 1000 central line (CL) days for ONCs (Table 2).

Figure 1. Monthly CLABSI rate and overall CLABSI rate among all units

A total of 123 pathogens isolated in 111 CLABSI episodes were observed among all units. Gram positive cocci were most frequently reported in PICUs and Enterobacteriaceae were the most common CLABSI pathogens in NICUs and ONCs. Overall 37% of gram-negative pathogens were resistant to carbapenems and 38.5 % were resistant to third generation cephalosporins.

Table 2. Pooled means and key percentiles of the distribution of CLABSI rates and CLU ratios by type of location

Type of acute care facility	CLABSI RATE				CLU RATIO			
	Pooled mean	25%	Median	75%	Pooled mean	25%	Median	75%
NICUs	6.02	3.83	6.96	9.9	0.15	0.06	0.13	0.26
PICUs	6.12	4.77	6.24	16.5	0.68	0.38	0.72	0.89
ONCs	2.78	1.65	2.48	3.65	0.84	0.79	0.83	0.97

CLABSI RATE=(no of CLABSIs/central line days)*1000
CLU RATIO=(no of central line days/ no of patient days)

Conclusions:

Nationally representative CLABSI rates were determined for pediatric patients. Of significant concern was the identification of high rates of carbapenem-resistant organisms. These data could be used to benchmark and serve as baseline data for the design and evaluation of infection control and antimicrobial stewardship interventions.