Optimization Of Anesthetic Management In Free Flap Reconstruction Surgeries “Before-after Study”

H. KAMGA¹, F.VIDAL², H.ENATEAU³, JL. HANOUZ⁴ - Caen University Hospital Center, France

1. Consultant of Anesthesia & Critical Care 2. Anesthesia & Critical Care Resident 3. Professor of Maxillofacial surgery 4. Professor of Anesthesia & Critical Care

BACKGROUND : Free flap reconstruction is nowadays the best technic after complex surgery of the head and neck. In our center, a professional practices evaluation showed a 74% rate of postoperative complication. The practices have been modified by virtue of the latest evidences and this study aim is to evaluate these modifications.

METHODS : This is a single center before-after design study. All patients who underwent free flap surgery within the study period were included. Table1 shows the points of difference between the two periods.

The main outcome is the rate of postoperative complications. The major complications are flap loss or any medical complications required ICU admission. The minor complications are surgical revision for wound disunion and surgical site infection. The Chi-2 is use for categorical data. An inverse weight adjustment of the propensity score was performed.

	PROTOCOL « Before »	PROTOCOL « After »
Period	2009 - 2014	2015 - 2017
Catecholamine	Dobutamine	Noradrenaline
Vascular filling	Liberal	Monitored
Blood Pressure Level	MAP = 90 mmHg	Basic MAP: No more than 30% less.
Regional Anesthesia	Optional	Systematic
Hemoglobin Objective	10 g/dL	7-8-9-10 g/dL according to clinical state of the
Depth of Anesthesia	Non monitored	Monitored

RESULTS AND DISCUSSION : 31 patients are in the "before" group and 34 in the "after". The characteristics of the patients are comparable. The rate of postoperative complication are shown in Figure1.

Optimization of preoperative care for free flap reconstruction reduces the rate of postoperative complications. This results is in approval with the recent recommendations of the ERAS group in major head and neck cancer surgery

Figure1: Postoperative complication rate