A Comparative Evaluation Of Quadratus Lumborum Block Versus Fascia Iliaca Nerve Block For Patients Undergoing Elective Hip Replacement Surgeries

hashmi j^{1*}, cusack b², hughes l³, singh v⁴, srinivasan k⁵.

Tallaght University Hospital,Dublin,Ireland

^{1,2,3,4} trainee anaesthesia, ^{*}corresponding author, ⁵ Consultant anaesthesia ,

Introduction

Total hip replacement is a commonly performed surgical procedure with up to 4500 procedures performed annually in Ireland. Multimodal analgesia together with peripheral nerve block is recommended as gold standard for management of lower limb joint replacements. The limitations with peripheral nerve blocks include limited duration and associated motor block which can delay rehabilitation.

Aim

The aim of our study was to compare the quality of analgesia and the effect on motor blockade of US-guided TQL with that of US-guided fascia iliaca block (FIB). We tested the primary hypothesises that a single injection TQL block when compared to single injection fascia iliaca block (FIB) will reduce cumulative morphine consumption in the first 24 hours. Secondary outcome analysed compared the motor block, quality of analgesia and other additional measures.

Study design

Statistical Analysis

Normally distributed data are presented as mean ± standard deviation, and the comparison between groups were performed using unpaired t-test and chi2 test

The comparison for VAS between group were performed using Mann Whitney U test .The comparison of Motor block between the two groups was done using Chi2 test.

P <0.05 was considered a statistically significant difference.

Results

Table 1 Comparison of baseline characteristics of study participants between groups:

Patient characteristics	Group A (N=24) N (%) Mean (SD)	Group B (N=24) N (%) Mean (SD)	Total N	P value
Sex				
Male	17 (60.7)	11 (39.3)	28	0.079
Female	7 (35.0)	13 (65.0)	20	
ASA				
1	3 (42.9)	4 (57.1)	7	0.714
2	16 (48.5)	17 (51.5)	33	
3	5 (62.5)	3 (37.5)	8	
Weight in kilograms	84.1 (14.1)	76.3 (16.2)		0.820
Operative side				
Left side	10 (50.0)	10 (50.0)	20	1.000
Right side	14 (50.0)	14 (50.0)	28	

Table. 2 Comparison of analgesia use between groups:

Analgesia use	Group A (N=24) N (%)	Group B (N=24) N (%)	Total N=48	P value*
Pre opioids use, 48 hours preceding surgery				
No	22 (48.9)	23 (51.1)	45	0.551
Yes	2 (66.7)	1 (33.3)	3	
Intra operative opioids use				
No	23 (50.0)	23 (50.0)	46	1.000
Yes	1 (50.0)	1 (50.0)	2	
Intra operative paracetamol administration				
No	3 (60.0)	2 (40.0)	5	0.637
Yes	21 (48.8)	22 (51.2)	43	
Intra operative NSAIDS administration				
No	12 (54.5)	10 (45.5)	22	0.562
Yes	12 (46.2)	14 (53.8)	26	
Intra operative other analgesics administration				
No	24 (50.0)	24 (50.0)	48	NA
Yes	0 (0.0)	0 (0.0)	0	

Table 3 Comparison of duration of surgery, morphine use and post-operative pain scale between the groups:

	Group A (N=24) Mean (SD)	Group B (N=24) Mean (SD)	P value*
Duration of surgery in minutes	98.6 (20.9)	95.6 (32.3)	0.704
Morphine usage in 24 hours #	38.3 (23.3)	35.8 (20.2)	0.699

Table 4. Comparison of post-operative pain using verbal rating scale between the groups

Post-operative pain monitoring	Group A Median (IQR)	Group B Median (IQR)	P value*
At rest – 6 hours post block	4 (2-5)	2 (0.2-3.7)	0.118
On movement – 6 hours post block	4 (2-7)	3 (1-6.5)	0.540
At rest – 24 hours post block	2.5 (1-4)	3.0 (2.0-5.0)	0.260
On movement – 24 hours post block	5 (3.2-6.7)	5.5 (4.0-8.0)	0.383

Table 5. Comparison of motor block (Modified Bromage scale) at 6 hours post block between the groups:

Modified Bromage scale category for motor block	Group A (N=24) N (%)	Group B (N=24) N (%)	Total N=48	P value*
Complete block (unable to move feet or knees)	1 (50.0)	1 (50.0)	2	0.497
Almost complete block (able to move feet only)	2 (50.0)	2 (50.0)	4	
Partial block (just able to move knees)	7 (43.8)	9 (56.3)	16	
Detectable weakness of hip flexion while supine (full flexion of knees)	9 (69.2)	4 (30.8)	13	
No detectable weakness of hip flexion while supine	2 (25.0)	6 (75.0)	8	
Able to perform partial knee bend	3 (60.0)	2 (40.0)	5	
Total	24 (50.0)	24 (50.0)	48	

Discussion

Our randomised controlled study demonstrated that QLB although as equally effective as FINB didn’t provide any superior pain control either in quality or duration of pain relief nor had any lesser motor block compared to FINB.

As the QLB is more challenging to do and have greater potential for side effects and complications like injury to liver, spleen and kidney we propose that FINB should remain the preferred option for hip replacement surgeries as it is technically easier to do with less likelihood of complications. We need more studies to confirm whether or not QLB can be a viable option .Further studies with ideally greater numbers of patients are required.

BIBLIOGRAPHY

1.Ueshima H,YoshiyamaS,Otake H. The ultrasound-guided continuous transmuscular quadratus lumborum block is an effective analgesia for total hip arthroplasty. Journal of clinical anesthesia. 2016;31:35.

2.La Colla L,Uskova A, Ben-David B. Single-shot Quadratus Lumborum Block for Postoperative Analgesia After Minimally Invasive Hip Arthroplasty: A New Alternative to Continuous Lumbar Plexus Block? Regional anesthesia and pain medicine. 2017;42(1):125-126.