Post tonsillectomy analgesia; quality improvement project

Al-Zahid S, Norris B, Elias-Jones W, Wilson H
Royal Devon and Exeter Hospital, Exeter, UK

Aims

- 1) To ensure patients post tonsillectomy are provided with adequate analgesia
- 2) To reduce GP attendance post tonsillectomy

Methods

- Prospective cohort study Nov 15 – July 16
- Review of analgesia on discharge summary
- Contact GP's to review patient attendance within one month post-tonsillectomy
- Gold standard (Local post tonsillectomy guideline)
 - Paediatrics:
Age appropriate Paracetamol + Ibuprofen
Tramadol for >50kg or >12 yrs
 - Adult:
Paracetamol + Ibuprofen + Tramadol

Mid-study intervention

April 2016 - Introduction of automated discharge summaries with IT help

Paediatrics:

Enter child's weight, D/C summary pre-populates with age appropriate analgesia

Adults:

Enter post tonsillectomy protocol, pre-populates paracetamol, Ibuprofen and Tramadol on discharge summary

Electronic discharge summary

Infection Control issues (eg MRSA carriage, resolved u.d. infection):					
Other Infection Control Issues?					
KNOWN	Select Details:	BMI:	kgm ⁻²		
ALLERGY:		Smoker:	Select		
+ GP TO DO PLEASE: Post-op pain regime SELECT					
DRUGS ON DISCHARGE:					
Drug - (block capitals)	Dose	Freq	Route	Notes	Qty
1					
2					
3					
4					
5					
Follow-up with in					
Comments:					
Signed:		Grade:	Sleep:	Date:	
Dictated:					
Clinically Ready by:					
GP	GP VIA PATIENT	MEDICAL RECORD	SECRETARY	PHARMACY / MEDICAL APPTS	

Results

Totals – 141 patients

Adults 58

Paediatrics 83

Analgesia protocol followed – 112

Analgesia protocol not followed – 29

Attended GP – 58 (41%)

Did not attend GP – 83 (59%)

9% lower GP attendance rate in group of patients where post tonsillectomy analgesia protocol followed
Not statistically significant (P=0.381)

GP attendance

No statistically significant reduction in GP attendance following intervention (P value 0.940)

Commonest reasons for GP attendance

Further analgesia

Antibiotics

Re-assurance

Discussion

GP attendance post tonsillectomy very high

Group with optimised analgesia showed reduction in GP attendance rate although not statistically significant

Attendance to GP post-tonsillectomy is multi-factorial and not necessarily always related to analgesia. Re-assurance was an important reason and reflects the close relationship between patients and their GP's.

Automated discharge summaries improve the provision of appropriate analgesia post tonsillectomy.

Royal Devon and Exeter
NHS Foundation Trust