Comparison between palpation and ultrasound of the punctured spinal level in pregnant women

Hermann S. Fernandes (MD, DESA), Shirley A. Santos (MD), André C. A. Ventura (MD), Fernando S. Nani (MD), Cristiane M. F. Haddad (MD), Joaquim E. Vieira (MD, PhD)

Anaesthesiology Department – Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo – Brazil

Rationale and Objectives

Neuroaxial anesthesia is a widely used technique in obstetrics, with several advantages over general anesthesia: it avoids the manipulation of the airway, allows the woman to interact with the child at birth, lower risk of aspiration of gastric contents. However, it is not risk-free, such as hypotension, total spinal anesthesia, spinal cord injury.

Iatrogenic neurological damage after spinal anesthesia is a rare event in daily practice. One of the reasons related to this type of injury is the misleading punctured level. There is evidence in the literature that the lower end of the spinal cord is related to L2 in 43% of women and only 27% in men. There are articles that relate up to 63% of error in the punctured level. Therefore, this project aimed to evaluate the puncture level accuracy rate in pregnant patients.

Methods

An observational study was carried out in a university hospital. Patients over 18 years of age who underwent neuraxial anesthesia or analgesia during labor or cesarean delivery were included. After the neuraxial technique (epidural, spinal or combined spinal-epidural), the site of successful puncture was marked.

After the end of the procedure the patient was then placed back in the same position where the initial puncture was performed (sitting down position) and the patient was examined with ultrasound. It starts with the probe on vertical paramedian position, on lower back. After visualization of the sacrum, the probe is moved cephalad, keeping the visualization of the vertebral laminae. It is possible to count the spaces, from the sacrum to cephalad, until the puncture site is reached.

Figure 1. Technique for lumbar spine ultrasound

Results

A total of 44 patients were analyzed from April to December of 2017. The mean age was 29.25 ± 6.44 years and mean BMI of 30.74 ± 8.11 .

The incompatibility rate between the two methods was 45.45%. As showed on table 1, by palpation, the most frequent believed space punctured was L3-L4 (68.2%), but it was not correct in 46.66% of these cases, when checked with ultrasound. Table 2 shows the incompatibility rate by experience of the professional.

LEVEL OF PUNCTURE	PALPATION N (%)	ULTRASOUND N (%)
L1-L2	-	1 (2.3%)
L2-L3	4 (9.1%)	14 (31.8%)
L3-L4	30 (68.2%)	21 (47.7%)
L4-L5	10 (22.7%)	8 (18.2%)
L5-S1	-	-

Table 1

EXPERIENCE OF PROFESSIONAL	INCOMPATIBILITY RATE PALPATION - ULTRASOUND
1 st YEAR MEDICAL RESIDENT (N=8)	62.5%
2 nd YEAR MEDICAL RESIDENT (N=28)	55.5%
3 rd YEAR MEDICAL RESIDENT (N=3)	0%
ASSISTANT ANESTHETIST (N=5)	0%

Table 2

Conclusions

A non-negligible error rate was observed, however, with results compatible with the literature, and the level is usually punctured at the immediately superior level, when it is not the same.

References

- Eur J Anaesthesiol. 2015 Jul;32(7):499-505. Spinal ultrasound versus palpation for epidural catheter insertion in labour: A randomised controlled trial. Arzola C1, Mikhael R, Margarido C, Carvalho JC.
- Int J Obstet Anesth. 2016 Dec;28:12-16. Epub 2016 Jul 20. Ultrasound to identify the lumbar space in women with impalpable bony landmarks presenting for elective caesarean delivery under spinal anaesthesia: a randomised trial. Creaney M, Mullane D, Casby C, Tan T.
- Schlotterbeck H et al. Ultrasonographic control of the puncture level for lumbar neuraxial block in obstetric anaesthesia, Br J Anaesth 2008; 100: 230–4

Euroanaesthesia
The European Anaesthesiology Congress

2018