Childhood medical history and psychosis in adult life

Findings from the Bologna EU-GEI incidence and case-control study

Giuseppe D'Andrea, MD,^{1 4} Federico Suprani, MD,⁴ Elena Tolomelli, MD,² Monia Gennari, MD, PhD² Marcello Lanari, MD,² Giacomo Faldella, MD ² Roberto Muratori, MD,³ Domenico Berardi, MD,^{1 3} Ilaria Tarricone, MD, PhD,^{2 3 4}

¹Department of Biomedical and NeuroMotor Sciences, University of Bologna ²Department of Medical and Surgical Sciences, University of Bologna ³Department of Mental Health, AUSL Bologna, ⁴Bologna Transcultural Psychosomatic Team (BoTPT), Department of Medical and Surgical Sciences, University of Bologna

Introduction

Childhood adversity is a widely replicated risk factor for psychosis. Abuse and maltreatment during infancy have been associated with a more than twofold increase in the odds of psychosis.

Severe medical conditions that require hospital care can represent a challenging adversity for children; only one prior study has investigated the relationship between childhood medical history and risk of psychosis in adult life, finding that severe injuries during adolescence may predict later onset of psychosis.

The aim of this retrospective case-control study was to determine if a history of pediatric hospitalization or Emergency Room (ER) visits for any non-psychiatric medical reason could increase the odds of psychosis onset in adult life.

Methods

A sample of 70 cases with first-episode psychosis (FEP) and 65 controls was recruited between May 2010 and April 2015 in Bologna within the EU-GEI project.

- Cases were recruited assessing all individuals referring for the first time to the Community Mental Health Service of Bologna. Patients aged 18-64 and diagnosed with FEP were included.
- Controls were recruited in the same time span and catchment area. They had to be aged 18-64 and never treated with antipsychotic medications.

History of ER visits and hospital admissions before the age of 18 were retrospectively retrieved from the available digital records of three public hospitals in the city of Bologna. Data collected included age at the moment of the event and discharge diagnosis. Results were adjusted for age, sex, migrant status, cannabis use (CEQ), parental social class (SES), and childhood trauma (CTQ). All measures were collected with validated instruments.

Results

≥1 hospitalization: 20.0% vs 6.2% p=.02

≥2 hospitalizations 10.0% vs 1.5% p=.06

≥1 ER visit: 21.7% vs 4.6% p<.001

≥2 ER visits: 17.0% vs 3.1% p=.01

The number of admissions between cases and controls was significantly different in the age group 12-17, regarding both hospital and ER admissions (p=.02).

Both hospitalization (OR=3.81, 95% CI=1.19-12.27, p=.01) and ER visit (OR=7.69, 95% CI=2.16-27.49, p<.001) increased the odds of psychosis in adult life.

In multivariable logistic regression analysis the association remained significant after controlling for socio-demographic factors (age, sex, migrant status, and parental social class), cannabis use, and trauma history.

Conclusion

A robust association between hospitalizations before age of 18 and FEP was observed. Children and adolescents with this clinical history may be at very high risk of psychotic disorders. Further studies with a prospective design and a larger sample are needed to confirm this association.

References

- Varese F, Smeets F, Drukker M, Lieverse R, Lataster T, Viechtbauer W, Read J, van Os J, Bentall RP. Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull.* 2012 Jun;38(4):661-71
- S. Luoma, H. Hakko, M. Marttunen, A. Taanila, S. Lindeman, Severe injuries in adolescence predict psychosis: A nested case control study of the Northern Finland 1966 Birth Cohort, *Schizophr. Res.* 118 (2010) 48–53. doi:10.1016/j.schres.2009.12.031
- European Network of National Networks studying Gene-Environment Interactions in Schizophrenia (EU-GEI), van Os J, Rutten BP, et al. Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. *Schizophr Bull.* 2014;40(4):729-736
- Barker V, Gumley A, Schwannauer M, Lawrie SM. An integrated biopsychosocial model of childhood maltreatment and psychosis. *Br J Psychiatry.* 2015 Mar;206(3):177-80