Predictor Factors Of Postoperative Pain In Total Hip And Knee Arthroplasties

Bárbara Gouveia¹, Sara Fonseca¹, Daniela Xará¹, Inês Valdoleiros ², Sofia Ferraz ¹, André Sá Rodrigues³

1 Centro Hospitalar São João, Porto, 2 Centro Hospitalar do Baixo Vouga, Aveiro, 3 Hospital da Prelada, Porto

BACKGROUND AND GOAL OF STUDY

Major orthopedic surgeries are associated with postoperative pain (POP)⁽¹⁾. The purpose of this study was to identify predictor factors of POP associated with the total knee (TKA) and total hip (THA) arthroplasties.

MATERIALS AND METHODS

Prospective cohort study

Population: 95 adult patients admitted to elective THA and TKA, between september and december 2014

Table I– Variables analyzed in the pre and postoperative questionnaires

Preoperative questionnaire	Postoperative questionnaire
<ul style="list-style-type: none">Demographic variablesType of arthroplastyASA physical statusBody Mass Index (BMI)Preoperative chronic painDepression diagnosisDepressive profileMedication for depression	<ul style="list-style-type: none">Intensity of POP (<i>Brief Pain Inventory</i>)- Minimum- Average- Maximum- At the time

Descriptive analysis of variables was performed and non-parametric tests were used in the analysis of POP levels. It was performed a multiple linear regression to determine the significant predictors of the POP.

RESULTS

We included 95 patients, all caucasian, 58 of which underwent TKA and 37 to THA.

Table II – POP according to sex, age, BMI, ASA physical status, type of arthroplasty, depression diagnosis, depression medications, preoperative chronic pain

		Minimal POP	p	Average POP	p	Maximum POP	p	At the time POP	p
Sex	Female	1,47+1,77 (0-6)	0,717	4,84+1,59 (0-8)	0,020*	6,97+2,19 (3-10)	0,010*	2,49+2,71 (0-9)	0,238
	Male	1,15+1,26 (0-4)		4,04+1,81 (1-8)		5,74+1,87 (3-10)		3,19+2,59 (0-7)	
Age	<65	1,24+1,58 (0-6)	0,379	4,46+1,53 (0-8)	0,416	6,44+2,01 (3-10)	0,383	2,53+2,83 (0-8)	0,417
	≥ 65	1,56+1,73 (0-6)		4,8+1,87 (1-8)		6,85+2,36 (3-10)		2,88+2,5 (0-9)	
BMI	≤ 18	0,5+0,71 (0-1)	0,088	4,5+3,54 (2-7)	0,727	6,5+4,95 (3-10)	0,583	1,5+2,12 (0-3)	0,526
	>18 and <25	0,67+1,11 (0-3)		4,13+1,3 (2-6)		6+2,14 (3-9)		3,53+2,88 (0-8)	
	≥ 25and<30	1,28+1,32 (0-4)		4,58+1,69 (0-8)		6,58+2,05 (3-10)		2,47+2,43 (0-7)	
	≥ 30and<35	1,52+1,93 (0-6)		4,85+1,66 (2-8)		7,15+2,21 (3-10)		2,42+3,05 (0-9)	
	≥ 35	3+ 2,27 (0-6)		4,88 +2,17 (2,8)		6,25+2,19 (3-10)		3,38+2,56 (0-7)	
ASA physical status	1	1+1 (0-2)	0,591	4,67+ 3,06 (2-8)	0,246	6+3,61 (3-10)	0,131	2+3,46 (0-6)	0,952
	2	1,31+1,62 (0-6)		4,52 + 1,61 (0-8)		6,49+2,11 (3-10)		2,72+2,7 (0-9)	
	3	2+1,96 (0-6)		5,38+1,89 (1-8)		8+1,91 (4-10)		2,77+2,83 (0-7)	
	4	1+0 (1-1)		4+0 (4-4)		6+0 (6-6)		3+0 (3-3)	
Type of arthroplasty	THA	0,89+1,17 (0-4)	0,052	3,92+1,85 (0-8)	0,002	5,84+2,2 (3-10)	0,005	2,84+2,76 (0-8)	0,725
	TKA	1,69+1,83 (0-6)		5,05+1,42 (2-8)		7,12+2 (3-10)		2,58+2,65 (0-9)	
Depression diagnosis	No	1,41+1,6 (0-6)	0,655	4,37+1,81 (0-8)	0,070	6,1+1,94 (3-10)	0,003*	3,14+2,71 (0-8)	0,048*
	Yes	1,33+1,74 (0-6)		5+1,39 (2-8)		7,47+2,27 (3-10)		1,94+2,5 (0-9)	
Depression medication	No	0,67+0,82 (0-2)	0,509	5,17+1,17 (4-7)	0,777	8+2,76 (3-10)	0,437	1+1,26 (0-3)	0,475
	Yes	1,47+1,85 (0-6)		4,97+1,45 (2-8)		7,37+2,2 (3-10)		2,13+2,65 (0-9)	
Chronic pain	No	0,92+1,5 (0-6)	0,078	3,84+1,62 (2-8)	0,003*	5,92+2,31 (3-10)	0,051	1,5+2,43 (0-7)	0,007*
	Yes	1,54+1,67 (0-6)		4,89+1,63 (0-8)		6,87+2,07 (3-10)		3,09+2,66 (0-9)	

* p < 0,05;

Table III– POP according to the type of arthroplasty by sex and age

		Minimal POP	p	Average POP	p	Maximum POP	p	At the Time POP	p
Female	THA	7	0,159	4,1 ± 1,79	0,025*	6,1 ± 2,36	0,033*	2,67 ± 2,9	0,777
	TKA	1,72 ± 1,93		5,17 ± 1,39		7,36 ± 2		2,4 ± 2,64	
Male	THA	0,88 ± 1,15	0,167	3,69 ± 1,96	0,152	5,5 ± 2	0,321	3,06 ± 2,64	0,776
	TKA	1,55 ± 1,37		4,55 ± 1,51		6,09 ± 1,7		3,4 ± 2,63	
<65	THA	0,87 ± 1,14	0,258	3,87 ± 1,66	0,029*	5,78 ± 2,02	0,032*	2,83 ± 2,89	0,583
	TKA	1,52 ± 1,81		4,9 ± 1,27		6,94 ± 1,88		2,3 ± 2,81	
≥ 65	THA	0,93 ± 1,27	0,122	4 ± 2,18	0,052	5,93 ± 2,56	0,082	2,86 ± 2,66	0,989
	TKA	1,89 ± 1,87		5,22 ± 1,58		7,33 ± 2,15		2,89 ± 2,47	

* p < 0,05;

Table IV– Multiple linear regression

	Minimal POP		Average POP		Maximum POP		At the Time POP	
	B (IC 95%)	p	B (IC 95%)	p	B (IC 95%)	p	B (IC 95%)	p
BMI ≤ 18			-2,64 (-5,24//0,04)	0,047*				
BMI >18 and <25			-2,22 (-3,66//0,77)	0,003*				
BMI ≥ 25 and <30			-1,56 (-2,86//0,25)	0,020*				
BMI ≥ 30and <35			-1,13 (-2,49//0,23)	0,103				
Depression	0,08 (-0,69//0,84)	0,847			-0,71 (-1,69//0,28)	0,156	-1,57 (0,46//2,69)	0,006*
Type of Surgery - THA	-0,84 (-1,53//0,15)	0,018			-0,67 (-1,59//0,26)	0,156		
Chronic Pain	0,55 (-0,21//1,32)	0,155	0,31 (-0,47//1,09)	0,429	0,10 (-0,90//1,10)	0,848	1,71 (-2,96//0,47)	0,008*
R ²	0,280		0,210		0,385		0,201	

* p < 0,05;

DISCUSSION AND CONCLUSION

Some predictive factors of POP are described after orthopedic surgery, but the most consistently identified are: age, type of surgery, preoperative pain, anxiety and depression ^(2,3).

In this study, women and younger patients undergoing TKA had higher levels of POP, however, age and gender were not identified as predictors of POP and physical status ASA didn’t interfere with POP intensity. We also found an association between the presence of chronic pain in the preoperative period and the development of more intense POP. Type of surgery, BMI, presence of chronic pain and absence of depression were identified as pain predictors after TKA and THA.

The identification of these predictors contributes to the individualized optimization of analgesic regimens, with a consequent improvement in postoperative pain recovery after THA and TKA.

(1) Barrington JW, Dalury DF, EmJoshi GP, Stulberg BN. "Improving patient outcomes through advanced pain management techniques in total hip and knee arthroplasty." Am J Orthop, 2013; S1-S20.

(2) Ip HY, Abrishami A, Peng PW, Wong J, Chung F. "Predictors of postoperative pain and analgesic consumption: a qualitative systematic review." Anesthesiology, 2009; 657-77.

(3) Pinto PR, McIntyre T, Ferrero R, Almeida A, Araújo-Soares V. "Predictors of acute postsurgical pain and anxiety following primary total hip and knee arthroplasty." J Pain, 2013; 502-15.