Is Public Access Defibrillation effective for the elderly OHCA patients same as the younger OHCA patients?

Kyoko TSUKIGASE¹⁾ • Hideharu TANAKA^{1,2)} • Hiroshi TAKYU²⁾

1) Research Institute of Disaster management and EMS, Kokushikan University,

2) Graduate school of EMS, Kokushikan University

Background/Aim

There is little information on elderly patients transported to the hospital resulting from out-of-hospital cardiac arrest (OHCA) in the world.

The aim of this study is to determine the effectiveness of PAD in elderly patients with OHCA coupled with the use of an AED by bystanders compared younger OHCA patients.

Method

Eligible 4,571 OHCA patients got PAD at the scene who were extracted from the Japanese Utsutein style registry from 2005 to 2014 (total 1,176,363 patients), were included and classified to four age categories (15-64, 65-74, 75-84, 85-104). Analysed Initial ECG when EMT arrived at the scene as PAD effectiveness. Analyzed Initial ECG when EMT's arrived at the scene were analyzed for PAD effectiveness. Multiple logistic regression analyses were performed to identify independent predictors of the good neurological outcome as the cerebral performance categories (CPC) 1 or 2.

Conclusion

Japan has been one of the most highly-advanced age societies in the world and more than 50% of OHCA patients are over 75 years of age. We have all data of elderly OHCA patients transported to the hospital because we don't have a system of termination of resuscitation.

We found CPC score results in elderly patients lower than younger patients even with bystander AED applications. We suggested that it is better to classify the age groups to analyze the effect of PAD in the advanced age society like Japan.

Results

	all (n=4571)	Age (years)				p
		15-64 (n=2101)	65-74 (n=925)	75-84 (n=801)	85-104 (n=744)	
Age, Mean (SD), y	65.6 (18.0)	50.0 (12.6)	69.4 (2.82)	79.5 (2.86)	90.3 (3.98)	
Females, n (%)	1194 (26.1)	249 (11.9)	164 (17.8)	293 (36.6)	488 (65.6)	<.0001
Cardiac origin, n (%)	3728 (81.6)	1879 (89.4)	809 (87.5)	590 (73.7)	450 (60.5)	<.0001
EMT Response time, mean (SD), min	9.51	9.64	9.59	9.32	9.23	
Resuscitation parameters						
Witnessed by Family	428 (9.4)	161 (7.7)	129 (13.9)	92 (11.5)	46 (6.2)	<.0001
Dispatcher instruction of CPR, n (%)	1673 (36.6)	696 (33.1)	299 (32.4)	341 (42.6)	337 (45.3)	<.0001
rescue breathing	2423 (53.2)	1054 (50.3)	386 (52.8)	435 (54.4)	448 (40.5)	<.0001
Chest compression	4348 (95.2)	1997 (95.1)	873 (94.6)	765 (95.6)	713 (95.8)	n.s.
defibrillation by EMT	1589 (34.8)	881 (42.0)	369 (39.9)	229 (28.6)	110 (14.8)	<.0001
Advanced airway by EMT	1354 (29.7)	541 (25.8)	267 (29.0)	271 (33.9)	275 (37.0)	<.0001
administered Epinephrine	692 (15.1)	288 (13.8)	152 (16.5)	137 (17.2)	115 (15.6)	n.s.
ROSC, n (%)	1984 (43.4)	1123 (53.4)	443 (47.9)	266 (33.2)	152 (20.4)	<.0001
30-day survival (%)	1708 (37.4)	1051 (50.0)	393 (42.4)	187 (23.3)	77 (10.3)	<.0001
Initial ECG when EMT arrived at the scene						
Shockable, n (%)	1256 (27.5)	704 (33.5)	306 (33.1)	169 (21.1)	77 (10.4)	
Non-shockable, n (%)	2107 (46.1)	677 (32.2)	355 (38.4)	490 (61.2)	585 (78.6)	
Assumed ROSC, n (%)	1208 (26.4)	720 (34.3)	264 (28.5)	142 (17.7)	82 (11.0)	
Neurological outcome, n (%)						
CPC1,2	1440 (31.5)	934 (44.5)	333 (36.0)	140 (17.5)	33 (4.4)	<.0001

Who was the bystander?

ECG wave when EMT arrived vs. Age group

*The shaded area stands for CPC 1 or 2.

In multivariate logistic regression analyses, in patients over 65 years of age, the following factors were associated with increased good neurologic outcome: younger age, non-medical intervention (advanced airway management, IV access, adrenaline administration) by EMT.

Contact

Previous Paper

KYOKO TSUKIGASE

Email: k-tsukigase@kokushikan.ac.jp
Research Institute of Disaster management and EMS
Kokushikan University

Our Website