Hypocalcaemia Following Laryngectomy

Research Question: what is the prevalence of hypocalcaemia after laryngectomy and is total thyroidectomy a risk factor?

Andrew S Harris Eduard Prades Carl D Passant Duncan R Ingrams
Aneurin Bevin University Health Board, The Royal Gwent Hospital, Newport

This is the largest study to ask this question and the first to follow patients up longer than three months.

Hypocalcaemia is a recognised complication of laryngectomy. However, there is a scarcity of research in this area. The prevalence and risk factors are unknown.

Postoperative hypocalcaemia causes extended admission (often to intensive care), requires long lines, blocks NG tubes, needs repeated blood tests and results in lifelong reliance on oral medication.

Post laryngectomy hypocalcaemia results in extra morbidity and increased health costs.

Consecutive total laryngectomies at The Royal Gwent Hospital identified from January 2006 to August 2017.

Electronic records retrospectively reviewed for operation notes, letters and investigation results.

Analysis of results for early (12-24 hours), protracted (4-6 weeks) and permanent hypocalcaemia (after 1 year).

Patient Demographics

	Thyroid in situ	Hemi thyroidectomy	Total thyroidectomy	Total
Total	14	69	7	90
Male				83%
Neck dissection	71%	99%	100%	92%
Salvage Procedure	71%	29%	14%	34%
Adjuvant Radiotherapy	29%	54%	86%	52%

Prevalence of hypocalcaemia

Exact logistic regression analysis shows that total thyroidectomy was significant at all time frames. Other variables did not have a significant effect.

	Number of patients	Odds ratio	95% CI	Model P-value
Early hypocalcaemia	90	15.5	2.2-181.9	0.002
Protracted hypocalcaemia	87	13.3	1.5-117.1	0.010
Permanent hypocalcaemia	60	22.7	1.9-376.5	0.005

Take home message

By choosing to remove more thyroid tissue during laryngectomy your patient is more likely to suffer early and long term hypocalcaemia.