

## Chapter

# Targeting the Aryl Hydrocarbon Receptor (AhR): A Review of the In-Silico Screening Approaches to Identify AhR Modulators

*Farag E.S. Mosa, Ayman O.S. El-Kadi and Khaled Barakat*

## Abstract

Aryl hydrocarbon receptor (AhR) is a biological sensor that integrates environmental, metabolic, and endogenous signals to control complex cellular responses in physiological and pathophysiological functions. The full-length AhR encompasses various domains, including a bHLH, a PAS A, a PAS B, and transactivation domains. With the exception of the PAS B and transactivation domains, the available 3D structures of AhR revealed structural details of its subdomains interactions as well as its interaction with other protein partners. Towards screening for novel AhR modulators homology modeling was employed to develop AhR-PAS B domain models. These models were validated using molecular dynamics simulations and binding site identification methods. Furthermore, docking of well-known AhR ligands assisted in confirming these binding pockets and discovering critical residues to host these ligands. In this context, virtual screening utilizing both ligand-based and structure-based methods screened large databases of small molecules to identify novel AhR agonists or antagonists and suggest hits from these screens for validation in an experimental biological test. Recently, machine-learning algorithms are being explored as a tool to enhance the screening process of AhR modulators and to minimize the errors associated with structure-based methods. This chapter reviews all *in silico* screening that were focused on identifying AhR modulators and discusses future perspectives towards this goal.

**Keywords:** human AhR, in silico, in vitro, AhR modulator, crystal structure, AhR modeling

## 1. Introduction

Six decades ago, researchers made extensive studies to answer a puzzling question. That was how administrating exogenous substances such as polycyclic aromatic hydrocarbons (PAHs) had a potent induction on xenobiotic-metabolizing enzymes in rats' livers [1, 2]. It was finally Alan Poland and his colleagues who finally answered this question in the early 1970s. Poland discovered a novel hepatic protein in complex with the polycyclic aromatic hydrocarbons compound, 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) [2]. The new protein was bound to TCDD in a potent affinity and was isolated from hepatic cytosolic fractions of mice C57BL/6, a mice model strain for studying aromatic hydrocarbon responsiveness.

This protein was later termed as the aryl hydrocarbon receptor (AhR) [2] and was identified as a ligand-activated transcription factor.

Later studies showed that AhR is expressed in several tissues including but not limited to; liver, lung, placenta, and heart and different cell types throughout the developmental periods of organ growth [3]. Further knockout studies in mice revealed essential functions for AhR in multiple physiological and pathophysiological pathways [4–6]. This accumulated knowledge over the last decades defined AhR as an environmental sensor for air pollutants and as a ligand-activated transcriptional factor, which regulates the expression of various genes, including enzymes responsible for xenobiotic metabolism [7].


AhR-mediates the toxicity of uncountable xenobiotics, and their triggered toxicity is accompanied by an overexpression and overactivation of AhR in cells. Thus, it increases the pathophysiological functions of AhR and could develop cancer in different organs such as the breast and liver. In addition, it can also lead to cardiovascular diseases, among other diseases [8, 9]. Thus, targeting AhR with a small molecule agonist/antagonist could efficiently inhibit several of the important hallmarks of various cancers [10].

Computational modeling and computer simulations continue to be an important tool for studying various biological mechanisms and for analyzing the interactions between biomolecular entities (*e.g.* proteins, DNA, and drugs). These methods use complex physiochemical and mathematical algorithms to predict the interactions between ligands and proteins at the atomistic level [11]. For instance, in the case of AhR, several computational studies have been reported to screen the chemical space for potential AhR modulators and to investigate how these ligands interact with the individual residues within AhR [12, 13]. This chapter will focus on summarizing the findings and computational approaches described in these studies. The chapter first discusses current knowledge related to AhR functions and describes all reported AhR crystal structures. It will then provide an overview on the different computational studies reported in the literature, which focused on investigating AhR dynamics and on identifying novel AhR modulators. We hope the information provided in this chapter can advance the development of AhR ligands and contribute to progress in this important research area.

## 2. AhR structure and functions

### 2.1 AhR domain structure

AhR is a member of the basic helix–loop–helix (bHLH)-PER- ARNT-SIM (PAS) family of transcription factors. The “PAS” term is an abbreviation for three proteins, namely, the *Drosophila* circadian rhythm protein period (Per), the mammalian AhR nuclear translocator (ARNT), and *Drosophila* neurogenic protein single-minded (Sim) [7, 14, 15]. Human AhR is a 848 amino acid with a molecular weight of ~96 kDa [16]. It includes two PAS domains, namely PAS A and PAS B, and interacts with the Aryl hydrocarbon nuclear translocator (ARNT) protein. Moreover, the PAS B domain involves two interaction sites: a ligand-binding site in which a bound ligand can modulate the AhR activity; and a direct binding interface for the HSP 90-chaperone protein. Additionally, AhR includes a basic helix loop helix motif located near its N-terminal domain, which is responsible for DNA binding as well as contributing to other protein–protein interactions. Finally, the transactivation (*i.e.*, glutamine-rich region) domain is located close to the C-terminal of AhR and binds to a co-activator [7] as shown in **Figure 1**.



**Figure 1.**  
*AhR domain structure and sub-domains' functions.*

## 2.2 The AhR ligands and their modes of action

The AhR PAS B domain can interact with both exogenous and endogenous chemicals from various origins. These interactions can induce different effects on AhR activity, leading to a wide range of physiological and toxicological downstream consequences. For example, several studies showed that environmental pollutants have been associated with developing cardiovascular diseases, cancer, and other diseases through AhR modulation [7, 17, 18]. Exogenous AhR ligands include various aromatic hydrocarbon molecules such as dioxins. One can be exposed to such ligands through contaminated food or environmental pollutants. Following exposure, their interaction with AhR can lead to several toxic effects, including organ dysfunctions, immunotoxicity, and carcinogenicity. On the other hand, endogenous AhR ligands are usually metabolic derivatives derived from cellular processes such as 6-Formylindolo (3,2- b) carbazole (FICZ). The interaction of these ligands with AhR is part of a normal functional response through AhR modulation [7, 19, 20].

## 2.3 AhR physiological and pathophysiological roles

AhR is an essential protein that contributes to countless biological pathways to establish its physiological role in developing the immune system and regulating xenobiotic enzymes [7, 15, 21]. AhR knockout mice models showed abnormal female reproductive functions and impairment in managing blood pressure [7]. The overactivation and constitutive activation of AhR have been associated with the initiation, promotion, progression, and invasion of cancer cells. For example, activating AhR by exogenous AhR ligands can have several effects, which includes inducing cell proliferation in the G1-S phase, silencing tumor suppressor genes, and activating proto-oncogenes in cancer cell lines.

Earlier findings showed that the exogenous AhR ligand, 2,3,7,8-tetrachlorodibenzo-p-dioxin TCDD promoted the degradation of cell-cell adhesion and expansion of cancer cells' motility by separating the Src kinase from the AhR protein complex. Furthermore, the activation of AhR via environmental pollutants can lead to a significant induction of xenobiotic-metabolizing enzymes, including CYP1A, which produces reactive intermediate metabolites and reactive oxygen species to promote tumor growth [14, 22]. In a nutshell, AhR resembles a machinery of genes, which controls xenobiotic-metabolizing enzymes in phases I and II, as shown in **Table 1**. Also, known AhR agonists such as TCDD and  $\beta$ -naphthoflavone have been shown to induce cellular hypertrophic actions on H9c2 cardiomyoblast cells. This was correlated with an increase in the levels of numerous cytochrome P450 genes, which could overcome by using an AhR antagonist [31].

On the positive side, experiments on a mouse model of induced colitis showed that the endogenous AhR agonist (FICZ), which has a strong binding affinity towards AhR, could block IL-6 and claudin-2 expression, and prevent any induced disorders in the intestinal barrier function through AhR activation [32].


Metabolism phase	Gene	Reference
Phase I	CYP1A1	[23]
Phase I	CYP1A2	[24]
Phase I	CYP1B1	[25]
Phase I	CYP2S1	[26]
Phase II	NQO1 NAD(P)H: Quinine oxidoreductase 1	[27]
Phase II	GSTA1 Glutathione transferase A1/2	[28]
Phase II	UGT1A6 Uridine diphosphate glucuronosyltransferase 1A6	[29]
Phase II	ALDH3A1 Aldehyde dehydrogenase 3A1	[30]

**Table 1.**  
*Xenobiotic metabolizing enzymes genes regulate via AhR pathway.*

Further protein knockout studies showed that AhR ligands play a fundamental role in autoimmune diseases through regulating Tregs and TH17 cell differentiation in the immune system. For example, FICZ inhibited Treg and TH17 cell development, accelerating experimental autoimmune encephalomyelitis in mice models [21, 33].

## 2.4 AhR signaling pathways

AhR is generally expressed in its inactive form in the cytoplasm as part of a protein complex encompassing a dimer heat shock protein, co-chaperone p23, an AhR-interacting protein, called AIP, and the protein kinase SRC (see **Figure 2**). The PAS



**Figure 2.**  
*Canonical pathway of aryl hydrocarbon receptor.*

B domain within AhR binds to one monomer of the HSP90 dimer and the second HSP90 monomer interacts with the AhR basic helix–loop–helix domain (bHLH) as well as with the PAS A domain [34]. As shown in **Figure 2**, the bHLH domain within AhR is also crucial for DNA binding in a process initiated by the binding of an AhR ligand within the PAS B domain and its interaction with the co-chaperone P23. Binding to P23 stabilizes AhR in the cytoplasm, protecting it from proteasomal degrading, and also maintains the PAS B domain of AhR in a unique conformation, suitable for strong ligand binding [35, 36].


Once an AhR ligand binds to the PAS B domain, it forms an AhR–ligand complex, including p23, SRC, and AIP (see **Figure 2**). This complex is transformed into an active state and then translocated inside the nucleus. Then in the nucleus, all complex components dissociate from the AhR–ligand complex, excluding an agonist and AhR protein. Subsequently, AhR forms an active heterodimer with ARNT and creates an AhR–ARNT complex. This complex is then recruited to the DNA via the Dioxin response element (DRE), exhibiting a common DNA compromise motif (5′-TNGCGTG-3′). This canonical AhR pathway increases the expression of various genes, including the principal ones in xenobiotic metabolism, AhR repressor (AHRR), and other genes [36].

### 3. AhR three-dimensional structures

Resolving the full-length three-dimensional structure of AhR has been a challenging exercise for the last two decades. Unfortunately, despite the many efforts towards this goal, there is no complete structure for the whole AhR protein. However, as discussed below, there are a few structures, which describe the number AhR domains. Although these structures do not reveal the exact overall AhR architecture, they can still provide useful information on the function of these separate domains. Giving computational modeling a favorable vantage point to construct reliable hypotheses for the full-length AhR organization for rational drug development and drug screening campaigns.

The first AHR 3D structure was reported in 2013 for the mouse PAS A domain (residues 110 to 267) at a resolution of 2.55 Å (PDB ID: 4M4X) (see **Figure 3**). This X-ray diffraction-based PAS A homodimer structure was obtained from recombinant *E. coli* expression. It contained a five  $\beta$ -sheet and elements order B-A-I-H-G, with four  $\alpha$ -helices ( $C\alpha$ ,  $D\alpha$ ,  $E\alpha$ , and  $F\alpha$ ) adjoining one side of the  $\beta$ -sheet. The interactions between the two PAS A monomers involved two distinctive areas within the A′  $\alpha$ -helices, revealing a strong interaction between Phe115, Leu116, and Ala119 from the A′  $\alpha$ -helix in one monomer with Val124, Phe260, and Ile262 from the  $\beta$ -sheet in the other monomer as shown in **Figure 3**. The protein structure revealed an undruggable pocket due to hydrophobic residues, and other residues such as Gln112 and Ile262 are essential in the interface for AHR dimerization, either homodimer or heterodimer with ARNT [37, 38].


Two more additional AhR structures were revealed in 2017 (see **Figure 4**). The two structures comprise multiple AhR domains and show a clear interaction between AhR and its dimerization partner, ARNT, as well as its interaction with two DNA strands. The two structures (PDB IDs: 5V0L and 5NJ8) [39, 40] were resolved at a resolution of 4.0 and 3.35 Å, respectively and revealed the complex formation among the bHLH and PAS A domains from human AhR and their interactions with ARNT and DNA. However, due to the observed high flexibility of the AhR PAS B domain and the transactive domain (C-terminal), none of these two subdomains were included in this architecture. However, both structures clearly explain the protein–protein interactions (PPI) and show clear interface regions for these



**Figure 3.** The upper figure represents crystal structure of homodimer mouse AhR-PAS a obtained from protein databank [37], and the lower figure shows the interactions between the two monomers.

interactions between the individual domains within AhR as well as their interactions with ARNET and DNA.

As shown in **Figure 4**, the first PPI interface is between the AhR-ARNT heterodimer with the two DNA strands. This interaction is mediated by DRE Ser36, His39, and Arg40 from the AhR bHLH domain and His79, Asp83, Arg86, and Arg87 from ARNT, as well as thymine and guanine from the DNA. The second PPI interface is between AhR and ARNT through different regions within the two proteins. These regions involve many hydrophobic interactions from both proteins and comprise residues Leu47, Leu50, Leu53, Val74, and Leu70 from the AhR bHLH domain and residues Ile109, Leu112, Val136, and Met139 in ARNT. The third PPI interface involves interactions between residues from the PAS A domain in both AhR and ARNT, mediated by residues Phe117, Leu118, Ala121, Leu122, Tyr137, Val126, Phe266, and Ile268 from AhR. The fourth, and final PPI interface encompasses the interdomain interactions between the AhR bHLH



**Figure 4.**  
*Crystal structure of human aryl hydrocarbon receptor in heterodimer with aryl hydrocarbon nuclear translocator and recruit on DNA in dioxin element response [39].*

and AhR PAS A domains, through residues Phe136, Ser151, Ile154, and Leu246 from the PAS A domain and Phe56, Val60, Leu72, Ala79, and Phe82 from the bHLH domain.

#### 4. Applications of computational methods in AhR modeling

The wealth of structural information described above on AhR provides an excellent opportunity to apply various computer-based simulations to study the dynamicity and structural organization of the various AhR domains. The applications of such computational tools not only can yield much needed insights on how these domains interact together within the AhR machinery, but can also offer detailed answers on their interactions with other AhR partners (*e.g.* ARNT, DNA, and chaperone proteins). It can also explain how a small molecule ligand can bind to AhR and how this can affect AhR functions, conformational dynamics or its interaction with other entities. Computational tools can also suggest novel-binding sites either within the AhR structure, or at the interfaces described above to either

stabilize these interaction (*i.e.* agonism) or block these interactions (*i.e.* antagonism). Most importantly, computational methods including virtual screening can be used as a high throughput-screening tool to identify compounds that can bind to these sites to modulate the AhR activity.

#### 4.1 Modeling the PAS B domain

Most of the *in silico* AhR screening campaigns till now focused on studying the PAS B domain. The PAS B domain is also known as the ligand-binding domain (LBD), where the ligands (agonists/antagonists) have been shown to bind [41]. Given that the available AhR structures (described above) lack this domain, computational methods played a key role in studying the interactions of ligands with this important region. Towards this goal, homology modeling was used to build 3-dimensional structures for this domain to allow the study of ligand binding to AhR. A homology model approach usually starts with identifying a similar template to the target domain (*i.e.*, the PAS B domain in this case). Once a template is identified, one uses various computational methods (*e.g.* sequence alignment, threading, and loop modeling) to construct the 3-dimensional structure of the target protein [42].

In many AhR studies, the human hypoxia inducible factors (HIF-2 $\alpha$ ) crystal structures served as templates for AhR-PAS B domain because it has the highest sequence similarity towards the AhR-PAS B domain. **Table 2** provides a list of the reported *in silico* studies that were conducted over the last few years by adopting various crystal structures of HIF-2 $\alpha$  as starting points to construct PAS B models. These studies were focused on understanding the roles played by the different PAS B residues in interacting with known AhR modulators and to screen for novel AhR ligands. Docking findings from these studies indicated that the binding cavity within the AhR-LBD can accommodate ligands with structural maximal dimensions of 14 Å X 12 Å X 5Å, and showed that their binding within AhR relies mainly on electronic properties [17]. Given this information, various computational methods including molecular modeling, molecular docking followed by MD simulations, and binding free energy calculations were used to provide insights about ligand interactions within the PAS B pockets [49].

For example, Bisson and his group established an agonist-optimized model of the human AhR-PAS B domain, followed by docking around five thousand chemical structures, including AhR agonists and antagonists, within the PAS B domain.

PDB ID	Structure method	Year of study	Reference
3H82	X-ray diffraction	2014	[41]
4GHI	X-ray diffraction	2014	[43]
3H3W	Electron microscopy	2016	[44]
3F1O, 3H7W, 3H82	X-ray diffraction	2018, 2018	[45, 46]
4XT2	X-ray diffraction	2019	[47]
3F1N, 3F1O, 3F1P, 3H7W, 3H82, 4GHI, 4GS9, 4XT2, 4ZP4, 4ZQD	X-ray diffraction	2019	[12]
3H82, 3H7W, 4ZQD	X-ray diffraction	2020	[48]

**Table 2.**

Report studies that used different crystal structures of human hypoxia inducible factors (HIF-2 $\alpha$ ).




Docking results were then filtered and the top five systems were subjected to long MD simulations (~ 60 ns) to study the conformational and dynamical changes in these generated complexes. Findings from Bisson's work revealed the importance of residues 307–329 in the PAS B domain, which were shown to be very flexible, acting as an access gate to the ligand-binding pocket. These residues can also adopt different conformations upon AhR ligands' binding and play a primary function in controlling the structural changes and accessibility of the ligands to the AhR ligand binding pocket [41].

#### 4.2 Interaction of the PAS B domain with different ligands

With the 3-dimensional structure of the PAS B domain in hand, many groups focused on studying its binding to different ligands (e.g., TCDD) (shown in **Figure 5**) [50] and investigated this binding reaction for different species. For example, TCDD studies on mouse AhR revealed a number of conserved residues that regulate the access of TCDD to the binding pocket [51–53]. These residues include Thr283, His285, Phe289, Tyr316, Ile319, Cys327, Met334, Phe345, Ala375, and Gln377 and have been also shown to control the internal size of the binding cavity [54–56]. Similarly, aromatic side chains of Phe 289, Phe 345, and Tyr 316 were shown to be important in stabilizing TCDD in its best mode of binding via non-covalent interaction [54].

Mutations at outer residues (e.g., Arg282, Thr311, Glu339, and Lys350) into alanine did not impact TCDD binding to AhR [57]. In the human AHR-LBD a mutation at Ala375 to Val and Leu decreases the binding affinity of TCDD and makes indirubin a less potent endogenous AhR ligand [45, 58]. Additional site-directed mutagenesis within AhR-LBD residues has been used to identify key residues promoting for ligand selectivity in AhR. These developed models provided a clear basis towards understanding the mechanism of ligand-dependent activation of AHR via its PAS B domain. In particular, the above mentioned molecular docking



**Figure 5.**  
Chemical structures of AhR ligands in this study obtained from pubchem database.

and mutagenesis analyses helped in identifying and confirming the binding pocket of TCDD and other AhR modulators [52, 57, 59, 60].

Examples of these models include those developed by Kim and her team, who constructed 3D models from several avian species including, chicken, albatross, and cormorant, and studied the sensitivity of dioxin derivatives against multiple AhR isoforms. All models were subjected to docking simulations with TCDD followed by MD simulations. Kim's results used the mean square displacement (MSD) of the MD trajectories as a stability indicator for the bound ligands. These findings revealed Ile324 and Ser380 from chicken AhR1 exhibited the least MSD values compared to all AhR-LBD residues in other avian species. The size of binding pocket was also shown to be variable among the different species. Moreover, stabilization of TCDD in the binding pocket of chicken AhR relied on the features of Ile324 and Ser380, which explained why chicken AhR is more sensitive to TCDD binding compared to other AhR isoforms [54, 61–63].

Further mutational and functional analysis studies were expanded to include additional AhR modulators other than TCDD. For example, the work of Faber and her team studied indirubin binding to AhR in both mouse and human. This study revealed that a mutation in His326Tyr and Ala349Thr in mouse AhR, and Tyr332 and Thr355 in human AhR can increase the potency of indole compounds, particularly, indirubin. Also, although indirubin and vemurafenib can fit within the same binding pocket in AhR, the two compounds showed two different modes of binding [45, 47]. For example, flutamide efficiently binds to residues inside the AHR-LBD with a high affinity in both mouse and human AHR to activate the AhR pathway [64]. It is important to note that, the biological response of AhR is dependant on the type of the bound ligand and has been shown to change based on the interaction of a given ligand with the residues forming the LBD in the PAS B domain [48, 65].

#### 4.3 Virtual screening and machine learning models applied to AhR

Over the last few decades, virtual screening has been used as a major tool to in hit identification campaigns against numerous biological targets [66]. In this regard, AhR is no exception and various *in silico* screening methods have been employed to identify new AhR modulators based on the developed 3D models for the PAS B domain [13]. These methods can be classified into two major groups; ligand-based methods (*e.g.*, quantitative structure–activity relationship (QSAR)) and structure-based methods (*e.g.*, docking and MD simulations). Ligand-based methods (LB) depend on the knowledge of known active/inactive molecules against a given target or disease to suggest new active chemical entities. Ligand-based methods are typically used when no information about the 3D structure of the target is available. They include QSAR, pharmacophore modeling and machine learning algorithms methods. QSAR models, for example, correlates the structural, physicochemical features, and biological mode of action of known compounds to build a mathematical model, which can be used to suggest new modifications to these structures for better activity or improved biophysical/biochemical properties [67–69].

Pharmacophore modeling maps the ligand–target interactions into a set of steric and electronic features structured in a specific 3D arrangement [70]. These pharmacophore models can be then used to screen millions of available chemical structural libraries for compounds that satisfy these pharmacophore features, which can be used for scaffold hopping and fragment-based drug design. On the other hand, structure–based methods require the knowledge of target protein crystal structure, or its 3D developed homology models. Ligands from a given database can be fitted into the active site of the target protein and can be ranked based on the predicted

binding affinities. In this context, molecular docking and molecular dynamics simulations are among the many valuable tools that can be used to predict the most probable mode of binding of a given ligand within the target. Furthermore, structure-based pharmacophore models can provide more detailed insights on the interaction of ligand with the binding site [69, 71, 72].

As discussed below, several AhR screening studies combined both methods to enhance the search for possible AhR candidates [67, 73]. The plethora of accumulated physicochemical, chemical and structural data on AhR modulators augmented this hit identification search with great tools to build reliable machine learning models, which require large datasets of chemical structures along with their interaction kinetics with AhR [74].

An example of AhR *in silico* screening studies is the one implemented by Xiao et al., who constructed 3D structures of the PAS B using the HIF-2 $\alpha$  as a template. Xiao used his model to study the effects of ~185 polybrominated diphenyl ethers (PBDEs), classified as organic pollutants, on AhR activation. This study combined molecular docking simulations, two-dimensional quantitative structure–activity relationship (2D-QSAR) models, and three-dimensional QSAR (3D-QSAR) models to analyze the local ligands’ interactions against a diverse set of PAS B configurations. Their result showed that bromide replacements in at the ortho- or meta-positions of PBDEs (BDE-49) as shown in the **Figure 5** exhibited the largest effect on PBDEs’ binding, mainly interacting with residues Met342, Thr290, Met334, and Phe289 in the binding site of the AHR-PAS B model in mouse and zebrafish [75, 76].

In a similar approach, Rath and his team built two human PAS B domain; a wild type and mutant (Val381 Ala, Val381Asn) models. Around 60 natural compounds from *Withania somnifera* were then docked within these models. Docking rustles were then refined using MD simulations for 50 ns. Findings from Rath’s study showed that withaferin A, withanolide A, withanolide B, withanolide D and withanone were effective as AhR ligands in all three models. In the meantime, withanolide A was more stable in the binding site and interacted with various residues in each model even after 50 ns of the MD simulations. Withanolide A was further validated experimentally in an *in vivo* zebrafish model to significantly reduce CYP1A1 expression. This was done in the presence of a strong AhR activator, namely benzo[a]pyrene in adult zebrafish brain when administrated together. Thus, withanolide A (see **Figure 5** and **Table 3**) neutralized the benzo[a]pyrene toxicity in zebrafish brain [77].

Compound name	Induction of AhR transcription	Binding free energy (kcal/mol)	Reference
Withanolide A	+	–7.5	[77]
Pinocembrin (5,7-Dihydroxyflavanone, R-form)	+	–2.9	[60]
5-hydroxy-7-methoxyflavone	+	–4.3	[60]
IMA-06201 (N-ethyl-N-phenyl-5-chloro-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxamide)	+	Not report	[46]
IMA-06504 (N-(4-trifluoromethylphenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide)	+	Not report	[46]

**Table 3.**  
 Activation of AhR transcription by chemical compounds that identified by *in silico* screening of different chemical libraries.

In another screening study, Mahiout, et al. identified IMA-06201 and IMA-06504 as two novel AhR agonists, with similar modes of binding to that of TCDD. Both compounds showed great stability in the central area of the AhR ligand-binding pocket. Furthermore, these AhR agonists were shown to be more efficient and more potent as selective AhR modulators than TCDD. To confirm that, Mahiout used CYP1A1 enzyme activity as a biomarker for AhR activation and compared the efficacy and potency of IMA-06201 and IMA-06504 (see **Figure 5** and **Table 3**) to that of TCDD in the presence and absence of the AhR antagonist, CH-223191, at different concentrations in rat hepatoma cell lines. Their results showed that the new compounds, IMA-06201 and IMA-06504, were able to induce CYP1A1 activity in a similar efficacy to that of TCDD, where CH-223191 was shown to block their CYP1A1 induction. Also, in an Ames test to assess the genotoxicity of the new identified compounds, IMA-06201 and IMA-06504 did not show mutagenic effects at low concentrations [46].

Machine-learning algorithms combined with QSAR have been recently used to screen for new AhR ligands. For instance, Matsuzaka used deep learning (DL) to construct machine-learning models to predict AhR activators. These models showed advantages on enhanced input data based on the 3D chemical structures of the compounds into these models, and their performance was better than traditional machine learning models [78]. To enhance the screening process of AhR ligands, Zhu established a virtual screening protocol from combining ligand-based and structure-based screening with supervised machine learning to screen around eight thousand from the pesticide databases to identify an agonistic effect on AHR activity. Zhu's results revealed sixteen compounds as AhR activators and these findings were validated in a zebrafish *in vivo* model to assess their AhR activation and exhibited induction in CYP 1a1 levels [79].

Towards improving the prediction accuracy of his model, Yang, et al. used machine learning algorithms to construct two-dimensional quantitative structure-activity relationship (2D-QSAR) models from multiple linear regression (MLR) and artificial neural network (ANN) algorithms. He used the pEC50 values of 60 dioxins derivatives as AhR activators to build. These models predicted the toxicity of 162 new dioxin derivatives, showing a good correlation between compounds' chemical structures and their IC50 and EC50 values.

Recently, Goya-Jorge employed various machine learning algorithms to build a set of QSAR models. These models adopted the adaboost (AdB), random forest (RF), gradient boosting (GB), support vector machine (SVM), and multilayer perceptron (MLP) as classifiers to examine around 1900 compounds from synthetic and natural sources on their AhR agonism. Around 40 compounds bearing the benzothiazole scaffold were classified as AhR agonists. In vitro validation of these hits showed that indole derivatives can serve as AhR ligands, including the endogenous substances [80, 81]. **Table 3** reports some of the top hits emerging from different *in silico* studies.

## 5. Current challenges in modeling AhR

Identifying novel AhR modulators using *in silico* approaches require establishing more comprehensive computational models of this target. These models should describe the detailed organization of the different AhR domains as well as its interaction with other protein/DNA partners. While the available crystal structures provide a glimpse of these missing pieces of information, there are still more to be done in this regard. For example, all currently available AhR crystal structures deposited in the protein data bank are lacking two important AhR domains, namely

the PAS B domain and the transactivation domain [39]. The transactivation domain is essential in AhR intercellular trafficking.

On the other hand, the PAS B domain interacts with an AhR ligand, which can modulate the AhR activity. While homology modeling has helped constructing acceptable models for this domain, the similarity of the templates used to build the PAS B domain is very low, leaving a lot of doubt about their accuracy. A crystal structure of the PAS B domain would be a great leap forward towards understanding the mode of action of AhR modulators and towards identifying better agonists/antagonists for this important target. Furthermore, there is a gap of knowledge on how AhR interact with other protein partners in the inactive state, including co-chaperone, AIP, and the protein kinase SRC. This builds an additional challenge to identify druggable pockets at their protein–protein interfaces [7, 82]. With the apparent advances in obtaining 3D experimental structures of protein (e.g. Cryo-electron microscopy (cryo-EM)) one expects several of these structural challenges can be solved in the near future, opening new gates for the computational science to identify new AhR modulators and to help understand its functional, structural and biological characterizes more clearly.

## 6. Executive summary


The AhR is a ligand-activated transcriptional factor. It regulates various genes' expression and plays a pathophysiological function in numerous diseases. Crystallography has been employed to resolve three crystal structures containing bHLH and PAS A domains from human and mouse origin and to identify four protein–protein interfaces. However, all these structures lacked the PAS B domain, which plays a fundamental role in ligands' binding domain to AhR. Computational and mutational studies revealed important residues that constitute the binding pockets within the PAS B domain. Towards identifying novel AhR modulators, several virtual screening and machine learning algorithms were constructed based on the available structural and pharmacological properties of known AhR ligands. Computational methods are extremely fast and intensely reduce the cost and time in screening millions of compounds to find compounds that could interact with the AhR. Recent studies employing these methods against AhR have been reviewed and discussed in this chapter. We hope the literature presented here can help advance the development of novel, selective and potent AhR modulators.

### Author details

Farag E.S. Mosa, Ayman O.S. El-Kadi\* and Khaled Barakat\*  
Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta,  
Edmonton, AB, Canada

\*Address all correspondence to: [kbarakat@ualberta.ca](mailto:kbarakat@ualberta.ca) and [aelkadi@ualberta.ca](mailto:aelkadi@ualberta.ca)

### IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] A.H. Conney, J.J. Burns, Stimulatory effect of foreign compounds on ascorbic acid biosynthesis and on drug-metabolizing enzymes, *Nature*. 184 (1959) 363-364. doi:10.1038/184363a0.
- [2] A. Poland, E. Glover, A.K.-J. of B. Chemistry, undefined 1976, Stereospecific, high affinity binding of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. Evidence that the binding species is receptor for induction of Elsevier. (n.d.). <https://www.sciencedirect.com/science/article/pii/S0021925817332052> (accessed January 24, 2021).
- [3] Y.Z. Jiang, K. Wang, R. Fang, J. Zheng, Expression of aryl hydrocarbon receptor in human placentas and fetal tissues, *J. Histochem. Cytochem.* 58 (2010) 679-685. doi:10.1369/jhc.2010.955955.
- [4] K.M. Burbach, A. Poland, C.A. Bradfield, Cloning of the Ah-receptor cDNA reveals a distinctive ligand-activated transcription factor., *Proc. Natl. Acad. Sci. U. S. A.* 89 (1992) 8185-8189. doi:10.1073/pnas.89.17.8185.
- [5] P. Fernandez-Salguero, T. Pineau, D.M. Hilbert, T. McPhail, S.S.T. Lee, S. Kimura, D.W. Nebert, S. Rudikoff, J.M. Ward, F.J. Gonzalez, Immune system impairment and hepatic fibrosis in mice lacking the dioxin-binding Ah receptor, *Science* (80-.). 268 (1995) 722-726. doi:10.1126/science.7732381.
- [6] K.W. Bock, Human and rodent aryl hydrocarbon receptor (AHR): From mediator of dioxin toxicity to physiologic AHR functions and therapeutic options, *Biol. Chem.* 398 (2017) 455-464. doi:10.1515/hsz-2016-0303.
- [7] L. Larigot, L. Juricek, J. Dairou, X. Coumoul, AhR signaling pathways and regulatory functions, *Biochim. Open.* 7 (2018) 1-9. doi:10.1016/j.biopen.2018.05.001.
- [8] I.A. Murray, A.D. Patterson, G.H. Perdew, Aryl hydrocarbon receptor ligands in cancer: Friend and foe, *Nat. Rev. Cancer.* 14 (2014) 801-814. doi:10.1038/nrc3846.
- [9] C. Esser, A. Rannug, The aryl hydrocarbon receptor in barrier organ physiology, immunology, and toxicology, *Pharmacol. Rev.* 67 (2015) 259-279. doi:10.1124/pr.114.009001.
- [10] S.K. Kolluri, U.H. Jin, S. Safe, Role of the aryl hydrocarbon receptor in carcinogenesis and potential as an anti-cancer drug target, *Arch. Toxicol.* 91 (2017) 2497-2513. doi:10.1007/s00204-017-1981-2.
- [11] A. Ganesan, K. Barakat, Applications of computer-aided approaches in the development of hepatitis C antiviral agents, *Expert Opin. Drug Discov.* 12 (2017) 407-425. doi:10.1080/17460441.2017.1291628.
- [12] S. Giani Tagliabue, S.C. Faber, S. Motta, M.S. Denison, L. Bonati, Modeling the binding of diverse ligands within the Ah receptor ligand binding domain, *Sci. Rep.* 9 (2019) 1-14. doi:10.1038/s41598-019-47138-z.
- [13] X. Lin, X. Li, X. Lin, A review on applications of computational methods in drug screening and design, *Molecules.* 25 (2020). doi:10.3390/molecules25061375.
- [14] J.P. Whitlock, Induction of cytochrome P4501A1, *Annu. Rev. Pharmacol. Toxicol.* 39 (1999) 103-125. doi:10.1146/annurev.pharmtox.39.1.103.
- [15] T. V. Beischlag, J.L. Morales, B.D. Hollingshead, G.H. Perdew, The aryl hydrocarbon receptor complex and the control of gene expression, *Crit. Rev.*

- Eukaryot. Gene Expr. 18 (2008) 207-250. doi:10.1615/  
CritRevEukarGeneExprv18.i3.20.
- [16] C. Rowlands, J.C. Rowlandsar, J.-A. Gustafssonbj, Aryl Hydrocarbon Receptor-Mediated Signal Transduction, Aryl Hydrocarbon Receptor-Mediated Signal Transduction Artic. Crit. Rev. Toxicol. 27 (1997) 109-110.  
doi:10.3109/10408449709021615.
- [17] J. Abel, T. Haarmann-Stemann, An introduction to the molecular basics of aryl hydrocarbon receptor biology, Biol. Chem. 391 (2010) 1235-1248.  
doi:10.1515/BC.2010.128.
- [18] E.M. Reyes-Reyes, I.N. Ramos, M.A. Tavera-Garcia, K.S. Ramos, The aryl hydrocarbon receptor agonist benzo(a) pyrene reactivates LINE-1 in HepG2 cells through canonical TGF- $\beta$ 1 signaling: Implications in hepatocellular carcinogenesis, Am. J. Cancer Res. 6 (2016) 1066-1077.
- [19] O. Ibraheem, T.H. Fatoki, J.M. Enibukun, B.C. Faleye, D.U. Momodu, In silico toxicological analyzes of selected toxic compounds from dumpsite or contaminated soils on human health, Nov. Biotechnol. Chim. 18 (2019) 144-153. doi:10.2478/nbec-2019-0017.
- [20] M. Adam, L. Chaubah, B.W. Bontes, N.A. Mulachelah, J. Veteran, M. East, J. Indonesia, Molecular Docking of Polycyclic Aromatic Hydrocarbons as Potentially Carcinogenic Molecules Through Binding with Aryl Hydrocarbon Receptor, J. Smart Bioprospecting Technol. P-ISSN. 01 (2019) 1. <https://doi.org/10.21776/ub.jsmartech.2019.001.01>.
- [21] F.J. Quintana, A.S. Basso, A.H. Iglesias, T. Korn, M.F. Farez, E. Bettelli, M. Caccamo, M. Oukka, H.L. Weiner, Control of Treg and TH17 cell differentiation by the aryl hydrocarbon receptor, Nature. 453 (2008) 65-71.  
doi:10.1038/nature06880.
- [22] P. Xue, J. Fu, Y. Zhou, The aryl hydrocarbon receptor and tumor immunity, Front. Immunol. 9 (2018) 1.  
doi:10.3389/fimmu.2018.00286.
- [23] F. Saatcioglu, D.J. Perry, D.S. Pasco, J.B. Fagan, Multiple DNA-binding factors interact with overlapping specificities at the aryl hydrocarbon response element of the cytochrome P450IA1 gene., Mol. Cell. Biol. 10 (1990) 6408-6416. doi:10.1128/mcb.10.12.6408.
- [24] W. Li, P.A. Harper, B.K. Tang, A.B. Okey, Regulation of cytochrome P450 enzymes by aryl hydrocarbon receptor in human cells. CYP1A2 expression in the LS180 colon carcinoma cell line after treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin or 3-methylcholanthrene, Biochem. Pharmacol. 56 (1998) 599-612.  
doi:10.1016/S0006-2952(98)00208-1.
- [25] L. Zhang, Ü. Savas, D.L. Alexander, C.R. Jefcoate, Characterization of the mouse Cyp1B1 gene: Identification of an enhancer region that directs aryl hydrocarbon receptor-mediated constitutive and induced expression: J. Biol. Chem. 273 (1998) 5174-5183.  
doi:10.1074/jbc.273.9.5174.
- [26] S.P. Rivera, S.T. Saarikoski, O. Hankinson, Identification of a novel dioxin-inducible cytochrome P450, Mol. Pharmacol. 61 (2002) 255-259.  
doi:10.1124/mol.61.2.255.
- [27] L. V. Favreau, C.B. Pickett, Transcriptional regulation of the rat NAD(P)H:Quinone reductase gene. Characterization of a DNA-protein interaction at the antioxidant responsive element and induction by 12-O-tetradecanoylphorbol 13-acetate, J. Biol. Chem. 268 (1993) 19875-19881.  
doi:10.1016/s0021-9258(19)36594-9.
- [28] M. Rojas, I. Cascorbi, K. Alexandrov, E. Kriek, G. Auburtin, L. Mayer, A. Kopp-Schneider, I. Roots, Modulation of benzo[a]pyrene

dioloxide-DNA adduct levels in human white blood cells by CYP1A1, GSTM1 and GSTT1 polymorphism, 2000.

[29] M.F. Yueh, Y.H. Huang, A. Hiller, S. Chen, N. Nguyen, R.H. Tukey, Involvement of the xenobiotic response element (XRE) in Ah receptor-mediated induction of human UDP-glucuronosyltransferase 1A1, *J. Biol. Chem.* 278 (2003) 15001-15006. doi:10.1074/jbc.M300645200.

[30] W.T. Kays, J. Piatigorsky, Aldehyde dehydrogenase class 3 expression: Identification of a cornea-preferred gene promoter in transgenic mice, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 13594-13599. doi:10.1073/pnas.94.25.13594.

[31] B.N.M. Zordoky, A.O.S. El-Kadi, 2, 3,7,8-Tetrachlorodibenzo-p-dioxin and  $\beta$ -naphthoflavone induce cellular hypertrophy in H9c2 cells by an aryl hydrocarbon receptor-dependant mechanism, *Toxicol. Vitro* 24 (2010) 863-871. doi:10.1016/j.tiv.2009.12.002.

[32] Y. Ma, Q. Wang, K. Yu, X. Fan, W. Xiao, Y. Cai, P. Xu, M. Yu, H. Yang, 6-Formylindolo(3,2-b)carbazole induced aryl hydrocarbon receptor activation prevents intestinal barrier dysfunction through regulation of claudin-2 expression, *Chem. Biol. Interact.* 288 (2018) 83-90. <https://doi.org/10.1016/j.cbi.2018.04.020>.

[33] B. Tong, X. Yuan, Y. Dou, X. Wu, G. Chou, Z. Wang, Y. Xia, Y. Dai, Norisoboldine, an isoquinoline alkaloid, acts as an aryl hydrocarbon receptor ligand to induce intestinal Treg cells and thereby attenuate arthritis, *Int. J. Biochem. Cell Biol.* 75 (2016) 63-73. doi:10.1016/j.biocel.2016.03.014.

[34] B.N. Fukunaga, M.R. Probst, S. Reisz-Porszasz, O. Hankinson, Identification of functional domains of the aryl hydrocarbon receptor, *J. Biol.*

*Chem.* 270 (1995) 29270-29278. doi:10.1074/jbc.270.49.29270.

[35] I. Pongratz, G.G.F. Mason, L. Poellinger, Dual roles of the 90-kDa heat shock protein hsp90 in modulating functional activities of the dioxin receptor. Evidence that the dioxin receptor functionally belongs to a subclass of nuclear receptors which require hsp90 both for ligand binding activity and repression of intrinsic DNA binding activity, *J. Biol. Chem.* 267 (1992) 13728-13734. doi:10.1016/s0021-9258(18)42274-0.

[36] V. Rothhammer, F.J. Quintana, The aryl hydrocarbon receptor: an environmental sensor integrating immune responses in health and disease, *Nat. Rev. Immunol.* 19 (2019) 184-197. doi:10.1038/s41577-019-0125-8.

[37] D. Wu, N. Potluri, Y. Kim, F. Rastinejad, Structure and Dimerization Properties of the Aryl Hydrocarbon Receptor PAS-A Domain, *Mol. Cell. Biol.* 33 (2013) 4346-4356. doi:10.1128/mcb.00698-13.

[38] J. V. De Souza, S. Reznikov, R. Zhu, A.K. Bronowska, Druggability assessment of mammalian Per-Arnt-Sim [PAS] domains using computational approaches, *Medchemcomm.* 10 (2019) 1126-1137. doi:10.1039/c9md00148d.

[39] K.W. Schulte, E. Green, A. Wilz, M. Platten, O. Daumke, Structural Basis for Aryl Hydrocarbon Receptor-Mediated Gene Activation, *Structure.* 25 (2017) 1025-1033.e3. doi:10.1016/j.str.2017.05.008.

[40] S.H. Seok, W. Lee, L. Jiang, K. Molugu, A. Zheng, Y. Li, S. Park, C.A. Bradfield, Y. Xing, Structural hierarchy controlling dimerization and target DNA recognition in the AHR transcriptional complex, *Proc. Natl. Acad. Sci. U. S. A.* 114 (2017) 5431-5436. doi:10.1073/pnas.1617035114.



- [41] A. Perkins, J. Phillips, N. Kerkvliet, R. Tanguay, G. Perdew, S. Kolluri, W. Bisson, A Structural Switch between Agonist and Antagonist Bound Conformations for a Ligand-Optimized Model of the Human Aryl Hydrocarbon Receptor Ligand Binding Domain, *Biology (Basel)*. 3 (2014) 645-669. doi:10.3390/biology3040645.
- [42] H. Jalily Hasani, K. Hassan Barakat, H. Jalily Hasani, K. Barakat, Homology Modeling: an Overview of Fundamentals and Tools Article in International Review on Modelling and Simulations · Computational drug design and molecular dynamics View project Interferon lambda signaling in the context of influenza vaccination View project Homology Modeling: an Overview of Fundamentals and Tools *Int. Rev. Model. Simulations*. 10 (2017). <https://doi.org/10.15866/iremos.v10i2.11412>.
- [43] A.J. Parks, M.P. Pollastri, M.E. Hahn, E.A. Stanford, O. Novikov, D.G. Franks, S.E. Haigh, S. Narasimhan, T.D. Ashton, T.G. Hopper, D. Kozakov, D. Beglov, S. Vajda, J.J. Schlezinger, D.H. Sherr, In silico identification of an aryl hydrocarbon receptor antagonist with biological activity in vitro and in vivo, *Mol. Pharmacol.* 86 (2014) 593-608. doi:10.1124/mol.114.093369.
- [44] I.S. Kim, J.H. Hwang, M. Hirano, H. Iwata, E.Y. Kim, In vitro and in silico evaluation of transactivation potencies of avian AHR1 and AHR2 by endogenous ligands: Implications for the physiological role of avian AHR2, *Comp. Biochem. Physiol. Part - C Toxicol. Pharmacol.* 187 (2016) 1-9. doi:10.1016/j.cbpc.2016.03.011.
- [45] S. Faber, A. Soshilov, S. Giani Tagliabue, L. Bonati, M. Denison, Comparative In Vitro and In Silico Analysis of the Selectivity of Indirubin as a Human Ah Receptor Agonist, *Int. J. Mol. Sci.* 19 (2018) 2692. doi:10.3390/ijms19092692.
- [46] S. Mahiout, S.G. Tagliabue, A. Nasri, I.M. Omoruyi, L. Pettersson, L. Bonati, R. Pohjanvirta, In vitro toxicity and in silico docking analysis of two novel selective AH-receptor modulators, *Toxicol. Vitro*. 52 (2018) 178-188. doi:10.1016/j.tiv.2018.06.010.
- [47] H.C. Hawerkamp, A. Kislat, P.A. Gerber, M. Pollet, K.M. Rolfes, A.A. Soshilov, M.S. Denison, A.A. Momin, S.T. Arold, A. Datsi, S.A. Braun, P. Oláh, M.E. Lacouture, J. Krutmann, T. Haarmann-Stemann, B. Homey, S. Meller, Vemurafenib acts as an aryl hydrocarbon receptor antagonist: Implications for inflammatory cutaneous adverse events, *Allergy*. 74 (2019) 2437-2448. doi:10.1111/all.13972.
- [48] J.H. Hwang, K. Kannan, T.J. Evans, H. Iwata, E.Y. Kim, Assessment of Risks of Dioxins for Aryl Hydrocarbon Receptor-Mediated Effects in Polar Bear (*Ursus maritimus*) by in Vitro and in Silico Approaches, *Environ. Sci. Technol.* 54 (2020) 1770-1781. doi:10.1021/acs.est.9b05941.
- [49] K.N. Chitralla, X. Yang, P. Nagarkatti, M. Nagarkatti, Comparative analysis of interactions between aryl hydrocarbon receptor ligand binding domain with its ligands: A computational study, *BMC Struct. Biol.* 18 (2018) 1-17. doi:10.1186/s12900-018-0095-2.
- [50] P. Ramadoss, G.H. Perdew, Use of 2-azido-3-[125I]iodo-7,8-dibromodibenzo-p-dioxin as a probe to determine the relative ligand affinity of human versus mouse aryl hydrocarbon receptor in cultured cells, *Mol. Pharmacol.* 66 (2004) 129-136. doi:10.1124/mol.66.1.129.
- [51] C. Odio, S.A. Holzman, M.S. Denison, D. Fraccalvieri, L. Bonati, D.G. Franks, M.E. Hahn, W.H. Powell, Specific ligand binding domain residues confer low dioxin responsiveness to AHR1 $\beta$  of *xenopus laevis*, *Biochemistry*.

52 (2013) 1746-1754. doi:10.1021/bi301722k.

[52] A. Pandini, A.A. Soshilov, Y. Song, J. Zhao, L. Bonati, M.S. Denison, Detection of the TCDD binding-fingerprint within the Ah receptor ligand binding domain by structurally driven mutagenesis and functional analysis, *Biochemistry*. 48 (2009) 5972-5983. doi:10.1021/bi900259z.

[53] M.S. Denison, A.A. Soshilov, G. He, D.E. Degroot, B. Zhao, Exactly the same but different: Promiscuity and diversity in the molecular mechanisms of action of the aryl hydrocarbon (dioxin) receptor, *Toxicol. Sci.* 124 (2011) 1-22. doi:10.1093/toxsci/kfr218.

[54] I. Motto, A. Bordogna, A.A. Soshilov, M.S. Denison, L. Bonati, New aryl hydrocarbon receptor homology model targeted to improve docking reliability, *J. Chem. Inf. Model.* 51 (2011) 2868-2881. doi:10.1021/ci2001617.

[55] S. Miyagi, K. Murata, K. Sashino, S. Sawamura, S. Uruno, S. Yoshimura, E. Akahoshi, M. Ishihara-Sugano, S. Itoh, N. Kurita, Binding affinity between AhR and exogenous/endogenous ligands: Molecular simulations and biological experiment, *Mol. Simul.* 41 (2015) 555-563. doi:10.1080/08927022.2014.899696.

[56] M. Kumar Gadhwal, S. Patil, P. D'mello, A. Joshi, Homology modeling of aryl hydrocarbon receptor and docking of agonists and antagonists, n.d.

[57] A. Pandini, M.S. Denison, Y. Song, A.A. Soshilov, L. Bonati, Structural and functional characterization of the aryl hydrocarbon receptor ligand binding domain by homology modeling and mutational analysis, *Biochemistry*. 46 (2007) 696-708. doi:10.1021/bi061460t.

[58] A.A. Soshilov, M.S. Denison, Ligand Promiscuity of Aryl Hydrocarbon

Receptor Agonists and Antagonists Revealed by Site-Directed Mutagenesis, *Mol. Cell. Biol.* 34 (2014) 1707-1719. doi:10.1128/mcb.01183-13.

[59] C. Knebel, J. Kebben, I. Eberini, L. Palazzolo, H.S. Hammer, R.D. Süßmuth, T. Heise, S. Hessel-Pras, A. Lampen, A. Braeuning, P. Marx-Stoelting, Propiconazole is an activator of AHR and causes concentration additive effects with an established AHR ligand, *Arch. Toxicol.* 92 (2018) 3471-3486. doi:10.1007/s00204-018-2321-x.

[60] W.H. Bisson, D.C. Koch, E.F. O'Donnell, S.M. Khalil, N.I. Kerkvliet, R.L. Tanguay, R. Abagyan, S.K. Kolluri, Modeling of the aryl hydrocarbon receptor (AhR) ligand binding domain and its utility in virtual ligand screening to predict new AhR ligands, *J. Med. Chem.* 52 (2009) 5635-5641. doi:10.1021/jm900199u.

[61] J.-H. Hwang, S. Bak, E.-Y. Kim, In Silico Analysis of the Interaction of Avian Aryl Hydrocarbon Receptors and Dioxins to Decipher Isoform-, Ligand-, and Species-Specific Activations, *Artic. Environ. Sci. Technol.* 49 (2015) 3795-3804. doi:10.1021/es505733f.

[62] B. Wu, Y. Zhang, J. Kong, X. Zhang, S. Cheng, In silico predication of nuclear hormone receptors for organic pollutants by homology modeling and molecular docking, *Toxicol. Lett.* 191 (2009) 69-73. doi:10.1016/j.toxlet.2009.08.005.

[63] Ž. Jakopin, Assessment of the endocrine-disrupting potential of halogenated parabens: An in silico approach, *Chemosphere.* 264 (2021) 128447. doi:10.1016/j.chemosphere.2020.128447.

[64] X. Gao, C. Xie, Y. Wang, Y. Luo, T. Yagai, D. Sun, X. Qin, K.W. Krausz, F.J. Gonzalez, The antiandrogen flutamide is a novel aryl hydrocarbon receptor ligand that disrupts bile acid

- homeostasis in mice through induction of Abcc 4, *Biochem. Pharmacol.* 119 (2016) 93-104. doi:10.1016/j.bcp.2016.08.021.
- [65] D. Corrada, A.A. Soshilov, M.S. Denison, L. Bonati, Deciphering Dimerization Modes of PAS Domains: Computational and Experimental Analyses of the AhR:ARNT Complex Reveal New Insights Into the Mechanisms of AhR Transformation, *PLOS Comput. Biol.* 12 (2016) e1004981. doi:10.1371/journal.pcbi.1004981.
- [66] T. Zhu, S. Cao, P.C. Su, R. Patel, D. Shah, H.B. Chokshi, R. Szukala, M.E. Johnson, K.E. Hevener, Hit identification and optimization in virtual screening: Practical recommendations based on a critical literature analysis, *J. Med. Chem.* 56 (2013) 6560-6572. doi:10.1021/jm301916b.
- [67] J. Vázquez, M. López, E. Gibert, E. Herrero, F. Javier Luque, Merging ligand-based and structure-based methods in drug discovery: an overview of combined virtual screening approaches, *Molecules.* 25 (2020) 4723. doi:10.3390/molecules25204723.
- [68] C. Acharya, A. Coop, J. E. Polli, A. D. Mac Kerell, Recent Advances in Ligand-Based Drug Design: Relevance and Utility of the Conformationally Sampled Pharmacophore Approach, *Curr. Comput. Aided-Drug Des.* 7 (2010) 10-22. doi:10.2174/157340911793743547.
- [69] S.U. Khan, N. Ahemad, L.H. Chuah, R. Naidu, T.T. Htar, Sequential ligand- and structure-based virtual screening approach for the identification of potential G protein-coupled estrogen receptor-1 (GPER-1) modulators, *RSC Adv.* 9 (2019) 2525-2538. doi:10.1039/c8ra09318k.
- [70] X. Qing, X.Y. Lee, J. De Raeymaeker, J.R. Tame, K.Y. Zhang, M. De Maeyer, A.R. Voet, Video abstract Pharmacophore modeling: advances, limitations, and current utility in drug discovery, *J. Receptor. Ligand Channel Res.* 16 (2021). doi:10.2147/JRLCR.S46843.
- [71] Z. Wang, H. Sun, C. Shen, X. Hu, J. Gao, D. Li, D. Cao, T. Hou, Combined strategies in structure-based virtual screening, *Phys. Chem. Chem. Phys.* 22 (2020) 3149-3159. doi:10.1039/c9cp06303j.
- [72] M. Wieder, A. Garon, U. Perricone, S. Boresch, T. Seidel, A.M. Almerico, T. Langer, Common Hits Approach: Combining Pharmacophore Modeling and Molecular Dynamics Simulations, *J. Chem. Inf. Model.* 57 (2017) 365-385. doi:10.1021/acs.jcim.6b00674.
- [73] S. Debnath, T. Debnath, S. Bhaumik, S. Majumdar, A.M. Kalle, V. Aparna, Discovery of novel potential selective HDAC8 inhibitors by combine ligand-based, structure-based virtual screening and in-vitro biological evaluation, *Sci. Rep.* 9 (2019) 1-14. doi:10.1038/s41598-019-53376-y.
- [74] V.K. Gupta, P.S. Rana, Toxicity prediction of small drug molecules of aryl hydrocarbon receptor using a proposed ensemble model, *Turkish J. Electr. Eng. Comput. Sci.* 27 (2019) 2833-2849. doi:10.3906/elk-1809-9.
- [75] H. Xiao, N. Mei, Q. Chi, X. Wang, Comprehensive binding analysis of polybrominated diphenyl ethers and aryl hydrocarbon receptor via an integrated molecular modeling approach, *Chemosphere.* 262 (2021) 128356. doi:10.1016/j.chemosphere.2020.128356.
- [76] A.L.W. Po, W. Zhang, What lessons can be learnt from withdrawal of mibefradil from the market?, *Lancet.* 351 (1998) 1829-1830. doi:10.1016/S0140-6736(05)78800-0.

[77] S.N. Rath, L. Jena, M. Patri, Understanding ligands driven mechanism of wild and mutant aryl hydrocarbon receptor in presence of phytochemicals combating Parkinson's disease: an in silico and in vivo study, *J. Biomol. Struct. Dyn.* 38 (2020) 807-826. doi:10.1080/07391102.2019.1590240.

[78] Y. Matsuzaka, T. Hosaka, A. Ogaito, K. Yoshinari, Y. Uesawa, Prediction Model of Aryl Hydrocarbon Receptor Activation by a Novel QSAR Approach, Deep Snap-Deep Learning, *Molecules*. 25 (2020) 1317. doi:10.3390/molecules25061317.

[79] K. Zhu, C. Shen, C. Tang, Y. Zhou, C. He, Z. Zuo, Improvement in the screening performance of potential aryl hydrocarbon receptor ligands by using supervised machine learning, *Chemosphere*. 265 (2021) 129099. doi:10.1016/j.chemosphere.2020.129099.

[80] E. Goya-Jorge, R.M. Giner, M. Sylla-Iyarreta Veitía, R. Gozalbes, S.J. Barigye, Predictive modeling of aryl hydrocarbon receptor (AhR) agonism, *Chemosphere*. 256 (2020) 127068. doi:10.1016/j.chemosphere.2020.127068.

[81] Z. Ek Dvo R Ak, K. Poulíkov, S. Mani, Indole scaffolds as a promising class of the aryl hydrocarbon receptor ligands, (2021). doi:10.1016/j.ejmech.2021.113231.

[82] A. Tkachenko, F. Henkler, J. Brinkmann, J. Sowada, D. Genkinger, C. Kern, T. Tralau, A. Luch, The Q-rich/PST domain of the AHR regulates both ligand-induced nuclear transport and nucleocytoplasmic shuttling, *Sci. Rep.* 6 (2016) 1-11. doi:10.1038/srep32009.