CASE SERIES OF LIVER ABSCESES IN CHILDREN

D. Dharmapalan¹, A. Bagde¹, N. Manwani¹, S. Patil¹, G. Sharma¹, N. Meshram¹, R. Verma¹, P. Patil¹, H. Lahoti², V. Yewale¹.

¹Apollo hospitals- Navi Mumbai, Department of Pediatrics, Navi Mumbai, India.

²Apollo hospitals- Navi Mumbai, Department of Pediatric Surgery, Navi Mumbai, India

Background

Liver abscesses in children pose both a diagnostic and therapeutic challenge in tropical countries. A case series of 4 liver abscesses managed between March 2017-18 is presented with retrospective analysis of clinical, microbiological, radiological characteristics and therapeutic methods with clinical outcome.

Case series Summary

Of the 4 children diagnosed with liver abscess, all were males with mean age of 10.8 years. Fever was present in all, loose motions in two, one presented with right shoulder pain. All had tender right hypochondrium. Leucocytosis above 15000/cmm was present in 3 of 4 children. Deranged coagulopathy was noted in one. One child had micro abscesses involving both lobes. Rest had right lobe involvement (segment 6 being the commonest followed by segment 7). Pleural effusion was noted in two patients. Aspiration of abscess was required in two patients. Organism could be detected only in one. There was no bacteremia in any.

One child was initially treated as amoebic abscess on sonography finding suggestive of amoebic abscess and positive serology for *E. histolytica* antibodies but had to be restarted on antibiotics due to poor clinical response after 10 days, Fig 1. No organism was detected.

Another child presented with fever and weight loss, was on antitubercular treatment for pulmonary Tb since 5 months. Amoebic abscess was suspected due to concomitant history of amoebic abscess in father but antibodies to *E. histolytica* was negative, Fig 2. The abscess improved with Metronidazole and Diloxanide furoate.

Mean defervescence of fever was 10.75 days, mean total antibiotic duration was 17.5 days. The commonest antibiotic combination used was Co-amoxiclav plus Metronidazole plus Amikacin.

High degree of suspicion is required in to diagnose liver abscesses due to non-specific complaints. Despite microbiological and radiological facilities, it is challenging to differentiate pyogenic from amoebic liver abscesses.

Fig 1: Xray chest in a child who was readmitted with pleural effusion within 2 weeks of primary treatment as amoebic abscess, (had positive *E. histolytica* serology and normal Xray Chest on prior admission), there was mild resolution of abscess and required antibacterial treatment.

Fig 2A

Fig 2A: Xray Chest in a child on antitubercular treatment since 5 months for pulmonary TB, presented with fever since 10 days and weight loss.

Fig 2B

Fig 2B: CT scan showed liver abscess. Had recent history of father diagnosed with amoebic abscess. (child had negative *E. histolytica* serology). Was managed successfully as amoebic liver abscess.