Meal-sequence and postprandial glucose excursion: A time course study to find optimal component intervals in a healthy Japanese diet

¹SAKI OKAMOTO, ^{1,2}HITOSHI KUWATA, ²DAISUKE YABE, ^{1,2}YOSHIYUKI HAMAMOTO, ^{1,2}TAKESHI KUROSE AND ^{1,2}YUTAKA SEINO

¹CENTER FOR DIABETES, ENDOCRINOLOGY AND METABOLISM, KANSAI ELECTRIC POWER HOSPITAL, OSAKA, JAPAN.

²YUTAKA SEINO DISTINGUISHED CENTER FOR DIABETES RESEARCH, KANSAI ELECTRIC POWER MEDICAL RESEARCH INSTITUTE, KOBE, JAPAN;

Backgrounds and aims

The postprandial glucose concentration is influenced by a number of factors including meal size and composition, gastric emptying and intestinal glucose absorption rates, and the secretion and action of insulin, glucagon and the incretins. We previously established that eating fish or meat 15 minutes before rice enhances GLP-1 secretion and improves postprandial glucose excursion in Japanese individuals with type 2 diabetes as well as healthy individuals. However, the 15 minute-interval is not practical in everyday meals. In this study, we attempted to clarify the optimal time interval between eating non-carbohydrate food and rice to minimize postprandial glucose elevation.

Meal-sequence (with 15-min interval)		→	→
Incretin	GLP-1	↑↑	↑↑
	GIP	↑	↑↑
Gastric emptying		delayed	delayed
Postprandial glucose elevation		Improved	Improved

Preload of fish or meat enhances incretin secretion; delays gastric emptying and ameliorates postprandial glucose elevation. Caution is required for meat preload as GIP in collaboration with high fats may cause obesity.

Kuwata H, Yabe D, Seino Y et al, Diabetologia 2016

Methods and subjects

Participants: This was a single centre, randomized, controlled trial involving fifteen healthy volunteers (n=15, age 31.0±4.5 years, HbA1c 5.9±0.2%). All participants were without diabetes, gastrointestinal tract disease including gastroparesis, history of gastrointestinal operation, cardiac disease, pulmonary disease, pancreatic disease, liver disease, renal disease, alcohol or drug abuse, glucose-lowering medication, diabetogenic medication or malignancy, or pregnancy and thyroid diseases.

Meal-sequence test: The total energy of the test meal was 550±36 kcal, protein; 16.5% , fat; 23.5% , carbohydrate; 60%.

Monitoring: Interstitial fluid glucose levels were measured every 15 min by continuous glucose monitoring (flush CGM; Abbott® Real time monitoring).

Statistical analysis: Results are reported as mean±SEM. Data analysis was performed with ANOVA. P value <0.05 was taken to indicate significant differences.

Results

Preloading of other foods of a meal before rice (-15 min, -10 min, and -5 min) resulted in significant improvement in postprandial glucose elevation compared to consuming both components at the same time (0 min) or consuming the rice before the other foods (+15 min). We also found that there was no significant difference after between preloading at -5, -10 or -15 minutes before rice. Each value was indicates as average ± SEM; * indicates p<0.05 versus + 15 min.

Conclusion

We show in this study that preloading of other foods before rice results in more favourable post prandial glucose excursion. We also find that such meal-sequencing shows beneficial effects with as little as 5 min other foods preloading. The current findings show the effectiveness of meal-sequencing in medical nutritional therapy in control postprandial glucose elevations.

COI

A position of a board member or advisor: None
Honoraria for lectures: AstraZeneca K.K., ARKRAY, Inc., Kao Corporation, Sanofi K.K., Taisho Toyama Pharmaceutical Co., Ltd., Takeda Pharmaceutical Company Limited., Novo Nordisk Pharma Ltd., MSD K.K.
Clinical commissioned / joint research grant: Boehringer Ingelheim Japan, Inc., Eli Lilly and Company, MSD K.K., Taisho Toyama Pharmaceutical Co., TERUMO CORPORATION
Scholarship grant: ARKRAY, Inc., ONO PHARMACEUTICAL CO., LTD., Taisho Toyama Pharmaceutical Co., Ltd., Takeda Pharmaceutical Company Limited., Novo Nordisk Pharma Ltd.