Pharmacokinetics and Pharmacodynamics of Mycophenolate in Patients After Renal Transplantation

Thomas Rath¹ and Manfred Küpper²

¹*Department of Nephrology and Transplantation Medicine,
Westfalz-Klinikum GmbH, Kaiserslautern*

²*HPLC-Laboratory, Institute for Immunology and Genetics, Kaiserslautern
Germany*

1. Introduction

Mycophenolic-acid (MPA) is a selective, non-competitive inhibitor of Inosine-Monophosphate-Dyhydrogenase (IMPDH) leading to the inhibition of the de-novo synthesis of guanosine-nucleotides. In human lymphocytes inhibition of IMPDH results in altered cellular proliferation with arrest in the S-phase of the cell cycle. Due to the absence of a salvage pathway, proliferating activated t-cells are severely affected by the inhibitory effects of MPA (1-3). For patients after renal transplantation MPA is used either as mycophenolate-mofetil (MMF, Cellcept) or as enteric-coated mycophenolate-Sodium (EC-MPS, Myfortic) in daily doses of 2000 mg respectively 1440 mg per day.

Since its introduction in immunosuppressive therapy more than ten years ago, Mycophenolate-Mofetil (MMF) is an established part of immunosuppressive therapy after renal transplantation. Still in the first publication of the landmark Tricontinental trial because of possibly dose-related side effects of the drug (CMV-infection, gastrointestinal disturbances, and increased cancer risk) the need for individualization depending on clinical course or other factors was mentioned (4).

The usefulness of pharmacokinetic measurements of MMF was shown in early studies stating that the Area-under the curve (AUC) of MMF is predictive of the likelihood of allograft rejection after renal transplantation in patients receiving mycophenolate mofetil (5). To facilitate therapeutic drug monitoring different limited sampling strategies for adult and pediatric patients after renal transplantation were established (6-13).

The two available preparations of MPA (MMF, EC-MPS) showed equivalent drug exposure measured by MPA-AUC when applied to the patients in equimolar doses. Therefore, both preparations are seen as equipotent (14-16).

2. Pharmacokinetics of MPA

MPA trough levels show relevant inter- and intraindividual variability especially in patients with elevated serum creatinine and proteinuria (17-19). Clinically important, low trough

levels are associated with an increased frequency of rejection (20), whereas elevated MPA trough levels are related to an increased risk for infections (21). Nevertheless, relevant correlations between MPA trough levels and MPA-AUC values could not be detected, therefore the usefulness of measuring trough levels in routine care of renal transplant recipients is doubted (22-24).

2.1 Effect of immunosuppressive therapy on MPA pharmacokinetics

Concomitant immunosuppressive therapy has major influence on MPA pharmacokinetics. For patients on Cyclosporine (CsA) therapy lower MPA trough levels are observed (25). In addition, MPA trough levels increased after discontinuation of CsA resulting in almost a doubling of MPA trough concentrations (26). In general, the variability of MPA-AUC in patients with concomitant cyclosporine and steroid therapy seems to be low (27). However, in 154 patients with an immunosuppressive therapy consisting of CsA, prednisone and MMF the mean MPA AUC increased after 21 days although mean MMF dose was reduced (28).

For patients treated with tacrolimus (TAC) increased MPA trough levels are reported (29). Additionally, a randomized trial with 150 participants, patients receiving TAC and MMF displayed significantly higher MPA trough levels and higher MPA exposure measured by MPA-AUC than those receiving CsA and the same dose of MMF. Equivalent MPA levels could only be attained in patients receiving CsA by increasing the MMF dose by 50% (30). Similar results were obtained for pediatric renal transplantation (31). Interestingly, at least in Japanese renal transplant patients, a difference for MPA-AUC in patients with different tacrolimus-trough levels could not be detected (32). At least, for renal transplant recipients limited sample strategies for MPA-AUC with concomitant medication of tacrolimus are established (33).

In patients with Sirolimus (SRL) MPA exposure in the presence of SRL is higher than MPA exposure with CsA. Therefore it was recommended, that the MMF dose should be reduced to 0.75 g twice a day in patients receiving SRL to obtain MPA-AUC levels comparable to that in patients treated with CsA and MMF 1 g twice a day (8). These results were confirmed in a pharmacokinetic study in 31 renal transplant patients (34). It was also shown, that although MPA peak concentration and time to peak concentration was comparable, the MPA-AUC was higher in patients receiving SRL instead of CsA (35).

Steroids have been shown to induce the hepatic glucuronyltransferase (GT) expression enhancing the activity of uridine diphosphate-GT, the enzyme responsible for mycophenolic acid (MPA) metabolism. Therefore, also for steroids interactions with MPA are reported. During a steroid tapering and withdrawal phase in 26 patients MPA trough levels progressively increased and plasma MPA clearance declined (36).

2.2 Effect of concomitant therapy on MPA pharmacokinetics

For patients in the maintenance phase after transplantation it is known, that MPA-AUC increases with declining transplant function (37). This effect may be modulated by concomitant medication. Beside immunosuppression, patients after renal transplantation have to use antiviral prophylaxis. At least for Ganciclovir no effect on MPA clearance in kidney transplant recipients was reported (38).

With respect to the use of proton pump inhibitors the published results are unequivocally. In Japanese patients, the peak MPA-concentrations were lower with 30 mg lansoprazole

than with 10 mg rabeprazole or without PPI. For patients with cytochrome (CYP) 2C19, and multidrug resistance (MDR)1 C3435T polymorphisms this was also seen for the MPA-AUC (39).

Patients after heart transplantation with PPI co-medication show significantly lower MPA plasma concentrations resulting in lower drug exposure exposing the patients at a higher risk for acute rejection (40). Also in patients with autoimmune diseases the co-medication of pantoprazole with MMF significantly influences the drug exposure and immunosuppressive potency of MMF (41). In contrast, the recently published sub-analysis of the CLEAR-study reported no difference in MPA-AUC in patients with or without PPI-therapy when a 3g/d loading dose of MMF for 5 days used. However, MPA concentrations 2 h and 12 h after MMF intake were reduced (42).

At least for heart transplant recipients no influence of pantoprazole on EC-MPS pharmacokinetics could be disclosed (43).

Own results in 74 patients in the early and maintenance phase after renal transplantation showed a relevant reduction in normalized MPA-AUC (40,9 +/- 19,7 vs. 26,1 +/- 11,7 mg/l*h; $p < 0,01$) in patients with PPI co-medication. A difference between patients using either omeprazole or pantoprazole in MPA-AUC could not be detected. (Rath et al., Congress of the German Transplantation Society, 2009)

3. Clinical relevance of MPA-AUC

3.1 MPA-AUC and acute rejection

In different clinical trials, MPA drug exposure was correlated with the occurrence of biopsy proven acute rejection (BPAR). In a double blind trial aiming for three predefined target MPA AUC values the incidence of BPAR was lower in patients with MPA AUC values between 30 and 60 $\mu\text{g} \times \text{h}/\text{ml}$ (28). Similar results were reported for a group of 46 stable patients after renal transplantation, with better graft function in patients with a MPA AUC > 40 $\mu\text{g}/\text{ml} \times \text{h}$ and for pediatric renal transplantation (20;44;45).

Three randomized trials, the OPTICEPT study, the APOMYGRE-trial (Adaption de Posologie du MMF en Greffe Renale) and the FDCC study (fixed-dose versus concentration controlled) investigated the benefit of therapeutic drug monitoring for MMF in renal transplant recipients.

The APOMYGRE Trial was a study in 137 allograft recipients treated with basiliximab, cyclosporine A, corticosteroids and MMF. Patients were randomized to receive either concentration-controlled doses or fixed-dose MMF. A novel Bayesian estimator of MPA AUC based on three-point sampling was used to individualize MMF doses. At month 12, the concentration-controlled group had fewer treatment failures and acute rejection episodes. Therefore, the authors conclude, that therapeutic MPA monitoring using a limited sampling strategy can reduce the risk of treatment failure and acute rejection in renal allograft recipients 12 months post-transplant with no increase in adverse events (46).

The FDCC study was a randomized trial in 901 patients after renal transplantation allocating patients to receive MMF either in a fix dose or in a concentration controlled manner aiming at a predefined MPA AUC of 45 $\text{mg} \times \text{h}/\text{L}$. In general, there was no difference in the incidence of primary treatment failure or biopsy proved rejection. However, MPA-AUC levels at day 3 after transplantation predicts the incidence of BPAR in the first year (47).

The OPTICEPT study was a 2-year, open-label, randomized, multicenter trial comparing the efficacy and safety of concentration-controlled MMF dosing with a fixed-dose regimen in 720 kidney recipients. In patients with Tacrolimus, those with higher MMF exposure had less rejection episodes (48).

Similarly, a recently published substudy of the FDCC-trial in patients with delayed graft function disclosed significantly lower dose-corrected MPA AUC on Day 3 and Day 10 in this patient group (49).

3.2 MPA-pharmacokinetics and gastro-intestinal side effects

It is known, that side effects of MMF are causing dose reductions in approximately 60% of the patients leading to a cumulative and increasing risk for acute rejection (50). In addition, gastrointestinal (GIT) side effects affect medical adherence of the patients with consecutive risk for graft failure (51). In addition, dose reductions of MMF are related to increased costs, mainly due to frequent hospitalization of the patients (52).

USRDS data of 3589 patients with MMF prescription and GIT complaints revealed that dosage reduction or discontinuation of mycophenolate mofetil in the first 6 months after diagnosis of GI complications was associated with significantly increased risk of graft failure and increased healthcare costs in adult renal transplant recipients (53). Another report from USRDS data of 3675 patients with gastrointestinal complications under MMF and subsequent dose reduction also disclosed an increased risk for graft loss after dose reduction or discontinuation of MMF (54).

The enteric coated preparation of mycophenolate (EC-MPS) is attributed to a lower rate of gastrointestinal side effects, but in a prospective study based on patient questionnaires the rate of gastrointestinal side effects was nearly identical between the two formulations (55). In addition, a double-blind study comparing MMF and the newly developed enteric-coated formulation of MPA (EC-MPS) showed no advantage for either of the drugs (56). In contrast, a large, prospective study in more than 700 renal transplant recipients disclosed a significant improvement in gastrointestinal adverse events after conversion from MMF to EC-MPS (57). A study in patients with GIT complaints under MMF switching to EC-MPS indicates that converting patients with mild, moderate or severe GI complaints from MMF to EC-MPS significantly reduces GI-related symptom burden and improves patient functioning and well-being (58).

Also in liver transplant patients results are reported that converting patients with gastrointestinal complaints from MMF to equimolar doses of EC-MMF leads to a reduction of gastrointestinal-related symptom burden and frequency of stools (59).

There is some evidence from pharmacokinetic studies that elevated MPA exposure correlates with the occurrence of gastrointestinal side effects. Some authors suggest that gastrointestinal side effects are related to exposure of the active substance MPA (60). Others report, that the occurrence of possibly MMF-related side effects corresponds with MPA-AUC and MPA concentration 30 minutes after oral dose of 1000 mg (61). Also, a longitudinal study in 37 patients with 357 MPA measurements revealed higher trough levels in patients with MMF associated side effects (62). It is known that MPA trough levels >3 mg/l, peak levels >8.09 mg/l and MPA-AUC >37.6 mg*h/l may lead to adverse effects (63). Also in 31 patients after renal transplantation higher MPA-AUC (>60 mg*h/l) was

associated with side effects (64). Nevertheless, in a small pharmacokinetic study with 11 hispanic renal transplant patients treated with EC-MPS the MPA-AUC does not correlate with overall Gastrointestinal Symptom Rating Scale scores or subscale scores (65).

Also, a 5-year clinical follow-up study in 100 renal allograft recipients in whom MPA exposure was measured at 7 days, 6 weeks, 3 months, 1, 3, and 5 years post transplantation using abbreviated AUC measurements reported more episodes of leucopenia and anemia with MPA AUC(0-12h) ranges $>60 \text{ mg/L} \times \text{h}(-1)$. However, no association between incident episodes of diarrhea or infection and target MPA AUC (0-12 h) ranges (66).

4. Pharmacodynamics of MPA

4.1 Inhibition of IMPDH-activity

Recently, pharmacodynamic measurement of MPA was introduced into clinical practice. Especially with the use of reversed-phase HPLC, it is possible to monitor the immunosuppressive effect of MPA in its target cell population by quantifying the activity of IMPDH. This nonradioactive method for specific measurement of IMPDH activity in isolated peripheral mononuclear cells was developed by direct chromatographic determination of produced xanthosine 5'-monophosphate (XMP). In the canine model MPA in therapeutic doses leads to an 50% inhibition of IMPDH-activity (67). Application of a single dose of 1 g MMF in dialysis patients resulted in a significant inhibition of IMPDH activity in lysed mononuclear cells. IMPDH activity is inversely correlated to MPA blood concentrations and the IC (50) for in vitro inhibition of IMPDH activity was about 2 to 3 $\mu\text{g/L}$. (68). In addition, others report, that IMPDH-activity on peak concentration of MPA is approximately 40% and could be suppressed for 8 hours (69). In general, it is assumed, that IMPDH activity has a substantial interindividual, but low intraindividual variability (70). This was also shown in pediatric patients (71).

In addition, in renal allograft recipients an inverse relationship between plasma MPA and IMPDH activity within the dose interval was demonstrated and minimum IMPDH activity was a median 8 % of values pre-MMF dose, coinciding with the MPA peak. Six hours post-dose, IMPDH activity had returned to pre-dose values. Patients receiving MMF had a 4.5-fold higher pre-dose enzyme activity than transplanted patients without MMF (72). Long-term treatment with mycophenolate was associated with an induction of IMPDH activity (73). Also a study with 12 patients over two years showed an increase of type 1 IMPDH mRNA during the first 3 months following transplantation and reaching its maximal level during acute rejection episodes, whereas type2 IMPDH mRNA was stable (74). Interestingly, in 30 patients transplantation and the initiation of immunosuppressive therapy was associated with increased IMPDH1 and decreased IMPDH2 expression. In addition, patients with acute rejection during follow-up demonstrated higher IMPDH2 expression in pretransplant CD4+ cells than nonrejecting patients (75). Later, the same group described in detail the MMF concentration dependent modulation of IMPDH1 expression in renal allograft recipients (76).

4.2 IMPDH-activity and acute rejection

Measurement of IMPDH activity may be useful in estimating the degree of immunosuppression in individual patients in addition to applied MMF dose. When

comparing three patients groups with MMF doses of 1.0, 1.5 and 2.0 g/d there was no correlation between MPA-AUC (0-12) values and MMF dose detectable. Also, the degree of inhibition of IMPDH activity was comparable in the three groups, indicating considerable interindividual pharmacodynamic variability (77). In a cross-sectional analysis patients experiencing acute rejection episodes had increased IMPDH activity during rejection episodes (78). Additional information was gained in a genotyping study in 191 kidney transplant patients. There, seventeen genetic variants were identified in the IMPDH1 gene with allele frequencies ranging from 0.2 to 42.7%. Two single-nucleotide polymorphisms, rs2278293 and rs2278294, were significantly associated with the incidence of biopsy-proven acute rejection in the first year post-transplantation (79). A similar study in 82 Japanese transplant recipients found no difference in the incidence of subclinical acute rejection between IMPDH1 rs2278293 or rs2278294 polymorphisms ($p = 0.243$ and 0.735 , respectively). However, the authors report that the risk of subclinical acute rejection for recipients who cannot adapt in therapeutic drug monitoring (TDM) of MPA seems to be influenced by IMPDH1 rs2278293 polymorphism (80). Also, in patients after renal transplantation high pre-transplant IMPDH-activity predisposes to subsequent MPA dose-reductions and increases the risk for acute rejection (81).

4.3 IMPDH activity and MMF or EC-MPS

There is some discussion about the degree of IMPDH suppression with either MMF or EC-MPS. In a single-center, crossover study in patients treated with MMF and EC-MPS IMPDH activity inversely followed MPA concentrations and was inhibited to a similar degree (approximately 85%) by both formulations. In addition, the calculated value for 50% IMPDH inhibition was identical for both drugs (16). However, when comparing the pharmacodynamic activity of MMF and EC-MPS a series of 260 measurements in 110 patients disclosed lower median IMPDH activity in the EC-MPS patients than in the MMF patients. This was especially pronounced in patients on 1440 mg/d EC-MPS compared with 2000 mg/d MMF (82).

Nevertheless, for EC-MPS a recently published pharmacokinetic study in 75 de-novo kidney transplant recipients randomly assigning the patients either to receive EC-MPS as standard dose or as intensified dose revealed in an exploratory analysis of IMPDH activity that the intensified regimen resulted in significantly lower IMPDH activity on day 3 after transplantation (83).

There is ongoing discussion about the effect of PPI therapy on pharmacokinetics of MMF and EC-MPS. In a cross-sectional analysis in 153 renal transplant recipients, we measured IMPDH-activity before the first daily dose of MMF or EC-MPS. We could not detect any statistical with respect to PPI intake, type or dosing of either MMF or EC-MPS (Congress of the German Transplant Society, 2009).

5. Measuring IMPDH-activity

5.1 Sample preparation

Peripheral blood is collected in 5ml tubes with Li-heparin as anti-coagulant and stored at room temperature. Heparin is superior to EDTA as anti-coagulant since it maintains cell viability for longer time. Within four hours after arrival of the sample to the lab, and within

no more than two days of collection, the peripheral mononuclear cell fraction is isolated by density centrifugation according to a modified protocol from Glander et al. (2001). Li-heparinized blood (2.5ml) is mixed with an equal volume of phosphate-buffered saline (PBS), carefully layered on 4ml Lymphodex (InnoTrain, Germany) density gradient centrifugation medium in a 15ml screw-cap polypropylene tube, and centrifuged at $1200 \times g$ for 15 min without brake at room temperature.

The mononuclear cell fraction is collected from the interphase and transferred into a fresh 15ml screw-cap tube with 5ml PBS for washing. The cells are washed only once with PBS since repeated washing steps might cause diffusion of mycophenolate from the cells, resulting in over-estimation of the residual IMPDH activity. After centrifugation at $1200 \times g$ for 10 min at room temperature, the supernatant is removed quantitatively. This step is crucial with respect to the assay validity, since only a minute fraction of the total mycophenolate is contained within the cells, while the vast majority (estimated 99%) is present in the plasma. Any trace of the supernatant might therefore still contain considerable amounts of mycophenolate, hence leading to a vast underestimation of the residual IMPDH activity. The cell pellet is resuspended in 250 μ l ice-cold HPLC-grade water, and 125 μ l of the sample are transferred into each of two 2ml screw-cap vials, one designated as working sample, the second as back-up. The vials are deep frozen at -80°C until assayed. In the same way, control cells from healthy probands are prepared; these cells will be included in each assay as an incubation control.

5.2 IMPDH activity assay

The residual IMPDH activity is assayed in a cell-free system. The patient samples and control cells are thawed at room temperature and vigorously vortexed for 30 seconds to support cell lysis; insoluble cell fragments are removed by centrifugation at $4000 \times g$ for 5 min at room temperature in a desktop centrifuge. Cell lysate (50 μ l) is added to 100 μ l incubation buffer containing 1 mmol/L inosine-monophosphate (IMP) as substrate, 0.5 mmol/L NAD as co-substrate, 72 mmol/L sodium dihydrogen-phosphate, and 180 mmol/L potassium chloride (pH = 7.5). After adjusting the volume to 180 μ l with distilled water, the samples are incubated at 37°C in a heating block. In presence of NAD, IMPDH converts inosine-monophosphate to xanthine 5'-mono-phosphate. In the subsequent high-performance liquid chromatography (HPLC) assay, the amount of synthesized xanthine 5'-monophosphate is determined together with the amount of AMP, which serves as an internal standard for normalization to the cell count.

After exactly 2.5 hours of incubation, the reaction is stopped by adding 20 μ l ice-cold 4mol/L perchloric acid. Precipitation of denatured protein is enhanced by incubating the samples at -20°C for 10 min. After centrifugation at 13000 rpm for 2 min in a desktop centrifuge, 170 μ l supernatant are transferred to a test tube containing 14 μ l 2.5 mol/L potassium carbonate solution for neutralization. The exact volume of potassium carbonate, required to achieve a final pH between pH 6 and pH 7, has to be determined for each lot of 4 mol/L perchloric acid and 2.5 mol/L potassium carbonate solution. Prior to HPLC analysis, the samples are deep-frozen at -20°C for at least 30 min, thawed, and centrifuged 5 min at 13000 rpm in a desktop centrifuge.

5.3 HPLC chromatography

Determination of the amounts of xanthine-monophosphate and adenosine-monophosphate is carried out by ion-pair reversed-phase high-performance liquid chromatography on a computerized isocratic HPLC system from Shimadzu (Kyoto, Japan) consisting of a system controller SCL-10A VP, an HPLC pump LC-10AT VP, an autoinjector SIL-10AF, a column oven CTO-10AS VP, and an UV-VIS detector SPD-10A VP, controlled by Shimadzu LC Solution data collection software.

For the assay 6 μ L of the samples are loaded onto a 250 mm x 3.1 mm Prontosil 120 to 5 ODS AQ column (Bischoff Chromatography, Leonberg, Germany). Column oven temperature is set to 40°C. Chromatographic separation is achieved using a mobile phase containing 50 mmol/L potassium-dihydrogen-phosphate, 7 mmol/L tetra-n-butyl-ammonium hydrogen sulfate, and 6% (v/v) methanol at a flow rate of 1 mL/min. The analytes are detected at 254-nm wavelength. Incubation efficacy is verified by including a sample from a healthy volunteer as incubation control in each incubation cycle. For calibration, two standards containing 500 and 2500 pmol xanthin-monophosphate and adenosin-monophosphate, respectively, in 0.4% BSA solution are processed in several independent experiments, and repeatedly measured, like the patient specimen: protein denaturation with perchloric acid followed by neutralization with potassium carbonate. This calibration curve allows to deduct the amount of XMP synthesized during incubation and the amount of AMP in the sample. The specific IMPDH activity is then expressed as pmol XMP synthesized per second, which is normalized to 1 pmol of AMP [pmol XMP/(pmol AMP s)].

6. Summary and conclusion

Mycophenolic-acid (MPA) is a selective, non-competitive inhibitor of Inosine-Monophosphate-Dyhydrogenase (IMPDH) leading to the inhibition of the de-novo synthesis of guanosine-nucleotides. In human lymphocytes inhibition of IMPDH results in altered cellular proliferation with arrest in the S-phase of the cell cycle. Due to the absence of a salvage pathway, proliferating activated t-cells are severely affected by the inhibitory effects of MPA.

In patients after renal transplantation, MPA is a well-established part of immunosuppressive therapy, applied either as mycophenolate-mofetil (MMF, Cellcept) or as mycophenolate-sodium (MPS, Myfortic). MMF is used in prophylaxis of kidney rejection for nearly 15 years in daily doses of 2 – 3 g/d. The enteric-coated MPS is available since a few years; the recommended daily dose is 1440 mg/d. Both preparations are equipotent, when given in equimolar doses.

In recent years drug monitoring of MPA gained more and more attention proving its usefulness in clinical setting. Relevant information could be collected by measuring MPA drug exposure by calculating the MPA-Area under the curve (MPA-AUC) with pharmacokinetic modeling allowing estimating the degree of immunosuppression.

In different clinical studies MPA-AUC target concentrations of 30 – 60 μ g*h/ml were correlated to a low rate of rejections and less occurrence of drug induced side effects. Clinically important, MPA metabolism is influenced not only by the choice of immunosuppressive medication, but also by renal function and concomitant medication.

Recently pharmacodynamic measurement of MPA was introduced into clinical practice. Especially with the use of reversed-phase HPLC, it becomes possible to monitor the immunosuppressive effect of MPA in its target cell population by quantifying the activity of IMPDH. IMPDH activity is inversely correlated to MPA blood concentrations. Maximum inhibition of IMPDH activity ranges between 60% and 80% and the reported IC₅₀ of IMPDH activity corresponds to MPA blood levels of 2-3 μ /l. In renal transplant recipients, IMPDH shows relevant inter-individual variability. However, pre-transplant IMPDH activity was predictive for increased risk of rejection when additional dose reductions of MMF were necessary. In a cross-sectional studies better transplant function was associated with lower IMPDH-activity and probably the usage of EC-MPS. Pharmacokinetic and pharmacodynamic parameters of MPA are influenced by additional immunosuppression. In addition, concomitant therapy especially the use of proton-pump inhibitors affects MPA-levels, whereas an effect of IMPDH-activity, at least in renal transplant recipients could not be disclosed. Therefore, it can be concluded, that pharmacokinetic and pharmacodynamic measurements of MPA adds relevant information to improve clinical care of renal transplant recipients.

7. References

- [1] Allison AC, Eugui EM. Purine metabolism and immunosuppressive effects of mycophenolate mofetil (MMF). *Clin Transplant* 1996 Feb;10(1 Pt 2):77-84.
- [2] Cohn RG, Mirkovich A, Dunlap B, Burton P, Chiu SH, Eugui E, et al. Mycophenolic acid increases apoptosis, lysosomes and lipid droplets in human lymphoid and monocytic cell lines. *Transplantation* 1999 Aug 15;68(3):411-8.
- [3] Dayton JS, Lindsten T, Thompson CB, Mitchell BS. Effects of human T lymphocyte activation on inosine monophosphate dehydrogenase expression. *J Immunol* 1994 Feb 1;152(3):984-91.
- [4] A blinded, randomized clinical trial of mycophenolate mofetil for the prevention of acute rejection in cadaveric renal transplantation. The Tricontinental Mycophenolate Mofetil Renal Transplantation Study Group. *Transplantation* 1996 Apr 15;61(7):1029-37.
- [5] Hale MD, Nicholls AJ, Bullingham RE, Hene R, Hoitsma A, Squifflet JP, et al. The pharmacokinetic-pharmacodynamic relationship for mycophenolate mofetil in renal transplantation. *Clin Pharmacol Ther* 1998 Dec;64(6):672-83.
- [6] Johnson AG, Rigby RJ, Taylor PJ, Jones CE, Allen J, Franzen K, et al. The kinetics of mycophenolic acid and its glucuronide metabolite in adult kidney transplant recipients. *Clin Pharmacol Ther* 1999 Nov;66(5):492-500.
- [7] Willis C, Taylor PJ, Salm P, Tett SE, Pillans PI. Evaluation of limited sampling strategies for estimation of 12-hour mycophenolic acid area under the plasma concentration-time curve in adult renal transplant patients. *Ther Drug Monit* 2000 Oct;22(5):549-54.
- [8] El HW, Ficheux M, Debruyne D, Rognant N, Lobbedez T, Allard C, et al. Pharmacokinetics of mycophenolic acid in kidney transplant patients receiving sirolimus versus cyclosporine. *Transplant Proc* 2005 Mar;37(2):864-6.

- [9] Miura M, Satoh S, Niioka T, Kagaya H, Saito M, Hayakari M, et al. Limited sampling strategy for simultaneous estimation of the area under the concentration-time curve of tacrolimus and mycophenolic acid in adult renal transplant recipients. *Ther Drug Monit* 2008 Feb;30(1):52-9.
- [10] Mohammadpour AH, Nazemian F, Abtahi B, Naghibi M, Gholami K, Rezaee S, et al. Estimation of abbreviated mycophenolic acid area under the concentration-time curve during early posttransplant period by limited sampling strategy. *Transplant Proc* 2008 Dec;40(10):3668-72.
- [11] Filler G. Abbreviated mycophenolic acid AUC from C0, C1, C2, and C4 is preferable in children after renal transplantation on mycophenolate mofetil and tacrolimus therapy. *Transpl Int* 2004 Mar;17(3):120-5.
- [12] Schutz E, Armstrong VW, Shipkova M, Weber L, Niedmann PD, Lammersdorf T, et al. Limited sampling strategy for the determination of mycophenolic acid area under the curve in pediatric kidney recipients. German Study Group on MMF Therapy in Pediatric Renal Transplant Recipients. *Transplant Proc* 1998 Jun;30(4):1182-4.
- [13] Weber LT, Schutz E, Lamersdorf T, Shipkova M, Niedmann PD, Oellerich M, et al. Therapeutic drug monitoring of total and free mycophenolic acid (MPA) and limited sampling strategy for determination of MPA-AUC in paediatric renal transplant recipients. The German Study Group on Mycophenolate Mofetil (MMF) Therapy. *Nephrol Dial Transplant* 1999;14 Suppl 4:34-5.
- [14] Arns W, Breuer S, Choudhury S, Taccard G, Lee J, Binder V, et al. Enteric-coated mycophenolate sodium delivers bioequivalent MPA exposure compared with mycophenolate mofetil. *Clin Transplant* 2005 Apr;19(2):199-206.
- [15] Arns W, Breuer S, Choudhury S, Taccard G, Lee J, Binder V, et al. Enteric-coated mycophenolate sodium delivers bioequivalent MPA exposure compared with mycophenolate mofetil. *Clin Transplant* 2005 Apr;19(2):199-206.
- [16] Budde K, Bauer S, Hambach P, Hahn U, Roblitz H, Mai I, et al. Pharmacokinetic and pharmacodynamic comparison of enteric-coated mycophenolate sodium and mycophenolate mofetil in maintenance renal transplant patients. *Am J Transplant* 2007 Apr;7(4):888-98.
- [17] Merkel U, Lindner S, Vollandt R, Sperschneider H, Balogh A. Trough levels of mycophenolic acid and its glucuronidated metabolite in renal transplant recipients. *Int J Clin Pharmacol Ther* 2005 Aug;43(8):379-88.
- [18] Fernandez A, Marcen R, Pascual J, Martins J, Villafruela JJ, Cano T, et al. Mycophenolate mofetil levels in stable kidney transplant recipients. *Transplant Proc* 2007 Sep;39(7):2182-4.
- [19] Fernandez A, Martins J, Villafruela JJ, Marcen R, Pascual J, Cano T, et al. Variability of mycophenolate mofetil trough levels in stable kidney transplant patients. *Transplant Proc* 2007 Sep;39(7):2185-6.
- [20] Oellerich M, Shipkova M, Schutz E, Wieland E, Weber L, Tonshoff B, et al. Pharmacokinetic and metabolic investigations of mycophenolic acid in pediatric patients after renal transplantation: implications for therapeutic drug monitoring. German Study Group on Mycophenolate Mofetil Therapy in Pediatric Renal Transplant Recipients. *Ther Drug Monit* 2000 Feb;22(1):20-6.

- [21] Smak Gregoor PJ, van Gelder T, van Riemsdijk-van Overbeeke IC, Vossen AC, Ijzermans JN, Weimar W. Unusual presentation of herpes virus infections in renal transplant recipients exposed to high mycophenolic acid plasma concentrations. *Transpl Infect Dis* 2003 Jun;5(2):79-83.
- [22] Mardigyan V, Giannetti N, Cecere R, Besner JG, Cantarovich M. Best single time points to predict the area-under-the-curve in long-term heart transplant patients taking mycophenolate mofetil in combination with cyclosporine or tacrolimus. *J Heart Lung Transplant* 2005 Oct;24(10):1614-8.
- [23] Jirasiritham S, Sumethkul V, Mavichak V, Na-Bangchang K. The pharmacokinetics of mycophenolate mofetil in Thai kidney transplant recipients. *Transplant Proc* 2004 Sep;36(7):2076-8.
- [24] Pape L, Ehrich JH, Offner G. Long-term follow-up of pediatric transplant recipients: mycophenolic acid trough levels are not a good indicator for long-term graft function. *Clin Transplant* 2004 Oct;18(5):576-9.
- [25] Smak Gregoor PJ, van Gelder T, Hesse CJ, van der Mast BJ, van Besouw NM, Weimar W. Mycophenolic acid plasma concentrations in kidney allograft recipients with or without cyclosporin: a cross-sectional study. *Nephrol Dial Transplant* 1999 Mar;14(3):706-8.
- [26] Gregoor PJ, de Sevaux RG, Hene RJ, Hesse CJ, Hilbrands LB, Vos P, et al. Effect of cyclosporine on mycophenolic acid trough levels in kidney transplant recipients. *Transplantation* 1999 Nov 27;68(10):1603-6.
- [27] Sumethkul V, Na-Bangchang K, Kantachuvesiri S, Jirasiritham S. Standard dose enteric-coated mycophenolate sodium (myfortic) delivers rapid therapeutic mycophenolic acid exposure in kidney transplant recipients. *Transplant Proc* 2005 Mar;37(2):861-3.
- [28] van Gelder T, Hilbrands LB, Vanrenterghem Y, Weimar W, de Fijter JW, Squifflet JP, et al. A randomized double-blind, multicenter plasma concentration controlled study of the safety and efficacy of oral mycophenolate mofetil for the prevention of acute rejection after kidney transplantation. *Transplantation* 1999 Jul 27;68(2):261-6.
- [29] Hubner GI, Eismann R, Sziegoleit W. Drug interaction between mycophenolate mofetil and tacrolimus detectable within therapeutic mycophenolic acid monitoring in renal transplant patients. *Ther Drug Monit* 1999 Oct;21(5):536-9.
- [30] Zucker K, Rosen A, Tsaroucha A, de Faria L, Roth D, Ciancio G, et al. Unexpected augmentation of mycophenolic acid pharmacokinetics in renal transplant patients receiving tacrolimus and mycophenolate mofetil in combination therapy, and analogous in vitro findings. *Transpl Immunol* 1997 Sep;5(3):225-32.
- [31] Filler G, Zimmering M, Mai I. Pharmacokinetics of mycophenolate mofetil are influenced by concomitant immunosuppression. *Pediatr Nephrol* 2000 Feb;14(2):100-4.
- [32] Kagaya H, Miura M, Satoh S, Inoue K, Saito M, Inoue T, et al. No pharmacokinetic interactions between mycophenolic acid and tacrolimus in renal transplant recipients. *J Clin Pharm Ther* 2008 Apr;33(2):193-201.
- [33] Pawinski T, Hale M, Korecka M, Fitzsimmons WE, Shaw LM. Limited sampling strategy for the estimation of mycophenolic acid area under the curve in adult renal

- transplant patients treated with concomitant tacrolimus. *Clin Chem* 2002 Sep;48(9):1497-504.
- [34] Picard N, Premaud A, Rousseau A, Le MY, Marquet P. A comparison of the effect of ciclosporin and sirolimus on the pharmacokinetics of mycophenolate in renal transplant patients. *Br J Clin Pharmacol* 2006 Oct;62(4):477-84.
- [35] Cattaneo D, Merlini S, Zenoni S, Baldelli S, Gotti E, Remuzzi G, et al. Influence of co-medication with sirolimus or cyclosporine on mycophenolic acid pharmacokinetics in kidney transplantation. *Am J Transplant* 2005 Dec;5(12):2937-44.
- [36] Cattaneo D, Perico N, Gaspari F, Gotti E, Remuzzi G. Glucocorticoids interfere with mycophenolate mofetil bioavailability in kidney transplantation. *Kidney Int* 2002 Sep;62(3):1060-7.
- [37] Gonzalez-Roncero FM, Gentil MA, Brunet M, Algarra G, Pereira P, Cabello V, et al. Pharmacokinetics of mycophenolate mofetil in kidney transplant patients with renal insufficiency. *Transplant Proc* 2005 Nov;37(9):3749-51.
- [38] Wolfe EJ, Mathur V, Tomlanovich S, Jung D, Wong R, Griffy K, et al. Pharmacokinetics of mycophenolate mofetil and intravenous ganciclovir alone and in combination in renal transplant recipients. *Pharmacotherapy* 1997 May;17(3):591-8.
- [39] Miura M, Satoh S, Inoue K, Kagaya H, Saito M, Suzuki T, et al. Influence of lansoprazole and rabeprazole on mycophenolic acid pharmacokinetics one year after renal transplantation. *Ther Drug Monit* 2008 Feb;30(1):46-51.
- [40] Kofler S, Deutsch MA, Bigdeli AK, Shvets N, Vogeser M, Mueller TH, et al. Proton pump inhibitor co-medication reduces mycophenolate acid drug exposure in heart transplant recipients. *J Heart Lung Transplant* 2009 Jun;28(6):605-11.
- [41] Schaier M, Scholl C, Scharpf D, Hug F, Bonisch-Schmidt S, Dikow R, et al. Proton pump inhibitors interfere with the immunosuppressive potency of mycophenolate mofetil. *Rheumatology (Oxford)* 2010 Nov;49(11):2061-7.
- [42] Kiberd BA, Wrobel M, Dandavino R, Keown P, Gourishankar S. The role of proton pump inhibitors on early mycophenolic acid exposure in kidney transplantation: evidence from the CLEAR study. *Ther Drug Monit* 2011 Feb;33(1):120-3.
- [43] Kofler S, Wolf C, Shvets N, Sisic Z, Muller T, Behr J, et al. The proton pump inhibitor pantoprazole and its interaction with enteric-coated mycophenolate sodium in transplant recipients. *J Heart Lung Transplant* 2011 May;30(5):565-71.
- [44] Cattaneo D, Gaspari F, Ferrari S, Stucchi N, Del PL, Perico N, et al. Pharmacokinetics help optimizing mycophenolate mofetil dosing in kidney transplant patients. *Clin Transplant* 2001 Dec;15(6):402-9.
- [45] Weber LT, Shipkova M, Armstrong VW, Wagner N, Schutz E, Mehls O, et al. The pharmacokinetic-pharmacodynamic relationship for total and free mycophenolic Acid in pediatric renal transplant recipients: a report of the german study group on mycophenolate mofetil therapy. *J Am Soc Nephrol* 2002 Mar;13(3):759-68.
- [46] Le Meur Y, Buchler M, Thierry A, Caillard S, Villemain F, Lavaud S, et al. Individualized mycophenolate mofetil dosing based on drug exposure significantly improves patient outcomes after renal transplantation. *Am J Transplant* 2007 Nov;7(11):2496-503.

- [47] van GT, Silva HT, de Fijter JW, Budde K, Kuypers D, Tyden G, et al. Comparing mycophenolate mofetil regimens for de novo renal transplant recipients: the fixed-dose concentration-controlled trial. *Transplantation* 2008 Oct 27;86(8):1043-51.
- [48] Gaston RS, Kaplan B, Shah T, Cibrik D, Shaw LM, Angelis M, et al. Fixed- or controlled-dose mycophenolate mofetil with standard- or reduced-dose calcineurin inhibitors: the Opticept trial. *Am J Transplant* 2009 Jul;9(7):1607-19.
- [49] van GT, Silva HT, de FH, Budde K, Kuypers D, Mamelok RD, et al. How delayed graft function impacts exposure to mycophenolic acid in patients after renal transplantation. *Ther Drug Monit* 2011 Apr;33(2):155-64.
- [50] Knoll GA, MacDonald I, Khan A, Van Walraven C. Mycophenolate mofetil dose reduction and the risk of acute rejection after renal transplantation. *J Am Soc Nephrol* 2003 Sep;14(9):2381-6.
- [51] Takemoto SK, Pinsky BW, Schnitzler MA, Lentine KL, Willoughby LM, Burroughs TE, et al. A retrospective analysis of immunosuppression compliance, dose reduction and discontinuation in kidney transplant recipients. *Am J Transplant* 2007 Dec;7(12):2704-11.
- [52] Tierce JC, Porterfield-Baxa J, Petrilla AA, Kilburg A, Ferguson RM. Impact of mycophenolate mofetil (MMF)-related gastrointestinal complications and MMF dose alterations on transplant outcomes and healthcare costs in renal transplant recipients. *Clin Transplant* 2005 Dec;19(6):779-84.
- [53] Machnicki G, Ricci JF, Brennan DC, Schnitzler MA. Economic impact and long-term graft outcomes of mycophenolate mofetil dosage modifications following gastrointestinal complications in renal transplant recipients. *Pharmacoeconomics* 2008;26(11):951-67.
- [54] Bunnapradist S, Lentine KL, Burroughs TE, Pinsky BW, Hardinger KL, Brennan DC, et al. Mycophenolate mofetil dose reductions and discontinuations after gastrointestinal complications are associated with renal transplant graft failure. *Transplantation* 2006 Jul 15;82(1):102-7.
- [55] Kamar N, Oufroukhi L, Faure P, Ribes D, Cointault O, Lavayssiere L, et al. Questionnaire-based evaluation of gastrointestinal disorders in de novo renal-transplant patients receiving either mycophenolate mofetil or enteric-coated mycophenolate sodium. *Nephrol Dial Transplant* 2005 Oct;20(10):2231-6.
- [56] Budde K, Glander P, Diekmann F, Dragun D, Waiser J, Fritsche L, et al. Enteric-coated mycophenolate sodium: safe conversion from mycophenolate mofetil in maintenance renal transplant recipients. *Transplant Proc* 2004 Mar;36(2 Suppl):524S-7S.
- [57] Sanchez-Fructuoso A, Ruiz JC, Rengel M, Andres A, Morales JM, Beneyto I, et al. Use of mycophenolate sodium in stable renal transplant recipients in Spain: preliminary results of the MIDATA study. *Transplant Proc* 2009 Jul;41(6):2309-12.
- [58] Chan L, Mulgaonkar S, Walker R, Arns W, Ambuhl P, Schiavelli R. Patient-reported gastrointestinal symptom burden and health-related quality of life following conversion from mycophenolate mofetil to enteric-coated mycophenolate sodium. *Transplantation* 2006 May 15;81(9):1290-7.

- [59] Robaeys G, Cassiman D, Verslype C, Monbaliu D, Aerts R, Pirenne J, et al. Successful conversion from mycophenolate mofetil to enteric-coated mycophenolate sodium (myfortic) in liver transplant patients with gastrointestinal side effects. *Transplant Proc* 2009 Mar;41(2):610-3.
- [60] Arns W. Noninfectious gastrointestinal (GI) complications of mycophenolic acid therapy: a consequence of local GI toxicity? *Transplant Proc* 2007 Jan;39(1):88-93.
- [61] Mourad M, Malaise J, Chaib ED, De MM, Konig J, Schepers R, et al. Correlation of mycophenolic acid pharmacokinetic parameters with side effects in kidney transplant patients treated with mycophenolate mofetil. *Clin Chem* 2001 Jan;47(1):88-94.
- [62] Lu YP, Zhu YC, Liang MZ, Nan F, Yu Q, Wang L, et al. Therapeutic drug monitoring of mycophenolic acid can be used as predictor of clinical events for kidney transplant recipients treated with mycophenolate mofetil. *Transplant Proc* 2006 Sep;38(7):2048-50.
- [63] Mourad M, Malaise J, Chaib ED, De Meyer M, Konig J, Schepers R, et al. Pharmacokinetic basis for the efficient and safe use of low-dose mycophenolate mofetil in combination with tacrolimus in kidney transplantation. *Clin Chem* 2001;47(7):1241-8.
- [64] Mourad M, Malaise J, Chaib ED, De Meyer M, Konig J, Schepers R, et al. Correlation of mycophenolic acid pharmacokinetic parameters with side effects in kidney transplant patients treated with mycophenolate mofetil. *Clin Chem* 2001 Jan;47(1):88-94.
- [65] Shah T, Tellez-Corrales E, Yang JW, Qazi Y, Wang J, Wilson J, et al. The pharmacokinetics of enteric-coated mycophenolate sodium and its gastrointestinal side effects in de novo renal transplant recipients of Hispanic ethnicity. *Ther Drug Monit* 2011 Feb;33(1):45-9.
- [66] Kuypers DR, de JH, Naesens M, de LH, Halewijck E, Dekens M, et al. Current target ranges of mycophenolic acid exposure and drug-related adverse events: a 5-year, open-label, prospective, clinical follow-up study in renal allograft recipients. *Clin Ther* 2008 Apr;30(4):673-83.
- [67] Langman LJ, Shapiro AM, Lakey JR, LeGatt DF, Kneteman NM, Yatscoff RW. Pharmacodynamic assessment of mycophenolic acid-induced immunosuppression by measurement of inosine monophosphate dehydrogenase activity in a canine model. *Transplantation* 1996 Jan 15;61(1):87-92.
- [68] Glander P, Braun KP, Hambach P, Bauer S, Mai I, Roots I, et al. Non-radioactive determination of inosine 5'-monophosphate dehydrogenase (IMPDH) in peripheral mononuclear cells. *Clin Biochem* 2001 Oct;34(7):543-9.
- [69] Langman LJ, LeGatt DF, Halloran PF, Yatscoff RW. Pharmacodynamic assessment of mycophenolic acid-induced immunosuppression in renal transplant recipients. *Transplantation* 1996 Sep 15;62(5):666-72.
- [70] Glander P, Hambach P, Braun KP, Fritsche L, Waiser J, Mai I, et al. Effect of mycophenolate mofetil on IMP dehydrogenase after the first dose and after long-

- term treatment in renal transplant recipients. *Int J Clin Pharmacol Ther* 2003 Oct;41(10):470-6.
- [71] Fukuda T, Goebel J, Thogersen H, Maseck D, Cox S, Logan B, et al. Inosine Monophosphate Dehydrogenase (IMPDH) Activity as a Pharmacodynamic Biomarker of Mycophenolic Acid Effects in Pediatric Kidney Transplant Recipients. *J Clin Pharmacol* 2010 Apr 23.
- [72] Vethe NT, Mandla R, Line PD, Midtvedt K, Hartmann A, Bergan S. Inosine monophosphate dehydrogenase activity in renal allograft recipients during mycophenolate treatment. *Scand J Clin Lab Invest* 2006;66(1):31-44.
- [73] Sanquer S, Breil M, Baron C, Dhamane D, Astier A, Lang P. Induction of inosine monophosphate dehydrogenase activity after long-term treatment with mycophenolate mofetil. *Clin Pharmacol Ther* 1999 Jun;65(6):640-8.
- [74] Sanquer S, Maison P, Tomkiewicz C, Macquin-Mavier I, Legendre C, Barouki R, et al. Expression of inosine monophosphate dehydrogenase type I and type II after mycophenolate mofetil treatment: a 2-year follow-up in kidney transplantation. *Clin Pharmacol Ther* 2008 Feb;83(2):328-35.
- [75] Bremer S, Mandla R, Vethe NT, Rasmussen I, Rootwelt H, Line PD, et al. Expression of IMPDH1 and IMPDH2 after transplantation and initiation of immunosuppression. *Transplantation* 2008 Jan 15;85(1):55-61.
- [76] Bremer S, Vethe NT, Rootwelt H, Bergan S. Expression of IMPDH1 is regulated in response to mycophenolate concentration. *Int Immunopharmacol* 2009 Feb;9(2):173-80.
- [77] Brunet M, Martorell J, Oppenheimer F, Vilardell J, Millan O, Carrillo M, et al. Pharmacokinetics and pharmacodynamics of mycophenolic acid in stable renal transplant recipients treated with low doses of mycophenolate mofetil. *Transpl Int* 2000;13 Suppl 1:S301-S305.
- [78] Chiarelli LR, Molinaro M, Libetta C, Tinelli C, Cosmai L, Valentini G, et al. Inosine monophosphate dehydrogenase variability in renal transplant patients on long-term mycophenolate mofetil therapy. *Br J Clin Pharmacol* 2010 Jan;69(1):38-50.
- [79] Wang J, Yang JW, Zeevi A, Webber SA, Girnita DM, Selby R, et al. IMPDH1 gene polymorphisms and association with acute rejection in renal transplant patients. *Clin Pharmacol Ther* 2008 May;83(5):711-7.
- [80] Kagaya H, Miura M, Saito M, Habuchi T, Satoh S. Correlation of IMPDH1 gene polymorphisms with subclinical acute rejection and mycophenolic acid exposure parameters on day 28 after renal transplantation. *Basic Clin Pharmacol Toxicol* 2010 Aug;107(2):631-6.
- [81] Glander P, Hambach P, Braun KP, Fritsche L, Giessing M, Mai I, et al. Pre-transplant inosine monophosphate dehydrogenase activity is associated with clinical outcome after renal transplantation. *Am J Transplant* 2004 Dec;4(12):2045-51.
- [82] Rath T, Kupper M. Comparison of inosine-monophosphate-dehydrogenase activity in patients with enteric-coated mycophenolate sodium or mycophenolate mofetil after renal transplantation. *Transplant Proc* 2009 Jul;41(6):2524-8.

- [83] Glander P, Sommerer C, Arns W, Ariatabar T, Kramer S, Vogel EM, et al. Pharmacokinetics and pharmacodynamics of intensified versus standard dosing of mycophenolate sodium in renal transplant patients. Clin J Am Soc Nephrol 2010 Mar;5(3):503-11.

Renal Transplantation - Updates and Advances

Edited by Dr. Layron Long

ISBN 978-953-51-0173-4

Hard cover, 234 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

This book presents a nice international compilation of scholarly papers and chapters which address the latest advances in renal transplant surgery. These works cover a variety of topics; the last advance and success of renal transplant science: biochemistry, immunology, molecular genetics, pharmacology - pharmacogenetics, pediatric transplant and a few rare uropathies that warrant organ replacement.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Thomas Rath and Manfred Küpper (2012). Pharmacokinetics and Pharmacodynamics of Mycophenolate in Patients After Renal Transplantation, Renal Transplantation - Updates and Advances, Dr. Layron Long (Ed.), ISBN: 978-953-51-0173-4, InTech, Available from: <http://www.intechopen.com/books/renal-transplantation-updates-and-advances/pharmacokinetics-and-pharmacodynamics-of-mycophenolate-in-patients-after-renal-transplantation>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.