Are high levels of Lp(a) associated with familial hypercholesterolemia?

L Schreier¹, V Zago¹, L Cacciagiú¹, M Bursztyn¹, V Bañares², G López¹, G Berg¹, P Corral³.
1-Universidad de Buenos Aires, Departamento de Bioquímica Clínica-Laboratorio de Lípidos y Aterosclerosis, Buenos Aires, Argentina. 2-Centro Nacional de Genética Médica-ANLIS- Dr. Carlos Malbrán, Departamento de Genética Experimental, Buenos Aires, Argentina. 3-Universidad FASTA, Facultad de Medicina-Departamento de Investigación, Mar del Plata, Argentina.

- Lp(a) is a pro athero-thrombotic lipoprotein, which concentration is mainly genetically determined.
- Familial Hypercholesterolemia (FH)-characterized by high LDL-C levels from early age- is associated with premature atherosclerotic cardiovascular events.
- Lp(a) contains LDL in its structure. This leads to postulate that FH is associated with increased Lp(a)
- There are controversial results about whether FH patients have high Lp (a) levels or not.

Aim: To analyze the levels of Lp(a) in patients with clinical and/or genetic diagnosis of FH.

Methods:

- 48% of the patients were under hypolipidemic treatment LDL-C were corrected by factors depending on each type/dose of medication (Haralambos K, et al Atherosclerosis 2015, 240:190)

Biochemical profile

- Tot. Cholesterol
- Triglycerides
- HDL-C, LDL-C
- Apo B
- Lp(a)** (nmol/L)

Standardized and Automated methods

Automated Immunoturbidimetric assay with Polyclonal antibodies a-human Lp(a). Reference material: Preciset Lp(a) Gen2, traceability with SRM2B, IFCC/WHO, independent of the size apo (a).

Gene Panel Sequencing for FH patients*

- LDLR
- APOB
- PCSK9
- LDL-RAP1
- APOE

NGS: MiSeq (Illumina)

*The NGS and Bioinformatic analysis were performed in 2 Labs: 1-Boston Heart Diagnosis, Boston, MA-USA. 2-Instituto Nacional de Saúde Dr R. Jorge, Lisboa, Portugal.

Statistical Analysis: Non parametric Tests : U-Mann Whitney and Kruskal Wallis for multiple groups, and parametric tests: ANOVA and Scheffe.

Results:

No differences detected in Lp(a) concentration in patients grouped according to their Dutch Lipid Clinic Network (DLCN) Criteria for FH clinical diagnosis, n=140.

DLCN score	Lp(a) nmol/l Median (range)
Definite (>8), n=46	36.6 (2.0 - 618.0)
Probable (6-8), n=56	38.9 (1.3 - 523.2)
Possible (4-5) n=26	33.0 (3.1 - 311.0)
Unlikely (≤3) n=12	119.3 (4.9 - 297.0)

p=0.285; Test Kruskal-Wallis

No differences in LDL-C levels, between groups defined by Lp(a) tertiles.

p= 0.163 (ANOVA)

Lp(a) Tertiles (nmol/L)

Actual LDL-C after subtracting the contribution of Lp(a)-C: Tertile with the highest Lp(a) showed decreased LDL-C.

Estimation of Lp(a)-C:

- Lp(a) nmol/L/2.4=mg/dL
- Lp(a) mg/dl x 0.3= Lp(a)-C
- LDL-C -Lp(a)-C= true LDL-C

Lp(a) Tertiles (nmol/L)

p=0.017, ANOVA / Scheffé

Lp(a) in patients genotyped for FH (NGS for LDLR, APOB, PCSK9, APOE and LDLRap1),(n=95)

No differences were observed between patients with Positive or Negative genetic study.

Lp(a) Median (range), nmol/L	
Carriers of genetic variants (n=34)	Non-carriers of genetic variants (n=61)
32.1 (3.8-618.0)	44.6 (1.3-350.0)

p=0.214; U-Mann WhitneyTest

Conclusions: a-In a subgroup of our hypercholesterolemic cohort, we verified an independence between Lp(a) and LDL-C levels.
b- Lp(a) did not show associations with FH clinical diagnosis (DLCN) nor with FH genetic testing results.
c- However, it must be highlighted that the coexistence of increased LDL-C and Lp(a) levels impair the severity and prognosis of CVD
d-Testing for elevated Lp(a) during cascade screening for FH is effective in identifying relatives with high Lp(a) and increased risk ASCVD (SAFEHEART, Mata P et al, 2019, 73,9:1029)