Influence of preoperative disability in the recovery of elective cancer surgery

Braga A.¹, Silva D.¹, Carvalho B.¹, Neto S.¹, Santos A.¹, Abelha F.^{1,2}

¹Hospital São João, Dept of Anaesthesiology / ²Faculdade de Medicina Universidade do Porto (FMUP)

Background and Goal of Study:

Disability is defined by the World Health Organization (WHO) as difficulties in any area of functioning as they relate to environmental and personal factors. In surgical patients, it is assessed by WHO Disability Assessment Schedule (WHODAS) 2.0. The aim of this study was to evaluate the influence of preoperative disability in the recovery after cancer surgery.

Materials and Methods:

- Prospective observational study (Jun 2016 to Oct 2016)
 - **Inclusion criteria:** Patients scheduled for elective cancer surgeries with a curative intent.
 - **Exclusion criteria:** Age < 18 years old and inability to give informed consent.
 - **Preoperative disability:** ≥ 25 WHODAS 2.0.
 - **Post-Operative Quality of Recovery:**
 - **Quality of Recovery-15 (QoR-15)** at baseline (T0) and 24 hours (D1) after surgery.
 - **Post-Operative Quality of Recovery Scale (PQRS)** at baseline and after surgery at minute 15 (T15), 40 (T40) and days 1 and 3 (D1, D3), evaluating recovery in five domains.
- Recovery** was defined as return to baseline values or better for all questions within each domain.
- The **EuroQol five-dimension questionnaire (EQ-5D)**, and **Clinical Frailty Scale** were applied preoperatively.
 - The Chi-square, Fischer's exact or Mann-Whitney U tests were used for comparisons. Variables were considered significant if $p < 0.05$.

Results and Discussion:

- 148 included patients

Table 1 – Univariate analysis for demographics, EQ-5D, Frailty

	Non-Disability N=116	Disability N= 32	p value
Age, median [P25-P75]	62.5 [52.3-72.8]	74.0 [64.3-79.5]	<0.001 ^c
ASA $\geq$ III, n (%)	30 (25.9)	21 (65.6)	<0.001 ^a
RCRI > 2, n (%)	12 (10.3)	13 (40.6)	<0.001 ^a
EQ-5D VAS, median [P25-P75]	70 [50.0-80.0]	50 [32.5-58.75]	<0.001 ^c
Frailty, n (%)	19 (16.4)	26 (81.3)	<0.001 ^a

Table 2 – Recovery according to PQRS

Incomplete recovery PQRS: domains		Non-Disability	Disability	p value
physiological (%)	T15	100 (86.2)	30 (93.8)	0.202 ^a
	T40	79 (68.7)	27 (87.1)	0.030^b
	D1	24 (21.6)	14 (43.7)	0.013^a
nociceptive, n (%)	T15	20 (17.2)	15 (46.9)	<0.001 ^a
	T40	29 (25.2)	13 (41.9)	0.068 ^a
	D1	26 (23.2)	12 (38.7)	0.084 ^a
	D3	21 (18.9)	15 (51.7)	<0.001 ^a
Emotive, n (%)	T15	73 (62.9)	25 (78.1)	0.108 ^a
	T40	73 (63.5)	24 (77.4)	0.145 ^a
	D1	66 (58.4)	21 (70.0)	0.248 ^a
	D3	63 (56.3)	21 (75.0)	0.070 ^a
Cognitive, n (%)	T15	107 (92.2)	30 (93.8)	0.562 ^b
	T40	97 (84.3)	26 (83.9)	0.570 ^b
	D1	94 (83.2)	22 (73.3)	0.220 ^a
	D3	86 (86.0)	25 (83.3)	0.537 ^a
Funcional n (%)	T40	95 (82.6)	24 (77.4)	0.509 ^a
	D1	74 (66.7)	22 (68.8)	0.825 ^a
	D3	57 (51.8)	18 (60.0)	0.426 ^a
Complete recovery PQRS		Non-Disability	Disability	p value
T15, n (%)				
4 domains		6 (5.2)	0 (0)	0.189 ^a
3 domains		52 (44.8)	5 (15.6)	0.003^a
2 domains		104 (89.7)	21 (67.7)	0.002^a
T40, n (%)				
4 domains		17 (14.9)	0 (0)	0.013^a
3 domains		60 (52.6)	6 (19.4)	0.001^a
2 domains		100 (87.7)	25 (80.6)	0.230 ^a
D1, n (%)				
4 domains		16 (14.7)	3 (10.0)	0.374 ^a
3 domains		57 (52.3)	9 (30.0)	0.030^a
2 domains		91 (83.5)	19 (63.3)	0.016^a
D3, n (%)				
4 domains		29 (26.6)	2 (6.9)	0.024^a
3 domains		79 (72.5)	11 (37.9)	0.001^a
2 domains		105 (96.3)	24 (82.8)	0.020^a

Table 3 – Recovery according to QoR-15

QoR-15 T0, median [P25-P75]	134 [123-141]	105 [92-119]	<0.001 ^c
QoR-15 D1, median [P25-P75]	114 [101-127]	108 [87-122]	0.168 ^c

a) Chi-square; b) Fischer's exact; c) Mann-Whitney U tests

Conclusion:

Patients with disability had more comorbidities, a worst health status before surgery and revealed a poor quality of recovery, suggesting an adverse impact of disability in health status and in postoperative recovery.