Regional Anaesthesia and the Diagnosis of Compartment Syndrome:

A Systematic Review

Michael Shaw¹, Emil Shumanov²

1. Anaesthetics Trainee, Barts & The London School of Anaesthesia, London, UK. 2. Consultant Anaesthetist, Newham University Hospital, Barts Health NHS Trust, London, UK

Background

Acute compartment syndrome (ACS) is caused by tissue swelling within an encapsulated muscle – usually within 72 hours of traumatic or iatrogenic insult.^[1] This causes compression of blood vessels leading to compromised blood flow and ultimately, ischaemia. ACS is most likely to affect patients with tibial or forearm fractures; while other risk factors include age <35 years, injury severity and male gender.^[2]

Diagnosis of ACS is a time critical intervention, with significant morbidity and mortality in cases of delayed diagnosis.^[1] However, diagnosis may be challenging, with irretractable pain a significant symptom. Given the importance of pain in the diagnosis of ACS, there is evidence of reluctance of both surgeons and anaesthetists in the use of regional anaesthesia (RA) in patients at risk of ACS.^[3] Therefore, the aim of this systematic review was to explore what evidence, if any, exists on the use of RA and delayed diagnosis of ACS.

Method

We conducted a systematic literature review according to PRISMA recommendations. We used the Healthcare Database Advanced Search (HDAS) system provided by the National Institute for Health and Care Excellence (NICE). HDAS allows for search of multiple databases simultaneously to provide a comprehensive library of results. The search terms used were (“compartment syndrome” AND (“spinal anesthesia” OR “spinal anesthesia” OR “regional anaesthesia” OR “regional anesthesia”)). There were no historic or language restrictions. All relevant articles were assessed according to GRADE criteria.

References

1. Elliott KG, Johnstone AJ. Diagnosing acute compartment syndrome. *Journal of Bone and Joint Surgery*. 2003;85:625-632
2. Martin JT. Compartment syndromes: concepts and perspectives for the anesthesiologist. *Anesthesia and Analgesia*. 1992;75:275-283
3. Wordsworth H, Wickham A, Bellew B, et al. Attitudes towards safety of regional anaesthesia in trauma patients: A survey of surgeons and anaesthetists in a Major Trauma Centre and a Specialist Hand Unit. [Conference poster] Chelsea and Westminster NHS Trust. 2006
4. Kent ML, Buckenmaier CC. Battlefield regional anaesthesia: evolution and future concepts. *Techniques in Regional Anesthesia and Pain Management*. 2012;16:184-189
5. Stojadinovic A, Auton A, Peoples G, et al. Responding to challenges in modern combat casualty care: innovative use of advanced regional anaesthesia. *Pain Medicine*. 2006;7(4):330-338

Full reference list available upon request. (michaelwilliam.shaw@nhs.net)

Results

* Articles excluded as: 16 dealt with paediatrics and 11 with intravenous RA
† Articles excluded as: 15 unrelated to orthopaedic ACS and 23 unrelated to diagnosis of ACS following RA

Of the 28 included papers there were: 9 case reports, 12 review articles, 1 conference poster, 1 conference abstract, 1 editorial, 1 letter and 3 surveys. No articles discussed the diagnosis of RA in the context of neuroaxial (i.e. spinal) anaesthesia.

Discussion

Evidence suggesting delayed diagnosis of ACS in the context of RA was of very low quality by GRADE criteria. Of the case reports, those where RA was judged to contribute to delayed ACS diagnosis revealed significant flaws in aftercare of RA patients. 4 case reports describe early diagnosis of ACS in the presence of working RA. The surveys revealed between 4-32% of respondents had anecdotal experience of delayed ACS diagnosis with RA use.

The review articles were generally in favour of RA use. The review articles from the military^[4-5] are of particular interest due to the high prevalence of risk factors for ACS. They are unified in their support for RA following traumatic limb injuries and reported no cases or concerns that RA may be contributing to delayed ACS diagnosis.

Conclusions

- No prospective studies conducted on this topic
- Scant low quality evidence RA delays ACS diagnosis
- Many publications support RA use in at-risk patients
- Reluctance to use RA is common in clinician surveys
- Withholding RA denies powerful analgesia with little evidence of protecting against early ACS diagnosis