INCREASED SERUM CONCENTRATION OF C-TERMINAL PORTION OF PRO-ARGININE-VASOPRESSIN IN PRETERM NEWBORNS AND IMPAIRED RESPONSE TO ELEVATED EFFECTIVE OSMOLALITY IN THE FIRST DAYS OF LIFE

¹Anna Suchojad, ¹Anna Tarko, ²Barbara Przybył, ³Aniceta Brzozowska, ¹Iwona Maruniak-Chudek
¹Intensive Care and Neonatal Pathology Unit, Faculty of Medicine in Katowice, Medical University of Silesia, Katowice, Poland; ² Clinical Unit of Neonatology and Intensive Care, Multidisciplinary Hospital E. Warmiński, Bydgoszcz, Poland; ³Department of Pathophysiology, Faculty of Medicine in Katowice, Medical University of Silesia, Katowice, Poland

BACKGROUND AND AIMS

Copeptin (CTproAVP) is regarded as a new marker of dehydration, but its action in premature newborns is poorly investigated. The aim of this study was to evaluate the relation between serum effective osmolality and serum CTproAVP concentration and the analysis of factors potentially influencing CTproAVP in preterm newborns in their first days of life.

METHODS

155 newborns, 30+0(N=49) to 36+6(N=106) weeks (wks) of gestational age (GA) were enrolled. They underwent routine medical management based on clinical condition and additionally CTproAVP was assessed in the first 48 hours of life.

RESULTS

Markedly higher CTproAVP levels were observed in less mature neonates below 34 wks GA than in those ≥ 34 wks GA [198 (1-3Q:117–349) vs 151 (96-257) pmol/L, p=0.04], regardless of similar effective serum osmolality and gender. Neonates delivered by caesarean section (CS) and receiving total parenteral nutrition (TPN) presented higher values of CTproAVP. Serum CTproAVP concentrations were inversely related to GA (R=-0.205, p=0.01), birth weight (R=-0.201, p=0.01), Apgar 1' (R=-0.188, p=0.02) and Apgar 5' (R=-0.171, p=0.04), but not to effective osmolality. In multiple regression analysis only GA was explaining CTproAVP variability (β=0.168, p=0.04).

CONCLUSIONS

CTproAVP concentration should be considered mostly as a marker of stress during the early adaptation period. Higher concentrations of CTproAVP in immature newborns are rather related to stress factors accompanying medical interventions and not immaturity itself.

Scientific Grant of Medical University of Silesia
No: KNW-2-K19/N/6/N

Table 1. Demographic factors and clinical parameters of the study population presented for all preterm newborns and separately for two subgroups divided according to age. *-median

	All [N=150]	Newborns < 34 wks GA [N=49]	Newborns ≥34 wks GA [N=106]	Statistical significance
Gender (M/F)	93/62	30/19	63/43	P=0.97
Gestational age (wks)*	35 [33-36]	32 [32-33]	36 [35-36]	P=0.15
Birth weight (g)	2345 [2253-2436]	1843 [1729-1958]	2576 [2482-2670]	P<0.001
Apgar 1' (pts)*	9 [8-9]	8 [7-8]	9 [8-10]	P=0.63
Apgar 5' (pts)*	9 [8-10]	8 [8-9]	10 [9-10]	P=0.63
CTproAVP (pmol/L)*	157 [104-273]	198 [117-349]	151 [96-257]	P=0.01
Mode of delivery (CS/VG)	94/61	36/13	58/48	P=0.04
Effective serum osmolality (mOsm/L)	284 [283-286]	284 [281-287]	285 [283-286]	P=0.63
TPN	19	12	7	P=0.004
Urine osmolality (mOsm/l)*	222 [151-314]	156 [131-248]	256 [171-351]	P=0.01

Fig.1. CTproAVP in relation to Apgar 5'

Fig.2. Correlation between CTproAVP and birth weight