EVALUATION OF ULTRASOUND-GUIDED PERIPHERAL VENOUS CATHETERIZATION BY YOUNG RESIDENTS

Mehdi Hamdi^{*1}, Sana Boughariou¹, Said Guedri¹, Sameh Zakhama¹, Besma Zbidi¹, Mehdi Boussofara¹

¹Anesthesia, Trauma Center, Tunis, Tunisia

Background and Objectives:

Insertion of intravenous catheters is widely performed in the hospital, especially in emergency departments and intensive care units. While peripheral placement is the most common access, central venous catheters are used in case of peripheral cannulation failure. Ultrasound guidance may be helpful to determine the adequate catheter diameter and length as well as the adequate vein to catheterize. The ultrasound guidance technique has not been widely discussed in the literature. Identification can be performed in two ways: transversally, which allows measuring the diameter vein and verifying its compliance, easier at the first venous access, and longitudinal which has the advantage of viewing the catheter throughout its way to the vein.

The aim of our study was to compare two techniques for setting up peripheral vein catheter according to transverse and longitudinal techniques, in terms of feasibility and complications.

Materials and Methods:

The agreement of the hospital ethics committee was approved. A prospective study was conducted between July 2015 and April 2016. After their informed consent, 100 patients aged between 18 and 65 admitted to the operating room for abdominal surgery were randomized into 2 groups: GL (insertion of peripheral venous catheter under ultrasound guidance using the longitudinal section) and GT (we use the transverse section). We excluded patients with a BMI > 35 kg / m².

An ultrasound probe (Sonosite) flathead and high frequency was used for the identification of veins. Ultrasonography was performed in vascular mode with the use of color Doppler.

The residents, who have received theoretical and practical training, try to spot a peripheral vein in the upper limb and introduce an 18 gauge catheter. The procedure was performed in two different sections: transverse and longitudinal view. We collected anthropometric characteristics of the patients, durations of ultrasound guidance and procedure, the number of attempts and the difficulty score (evaluated 0-10).

For variables with a normal distribution, the data are presented as mean $\pm$ standard deviation. Mean comparisons shall be effected by paired or unpaired Student's t-test. Percentages were compared by Chi-Squared test. The bivariate correlation was analyzed using the Pearson test. Statistical analysis was performed using SPSS 22 software.

Scott Joing MD et al. Ultrasound-Guided Peripheral IV placement. NEJM 2012.10:366-68

Results:

No significant differences between both groups regarding the anthropometric criteria (weight = 74.4 ± 16.8 kg; height = 168.9 ± 18.6 cm and BMI = 23.9 ± 2.7 kg / m²).

	GL	GT	p
Brachial vein	29/50	32/50	0,23
Basilic vein	15/50	14/50	0,57
Cephalic vein	6/50	4/50	0,66
Peripheral vein catheterization scheduled difficult	4/30	6/30	0,39
Identification duration (min)	3,7 $\pm$ 1,6	2,9 $\pm$ 0,8	0,07
Average procedure time (min)	4,8 $\pm$ 1,9	3,6 $\pm$ 1,1	0,05
Failure incidence	5/50	3/50	0,17
Difficulty Score(0 to 10)	4,7 $\pm$ 1,4	3,8 $\pm$ 1,2	0,04
Patient satisfaction(0 to 10)	8,3 $\pm$ 1,4	8,5 $\pm$ 0,9	0,47

Conclusion:

The establishment of peripheral venous catheters is easier under ultrasound guidance. Although the transverse view seems easier than the longitudinal view, they allow, both to reduce the duration of the procedure and the number of attempts. And so they provide better patient's satisfaction and reduce the cost by decreasing the number of catheters used.