Identification of Characteristics of Neighbourhoods with High Incidence of Out-of-Hospital Cardiac Arrest & Low Bystander Cardiopulmonary Resuscitation Rates

Terry Brown¹, Claire Hawkes¹, Scott Booth¹, Rachael Fothergill², Sara Black³, Anna Bichman⁴, Helen Pocock⁵, Jasmeet Soar⁶, Julian Mark⁷, Jonathan Benger⁸, Gavin Perkins¹, OHCAO collaborators

Purpose of the Study:

The community in which a person sustains an out-of-hospital cardiac arrest (OHCA) influences the likelihood that they will receive bystander CPR (BCPR) & ultimately survive. The aim of this research was to identify high risk communities in England with a high OHCA/low BCPR incidence to enable direct-targeted strategies to be designed & implemented.

Materials & Methods:

Analysis was of 67,219 OHCA events that occurred between 2013 & 2015 in England, where resuscitation was commenced or continued by EMS personnel. OHCA location was geocoded where possible; postcode district (POD) was used as proxy for neighbourhood. Neighbourhood-level variables (from 2011 Census) were linked to each geocoded address. Associations between individual-level & neighbourhood-level characteristics & BCPR was assessed by hierarchical regression analysis.

Results:

Areas with high incidence (Figure 1) included parts of London/West Midlands, the south-east/north-east coasts, & Lake District. Table 1 compares the demographic characteristics PODs with a high & low incidence. On average the ethnic composition of the OHCA location was 84.3% white (83.7% where OHCA was witnessed & received BCPR). Incidence increased significantly with deprivation (IMD) from 7.0% in least deprived areas to 12.3% in most deprived areas.

Low BCPR rates (Figure 2) were observed in the North-East & Yorkshire, & parts of Milton Keynes, Cornwall/Somerset, Derby, Wiltshire, Kent & Leicester. However, areas of the country with low resident population, would be expected to have low BCPR rates. Table 1 also compares the demographic characteristics of PODs with high & low BCPR rates.

BCPR rate decreased significantly with POPD from 75.5% (<100/hectare) to 60.9% (>10,000/hectare; $p<0.05$). There was a significant difference between areas with highest & lowest deprivation (63.0% vs 67.9%; $p<0.05$). The OR for receiving BCPR for every 1% increase in white ethnic population was 1.01, whereas for non-white ethnic groups the OR was 0.99.

Areas of high risk (high OHCA incidence/low BCPR rate) are shown in Figure 3 & their demographic characteristics compared in Table 1. A total of 56 PODs were identified as high risk, many included the coastline or were in city/town centres. Generally, these areas had significantly higher POPD, more deprivation, lower proportion of managers & whites, & higher proportion of non-whites.

Regression analysis indicated that deprivation, the proportion of the population whose activities were limited, & the socio-economic classification & education breakdown within a POD were significant predictors of whether an OHCA received BCPR or not. Using the final model 29 PODs were predicted to have BCPR rates less than they actually were, 38% of these were in South-Central & 21% in North-East.

Table 1: Demographic characteristics of postcode districts

	OHCA incidence (/100,000)		BCPR (%)		Risk	
	High (≥170)	Low (<170)	High (≥39)	Low (<39)	High	Low
Population density (/ha)	22.1	20.2	21.4	17.2	23.0	19.8
IMD	5.0	5.9	5.8	5.5	5.1	5.9
% Aged >65y	18.6	17.4	17.6	17.7	17.3	17.8
% Managers	31.2	33.0	33.0	30.6	9.5	11.5
% Routine & Manual Occupations	31.5	30.8	30.5	33.1		
% White	86.6	89.1	88.1	90.4	86.7	89.2
% Non-white	10.2	8.3	9.0	7.4	10.2	8.2

Conclusions:

The most deprived areas of England have a high OHCA incidence & low BCPR rate. We have identified several high risk areas of England that could be targeted for future initiatives; a special focus should be given to holiday destinations & town/city centres.

1 University of Warwick; 2 London Ambulance Service; 3 South Western Ambulance Service
4 East Midlands Ambulance Service; 5 South Central Ambulance Service; 6 Resuscitation Council UK
7 Yorkshire Ambulance Service/ NASMed; 8 UWE Bristol