Crack Lung

An Unusual Presentation of Paediatric Respiratory Failure

S. Davies¹, B. Lakin¹. ¹Alder Hey Children's Hospital, Paediatrics, Liverpool, United Kingdom.

The Case

We present 15 year old girl who developed severe respiratory failure following illicit drug use in the context of an unusual social background.

Presentation was with progressive shortness of breath resistant to treatment, that required mechanical ventilation. Initially it was presumed pneumonia.

CXR and CT showed multiple military nodules but multiple investigations including for T.B were negative. s A lung biopsy showed pneumonitis, likely secondary to inhalation injury or ‘Crack-Lung’, due to its bronchio-central distribution.

Cocaine is an alkaloid derived from the leaves of *Erythroxylon coca* originally chewed by indigenous South Americans.

‘Crack Cocaine’ is a form of cocaine that can be smoked and provides a short but intense high.

‘Crack Lung’ refers to a syndrome of diffuse alveolar damage and haemorrhagic aveolitis that occurs within 48 hours of inhaling ‘Crack Cocaine’.

Presenting Features of Smoking Cocaine	
Fever	Sputum production
Chest Pain	Asthma Symptoms
Back Pain	Eosonophilic Pneumonia
Shortness of Breath	Pneumothorax

Re-exploration of the **social history** identified that the family had become involved with a homeless person who had recently been released from prison with a history of drug misuse and sex offences. He had introduced the girl to a variety of substance misuse, including crack cocaine.

Injury from smoking cocaine can occur from a variety of mechanisms including: thermal injury, direct toxicity to cells, cut substances, Vasospasm, propagation of inflammation

Social history can be neglected during an acute presentation especially when ‘the story fits’ but remains an important part of information gathering to formulate differential diagnosis. In this case, the subsequent story did not fit as the course was so severe without a definitive cause.

Fortunately, ‘Crack-Lung’ is not frequently encountered by paediatricians but children can become involved with risk taking behaviour and therefore diagnosis not usually expected.