Glycaemic variability is associated with microvascular complications in Emiratis with Type 2, but not Type 1 diabetes

Mohamed Suliman¹, Adam Buckley², Nader Lessan² and Maha T Barakat²

¹ Imperial College London Diabetes Centre, (Al Ain, United Arab Emirates)

² Imperial College London Diabetes Centre, (Abu Dhabi, United Arab Emirates)

Background

Microvascular complications of diabetes mellitus are a major source of morbidity. Variability in blood glucose may contribute to the risk of these complications.

Aims

To determine the contribution of HbA1c or early morning serum glucose contribute to the risk of microvascular complications, controlling for established risk factors.

Methods

ICLDC patient database was accessed to identify people of Emirati origin with type 1 (n=994) or type 2 (n=18,904) diabetes, who underwent retinal screening between 01/01/2015 and 01/04/2017, with at least five recorded measurements of HbA1c and plasma glucose.

“Average” and **“variability”** of each were expressed as the mean and standard deviation (SD).

Retinal disease in the more severely affected eye was classified as: “none” (ROM0), “background” (RIM0) and “significant” (any higher grade).

Microalbuminuria was used as the indicator of nephropathy and was defined as urinary albumin/creatinine ratio ≥ 2.5 mg/mmol in females or ≥ 3.5 mg/mmol in males at both of the two most recent measurements.

Smoking was graded in ordered fashion as: “never smoked”, “passive smoking”, “ex-smoker” and “current smoker”

Logistic regression models were used to examine the contribution of average and variability of HbA1c and glucose to the likelihood of retinal or renal endpoints, **controlling** for diabetes duration, smoking, and most recently measured systolic blood pressure (sBP). Interaction terms were included for means and SDs of both HbA1c and glucose.

Results

Patients with type 1 diabetes had higher mean HbA1c (8.8% cf 7.4%, $p < 0.0001$), mean glucose (11.7 mmol/L cf 9.1 mmol/L, $p < 0.0001$), SD HbA1c (1.18% cf 0.88%, $p < 0.0001$), and SD glucose (5.12 mmol/L cf 2.25 mmol/L, $p < 0.0001$) compared to those with type 2 diabetes.

Figures: Standard deviation (SD) of morning glucose (mmol/L) by grade of retinal disease in people with Type 1 (A) and Type 2 (B) diabetes. SD of HbA1c (%) by grade of retinal disease in people with Type 1 (C) and type 2 (D) diabetes.

In both type 1 and type 2 diabetes, independent and significant predictors of microvascular complications (retinopathy and nephropathy) included mean HbA1c and diabetes duration; in addition, systolic blood pressure and smoking were independent predictors of nephropathy.

SD HbA1c and SD Glucose were independent predictors of microvascular complications (retinopathy and nephropathy) in type 2 (table 1), but NOT type 1 diabetes.

In type 2 (but NOT type 1) diabetes, the interaction between SD and mean HbA1c was significant for nephropathy and background retinopathy end points, indicating the effect of HbA1c was greatest at lower values of mean HbA1c; the interaction between SD and mean glucose was significant for nephropathy and background retinopathy endpoints.

	Type 2						Type 1					
	Retinopathy			Nephropathy			Retinopathy			Nephropathy		
	Background	Significant		Background	Significant		Background	Significant		Background	Significant	
Intercept	OR	P	OR	P	OR	P	OR	P	OR	P	OR	P
mHbA1c (%)	1.52	<0.001	2.60	<0.001	1.74	<0.001	1.50	0.007	2.05	<0.001	1.45	0.012
mGlc (mmol/L)	1.09	0.004	0.98	0.335	1.07	0.022	1.17	0.062	0.97	0.674	0.98	0.809
SD HbA1c (%)	2.09	0.024	11.37	<0.001	2.40	0.012	1.72	0.602	5.21	0.094	1.30	0.783
SD Glucose (mmol/L)	1.27	<0.001	1.26	0.001	1.54	<0.001	1.13	0.526	0.76	0.128	1.03	0.865
Diabetes duration (years)	1.07	<0.001	1.12	<0.001	1.07	<0.001	1.21	<0.001	1.31	<0.001	1.14	<0.001
Smoker	1.14	<0.001	0.98	0.506	1.16	<0.001	1.24	0.013	1.76	<0.001	1.21	0.012
Systolic BP (mmHg)	1.00	<0.001	1.00	0.001	1.02	<0.001	1.01	0.150	1.01	0.018	1.02	0.001
mHbA1c:SDHbA1c interaction	0.93	0.058	0.77	<0.001	0.91	0.025	0.93	0.502	0.86	0.136	1.03	0.803
mGlc:SDGlc interaction	0.98	0.004	0.99	0.090	0.98	<0.001	0.98	0.177	1.01	0.251	1.01	0.635

Table: The contribution of average and variability of HbA1c and glucose to the likelihood of retinal or renal endpoints in type 1 and type 2 diabetes patients. OR: odd ratio; SD: standard deviation, BP: blood pressure; mHbA1c: mean HbA1c; mGlc: mean glucose.

Discussion

Glycaemic variability is an independent predictor of microvascular complications in type 2 diabetes, but its predictive power diminishes with increasing average HbA1c and glucose.

Neither variability of glucose nor HbA1c significantly predicted microvascular outcomes in Emirati people with type 1 diabetes, despite higher variability of both than found in people with type 2 diabetes.

Acknowledgements

This study was supported by Imperial College London Diabetes Centre