Dr Asanga Fernando¹ (Consultant in Cancer Psychiatry, Clinical Director of Simulation), **Dr Chrishanthi Rajasooriyar**² (Consultant Clinical Oncologist), **Dr Nuradh Joseph**³ (Consultant Clinical Oncologist), **Dr Kesharie De Silva**⁴ (Consultant Anaesthetist), **Dr Kobika Sritharan**⁵ (Oncology research fellow), **Mr Christopher Broom**¹ (simulation specialist), **Dr Tim Benepal**¹ (Consultant Medical Oncologist)

1. St George's University Hospitals NHS Foundation trust. London. U.K. 2. Tellippalai Trail Cancer Hospital. Jaffna. Sri Lanka. 3. Ministry of Health. Sri Lanka. 4. Karapitiya Teaching Hospital. Galle. Sri Lanka. 5. The Royal Marsden NHS Foundation trust. London. U.K. (Contact: asanga.fernando@stgeorges.nhs.uk @asangafern)

A collaborative approach to evaluating the effectiveness of cross cultural training in psycho-oncology.

BACKGROUND & AIMS

A successful U.K designed inter-professional education programme at the interface of cancer, mental health and end of life care was adapted in collaboration with healthcare professionals in Sri Lanka to a series of half day workshops reflecting local priorities about management of mental illness co-morbidity in cancer patients. This study aims to evaluate the effectiveness of education programme by analysis of pre v post intervention changes to participant knowledge, confidence and attitudes to co-morbid mental disorder in cancer patients, in different environments across Sri Lanka.

METHODS

A series of three inter-professional workshops were conducted across Sri Lanka in March 2018, in Jaffna, Galle and at the National Cancer Institute, Maharagama with analysis of pre v post intervention questionnaires exploring participants' knowledge of assessing and managing co-morbid mental illness in cancer patients (by MCQ), participants' confidence at managing co-morbid mental disorder (by self rated 1-5 Likert scales), and participants' attitudes to co-morbid mental disorder in cancer patients (using questions from validated questionnaires).

RESULTS

(156 participants, 50% questionnaires included, n=78)

Qualitative feedback

- **Which two things have you learned today?** – Depression screening, Mx Depression, importance of EoLC
- **What is likely to Change your clinical Practice?** – Screening Head & Neck pts for depression, psych factors at End of Life

Knowledge based questions

- 4 questions
- Total 78 responses
- Increased from 31% to 74%
- Significant difference, p=0.0001

Confidence based questions

p=0.0001

Questions

- 11 Risk assessment in suicidal patient
- 12 Screening for depression
- 13 Managing an agitated patient
- 14 Managing a patient at the end of life
- 15 Breaking bad news

Significant improvement in confidence for each theme, p=0.0001, paired t-test

Attitude based questions

Have the participants changed their views towards mental health and patients with mental health problems?

5. If you were buying a house, would the presence of a mental health facility in the area you were planning to buy in, cause you to think twice about buying? YES/NO
6. Mental illness is an illness like any other YES/NO
7. If you were suffering from depression, would you feel comfortable talking with your superior about it? YES/NO
8. If you were buying a house, would the presence of a diabetes clinic in the area you were planning to buy in, cause you to think twice about buying? YES/NO
9. If you were suffering from a chest infection, would you feel comfortable talking with your superior about it? YES/NO
10. Are you likely to offer less attentive care to a patient with a history of mental illness in the context of the patient being violent and agitated? YES/NO

CONCLUSIONS

There were statistically highly significant changes post workshop intervention (compared to pre) with regards to participants knowledge (p=0.0001) of assessing and managing co-morbid mental disorder in cancer and end of life patients. There were also highly significant changes (p=0.0001) in participants' confidence at managing co-morbid mental health concerns in cancer and end of life patients. There were also interesting changes with regards to participants' attitudes to co-morbid mental disorder post intervention – however these were not statistically significant. Participants gained specific skills in assessment and management of depression in high risk groups relevant to the Sri Lankan context such as head and neck cancer patients. The study demonstrates that whilst this workshop was relevant in improving knowledge and confidence to managing mental health co-morbidity in the cancer and end of life care setting, attitudinal beliefs about mental disorder are harder to change and the reasons behind this require further study in a Sri Lankan context.