PREEMPTIVE ETORICOXIB ADMINISTRATION VERSUS FEMORAL NERVE BLOCK FOR ANALGESIA AFTER ANTERIOR CRUCIATE LIGAMENT RECONSTRUCTION

Florescu-Cionac, S; Anastase, DM; Munteanu, AM; Mihailide N; Mogos S
Orthopedic Clinical Hospital Foisor, BUCHAREST, ROMANIA

Background Reconstruction of the anterior cruciate ligament (ACLR) is associated with a considerable degree of postoperative pain.

Objectives To evaluate the efficacy and safety of preemptive administration of etoricoxib 120 mg versus femoral nerve block (FNB) with ropivacaine 0,5% 20 ml + xilocaine 1% 20 ml + dexamethazone 4mg on postoperative pain for the first 24 hours postoperatively.

Materials & Methods After Ethics Committee approval, 50 patients ASA I-II scheduled for ACLR under spinal anesthesia were randomized into two groups: the study group(Group A-Arcoxia), who received one tablet of etoricoxib 1 hour prior to surgery and the control group (Group B-ACLR) who received the FNB performed under ultrasound neurostimulation after regression of anesthesia. All patients received the same postoperative multimodal analgesia.

The cumulative morphine at 24 hours postoperatively to maintain a 10-point numerical pain rating scale value of 3 or less (NRS), the duration of analgesia from initiation of spinal anaesthesia until the first analgesic requirement and the incidence of side effects were recorded to 24 hours after surgery.

Summary Demographic data and surgical technique were similar between groups. At 24 hours postoperative analgesia was more effective in the FNB group ($p < 0.05$) with a significantly decrease of morphine consumption ($p < 0.05$) than in the etoricoxib group. There were no reported side effects.

Conclusions The preemptive administration of etoricoxib 120 mg was effective and safe for the patients with ACLR; the postoperative FNB significantly extends the duration of postoperative analgesia and decreases morphine consumption but not statistically significant.

Tabel nr.1 Demographic data			
Parameters	Group Arcoxia	Group ACLR	P
Age (mean±SD)	29 ± 8.4	30 ± 9.3	NS
Weight (mean±SD)	79 ± 12	76 ± 15	NS
Height (mean±SD)	176 ± 7.6	175 ± 11	NS
Sex F/M	19/6	18/7	NS

Tabel nr.2 Morphine requirements and time lasting analgesia			
Parameters	Group Arcoxia	Group ACLR	P t-Student
Morphine requirements mg (mean±SD)	16 ± 7.9	12 ± 10	NS
Lasting analgesia min (mean±SD)	736 ± 73	896 ± 283	0.0042

References

1. Graber R. Regional Anesthesia For Postoperative Pain Control. Drugs & Diseases > Clinical Procedures, Medscape, dec 2017

2. Golladay GJ, Balch KR, Dalury DF, Satpathy J, Jiranek WA. Oral Multimodal Analgesia for Total Joint Arthroplasty. J Arthroplasty. 2017 Sep;32(9S):S69-S73

3. Boonriong et al. Comparing etoricoxib and celecoxib for preemptive analgesia for acute postoperative pain in patients undergoing arthroscopic anterior cruciate ligament reconstruction: a randomized controlled trial. BMC Musculoskeletal Disorders 2010, 11:246