Exploring Factors Influencing of Lifestyle on Postoperative Health Related Life Quality of Colorectal Cancer Patients

S.P. Yang¹, T.J. Wang², C.C. Huang³, H.L. Chen⁴, M.H. Shen⁵

1. Cathay General Hospital, Nurse Practitioner of General Surgery, Taipei, Taiwan R.O.C.
2. National Taipei University of Nursing and Health Sciences, School of Nursing, Taipei, Taiwan R.O.C.
3. Cathay General Hospital, General Surgery, Taipei, Taiwan R.O.C.
4. Cathay General Hospital, Nursing, Taipei, Taiwan R.O.C.
5. Cathay General Hospital, Colorectal Surgery, Taipei, Taiwan R.O.C.

Background

Colorectal cancer is the third leading cause of cancer death and its incidence rate is continuously increasing. Tumor resection is primary treatment for colorectal cancer. Lifestyle, such as smoking, drinking, exercise and diet habits were related to the risk of death with colorectal cancer. However, only few studies have explored the impact of lifestyle on health related life quality of colorectal cancer patients. The purpose of the study is to explore influences of healthy lifestyle on health related life quality of patients with colorectal cancer.

Method

The study was based on a descriptive correlational design. A convenient sample of 106 colorectal cancer patients scheduled for cancer resection surgeries was recruited from general surgery or rectal surgery clinics of a medical center in Taipei, Taiwan. Data were collected with self-reported questionnaires, including questions on demographic, Charlson comorbidity index, the Alcohol Use Disorders Identification Test, Taiwan version of International Physical Activity Questionnaire, and the Eating at America's Table Study-Quick Food Scan. Data on disease profiles were collected from the patients' chart. Data on health-related quality of life were also collected at one month post-operation by using Functional Assessment of Cancer Therapy- Colorectal instruments version 4. Descriptive statistics were used to describe participants' demographics, disease profiles, and other study variables. The frequency, percentage, mean, and standard division were used to descriptive the central tendency and dispersion of distributions of the study variables. T-tests, one-way ANOVA and Pearson's correlation coefficient analysis Pearson's correlation coefficient was used to analyze the influencing factors of health related life quality. Multivariate linear regression analyses were used to analyze the important predictive factors of patients' quality of life at one month.

Results

The average score of the health related quality of life was 107.43 (SD = 13.95 ; the full score is 136). Studies showed worker's, level of education above college, good of economic, surgery of laparoscopic approach, albumin level ≥ 3.5 g/dl and small amount of alcohol drinking had significantly higher health-related quality. Results of colorectal cancer patients' exercise was positive correlation with health related quality of life. Age, CCI and postoperative length of hospital stay were negative correlation with health related quality of life. Linear regression showed that age, the mean postoperative length of hospital stay and education level were the significant predictive factors of health related quality of life.

Conclusion

The results of the study supported that colorectal cancer patients who had more exercise also reported better health related quality of life. And patients who were elder, longer postoperative length of hospital stay, and below college level of education reported lower health-related quality of life . Therefore colorectal cancer patients were at risk of lower health related quality of life after colectomy should be evaluated closely before operation. The study inform us the important influencing factor of the patients' life quality as well as provide a foundation for developing interventions to decrease post-operative complications and promote health related quality of life.

Table 1.

Age , CCI, Smoking, Drinking, Exercise, Intake amount of fruit and vegetable, postoperative length of hospital day and health related quality of life (n=106)

Variables	Age	CCI	Smoking	Drinking	Exercise	Fruit & Vegetable	Length of hospital day	Quality of life
Age	1							
CCI	.274**	1						
Smoking	-.138	-.045	1					
Drinking	-.182	-.133	.439**	1				
Exercise	-.253	-.221*	-.025	.067	1			
Fruit & Vegetable	-.138	-.022	-.124	-.044	.260**	1		
Length of hospital day	.332**	.307**	.146	-.013	-.226*	-.067	1	
Quality of life	-.435**	-.215*	.050	.142	.277**	.110	-.358**	1

Note: Pearson's correlation, *. Correlation is significant at the 0.05 level (2-tailed).; ** Correlation is significant at the 0.01 level (2-tailed)

Table 2.

A important predictor of health-related quality of life for one month after surgery

Variables	Beta	t	p	Adjust R ²	F	VIF
Model				0.257	5.27	
1. Age	-0.27	-2.634	0.010			1.486
2. Postoperative length of hospital day	-0.23	-2.506	0.014			1.134
3. Education level						
High school VS elementary school	0.03	0.261	0.795			1.491
College VS elementary school	0.23	2.042	0.044			1.701