Attracting earlier referrals to a paediatric palliative care service: the vocabulary of enhanced supportive care

Hemsley J¹, Dinsdale A¹, Bayliss J¹, Wood A¹, Walters L¹, Downie J¹, Vemuri S¹, Barone G¹, McCulloch R¹, Henderson E¹, Stephenson A¹, Bluebond-Langner M^{1,2}. Great Ormond Street Hospital¹ Institute of Child Health²
Correspondence: june.hemsley@gosh.nhs.uk

Background

- Previous studies have shown cancer patients benefit from earlier referral to palliative care services^{1,2}.
- Resistance amongst cancer specialists to refer before curative treatment options are exhausted
- An NHS England initiative in adult cancer services aimed to improve early referral by implementing a framework of enhanced supportive care (ESC)³
- The subject - a tertiary level children's hospital with cancer services and a dedicated specialist palliative care service - agreed to trial this initiative within a paediatric cohort.

Aim and objectives

Increase rate of early referrals by cancer teams through

- Implementing a shared vocabulary of 'ESC' and 'poor prognosis' (table 1) to facilitate earlier referrals.
- Offering an ESC referral to patients with a poor prognosis within one week of patient and family being informed of prognosis.

Methods

- The vocabulary of ESC was integrated into the Principle Treatment Centre (PTC)
- An agreed definition of "poor prognosis" was developed in discussions between palliative care and cancer specialists.

Table 1. Definition of poor prognosis

Speciality	CancerType	Point of disease trajectory
Neuro-oncology	DIPG, HGG, HR MBL, ATRT, ETANTR, metastatic Ependymoma, Anaplastic astrocytoma grade III+	At diagnosis
	All brain tumours	1 st relapse / progression / refractory disease
Oncology	Rhabdoid, metastatic RMS	At diagnosis
	RMS, HR HBL, HR NBL (poor prognosis at day 80)	1 st relapse / progression / refractory disease
Haematology	Therapy-related AML	At diagnosis
	Burkitt's lymphoma, diffuse large B cell lymphoma, haematology post BMT	1st relapse / progression / refractory disease
All specialities	All malignancies	Multiple relapse / progression / persistent refractory disease (excl. LCH)
		Enrolment in phase I/II trial
		Commencement of medications for compassionate use
		Patient's lead H/O consultant confirms poor prognosis
	All malignancies with a confirmed life-limiting co-morbidity	Diagnosis of malignancy or life-limiting co-morbidity (whichever is later)

References:

1. Wiener et al (2015) Deeper conversations need not wait until the end, JCO, 33 (33) pp 3974;
2. Beecham et al (2016) Keeping all options open: parents' approaches to Advance Care Planning, Health Expectations, DOI: 10.1111/hex.12500
3. <https://www.england.nhs.uk/wp-content/uploads/2016/03/ca1-enhncd-supprtv-care-guid.pdf>

Defining Enhanced Supportive Care (ESC) in paediatric cancer care

- Symptom management and focus on quality of life
- In parallel with disease-directed treatment
- Poor prognosis or high symptom burden (>grade 2)

Table 2. Symptom assessment criteria

Symptom	Grade 0	Grade 1	Grade 2	Grade 3	Grade 4
Pain	Nil pain	Mild pain	Moderate pain; limiting instrumental ADL	Pain at rest; limiting self-care ADL	
Peripheral neuropathy	Nil peripheral neuropathy	Asymptomatic; loss of deep tendon reflexes or paraesthesia ; clinical or diagnostic observations only; intervention not indicated	Moderate pain; limiting instrumental ADL	Severe symptoms; limiting self-care ADL; assistive device indicated	Life-threatening consequences; urgent interventions indicated
Nausea	Nil nausea	Loss of appetite without alteration in eating habits	Oral intake decreased without significant weight loss, dehydration or malnutrition	Inadequate oral caloric or fluid intake; tube feeding, TPN or hospitalization indicated	
Vomit	Nil vomiting	1-2 episodes (separated by 5 minutes) in 24 hours	3-5 episodes (separated by 5 minutes) in 24 hours	>=6 episodes (separated by 5 minutes in 24 hours: tube feeding, TPN or hospitalization indicated	Life-threatening consequences; urgent interventions indicated
Agitation	Nil agitation	Mild mood alteration	Moderate mood alteration	Severe agitation; hospitalization not indicated	Life-threatening consequences; urgent interventions indicated
Breathlessness	Nil breathlessness	Shortness of breath with moderate exertion	Shortness of breath with minimal exertion; limiting instrumental ADL	Shortness of breath at rest; limiting self-care ADL	Life-threatening consequences; urgent interventions indicated
Constipation	Bowels open normally	Occasional or intermittent symptoms; occasional use of stool softeners, laxatives, dietary modification or enema	Persistent symptoms with regular use of laxatives or enemas; limiting instrumental ADL	Obstipation with manual evacuation indicated; limiting self-care ADL	Life-threatening consequences; urgent interventions indicated

Results

- Of the 36 patients identified with a new poor prognosis between October 2017 and March 2018, 27 (75%) were offered an ESC referral within one week.
- Of 6 patients identified with a high symptom burden between January 2019 and March 2019, 5 (83%) were met within 2 working days; 1 proceeded to palliative care.

Conclusion

- Attracting earlier referrals to a paediatric palliative care service is feasible through a change of vocabulary and reconfiguration of service.
- Next steps: consider applicability to patients with non-malignant life-limiting conditions.