ESP18-0609

SUPPURATIVE OTITIS MEDIA IN CHILDREN IN ANGOLA

M. Filipe¹, Å. Reimer², F. Uddén³, T. Pelkonen⁴, K. Riesbeck³.

¹ Department of ENT, Faculty of Medicine, University Agostinho Neto, Luanda, Angola.

² Näshälsan Höllviken AB, Höllviken, Sweden.

³ Clinical Microbiology, Dept. of Translational Medicine, Faculty of Medicine, Lund University, Malmö, Sweden.

⁴ Helsinki University Hospital, Children's Hospital, Helsinki, Finland and University of Helsinki, Helsinki, Finland.

Background: Chronic suppurative otitis media (CSOM) is an important cause of hearing loss in children and constitutes a serious health problem globally. We studied a cohort of children with otorrhoea in five provinces of Angola, a Sub-Saharan country.

Methods : Children attending services of oto-rhino-laryngology due to otorrhoea were submitted to full clinical examination and interview. Ear secretion and nasopharyngeal samples were collected, transported frozen, and cultured in Sweden using standard techniques

Results : Unilateral disease was detected in 76%, bilateral in 24% of children. Otorrhoea had lasted <15 days in 24%, and ≥ 1 year in 39%. Clinical hearing loss presented in 49% of children (96 of 196), and correlated with the duration of otorrhoea ($p = 0.003$). When comparing with the capital province Luanda, a poorer Lunda sul province, otorrhoea had lasted longer and hearing loss was more common ($p=0.02$). Ear discharge specimens ($n=133$) and nasopharyngeal swabs ($n=108$) yielded 376 and 212 individual isolates, respectively. *Proteus* spp. (13.3%), *Pseudomonas aeruginosa* (10.4%) and *Enterococcus* spp. (10.4%) were dominating in ear discharge. Pneumococci and *Staphylococcus aureus* were detected in 15.1% and 7.1% of nasopharyngeal samples, respectively. Resistance rate to quinolones exceeded 10% among *Enterobacteriaceae* and was 30.8% in *Staphylococcus aureus*, whereas 6.3% of *P. aeruginosa* was resistant.

Conclusions

- In Angola, otorrhoea is long-lasting, causes hearing loss, and associates with resource-limited living conditions.
- Bacterial cultures of ear discharge reveal difficult-to-treat Gram-negative bacteria.
- Ear infections should be treated early to avoid severe sequelae.