Incidence and risk factors associated with Postoperative Delirium

Norte G¹, Bento M¹, Vieira Marques F¹, Moreira M¹, Martins M¹, Matos F¹
¹Coimbra Hospital and University Centre, Dept. of Anaesthesiology, Coimbra, Portugal

Background and Goal of Study:

Postoperative delirium (POD) is characterized by an acute change in cognitive function after general anaesthesia and its prevalence can range from 5 to 51%. In this study, we assessed the incidence, risk factors and outcome of POD.

Materials and methods:

Prospective study, August 2015	4AT scale: <ul style="list-style-type: none">• Transfer to OR• Admission in PACU• Discharge from PACU• 24h after surgery Richmond Agitation-Sedation Scale (RASS): <ul style="list-style-type: none">• Hypoactive• Hyperactive• Sedation
Inclusion criteria: <ul style="list-style-type: none">• Elective surgery• General anaesthesia• > 18 yr-old• Admission to PACU Exclusion criteria: <ul style="list-style-type: none">• Severe dementia• Unable to communicate• Anoxic cerebral lesions• Neuromuscular disease	
Statistical analysis: X ² test, Mann-Whitney and Kruskal-Wallis tests.	

Results and discussion:

Table 1 – Description of the characteristics of the sample (using the median and interquartile range whenever possible). Comparison of the total sample characteristics with the characteristics of patients who developed delirium and those who did not develop delirium, regardless the study phase, with respective p-value. Assessment of delirium with 4AT scale.				
Variable	Total sample (n = 151)	With delirium (n = 28)	Without delirium (n = 123)	P-value
Age	63 (19, 93)	71.5 (29, 93)	62 (19, 93)	0.001
Gender, men	73 (48.34%)	18 (64.29%)	55 (44.72%)	0.061
ASA Classification	2 (1, 4)	2 (1, 3)	2 (1, 4)	0.045
1	8.6 %	3.6 %	9.8 %	
2	60.3 %	50 %	62.6 %	
3	30.5 %	46.4 %	26.8 %	
4	0.7 %		0.8 %	
Comorbidities				
Cardiovascular	72.8 %	89.3 %	69.1 %	0.030
Diabetes	29.1 %	39.3 %	26.8 %	0.190
Neurologic	7.3 %	10.7 %	6.5 %	0.429
Respiratory	23.8 %	21.4 %	24.4 %	0.740
Oncologic	24.5 %	32.1 %	22.8 %	0.298
Renal	8.6 %	21.4 %	5.7 %	0.016
Hepatic	3.3 %	3.6 %	3.3 %	1.000
Cognitive deficits	5.3 %	10.7 %	4.1 %	0.167
Anaesthesia duration (min)	150 (45, 420)	180 (45, 420)	150 (50, 360)	0.063
Modified Aldrete score				
Admission to PACU	12 (6, 14)	12 (6, 13)	12 (8, 14)	<0.001
Discharge from PACU	14 (12, 14)	14 (13, 14)	14 (12, 14)	0.065
Qualitative Pain scores				
Admission to PACU	2 (0, 4)	1.5 (0, 3)	2 (0, 4)	0.684
Discharge from PACU	0 (0, 3)	0 (0, 2)	0 (0, 3)	0.471
Infirmery	1 (0, 4)	1 (0, 3)	0 (0, 4)	0.504

References:

Inouye SK, Westendorp RG, Saczynski JS. Lancet 2014; 383:911-22

Table 2 – Factors associated with development of POD at all study phases.

	PACU admission	PACU discharge	Infirmery
Variable	P-value	P-value	P-value
Age	0.002	0.006	0.013
Modified Aldrete score at PACU admission	<0.001	0.001	0.001
Comorbidities			
Renal	0.042		
Cardiovascular	0.018		
Cognitive deficits		0.016	
Diabetes		0.024	0.023
Intraoperative metoclopramide	0.029		
ASA classification		<0.001	0.003

Table 3 – Signs of delirium assessed with 4AT scale.

RASS scale	n = 151
Signs of delirium at PACU admission	25 (16%)
Hypoactive	88%
Hyperactive	12%
Signs of delirium at PACU discharge	7 (4,6%)
Hypoactive	86%
Hyperactive	14%
Signs of delirium at Infirmery	7 (4,6%)
Hypoactive	86%
Hyperactive	14%

Conclusions:

- POD is common and with highest incidence at PACU admission;
- Hypoactive features are more prevalent;
- Marked decrease of POD during PACU stay, although some patients have delirium signs after discharge from PACU;
- No association with opioids or anaesthetic drugs;
- Routine delirium monitoring at PACU and control of comorbidities are fundamental