Haemodynamic monitoring with ECOM in elective off-pump coronary artery bypass grafting: A case-control study

Leclercq T., Schulz T., Meyer A., Lilot M., Fellahi J.-L.
Department of Anaesthesiology & Intensive Care, Louis Pradel Hospital, Lyon, France

Background: Perioperative haemodynamic optimization decreases morbidity and hospital length of stay following cardiac surgery. Off-pump coronary artery bypass grafting (OPCAB) is a simplified procedure which could accelerate postoperative recovery and limit admission to the intensive care unit (ICU). The Endotracheal Cardiac Output Monitor (ECOM) provides continuous cardiac output measurement *via* endotracheal bioimpedance in conjunction with an arterial catheter . We aimed to demonstrate that ECOM could improve haemodynamic management and shorten recovery following OPCAB surgery.

Materials and methods: After approval of local Ethical Committee, we prospectively included 20 consecutive adult patients scheduled for elective OPCAB in the Teaching University Hospital Louis Pradel (Lyon, France) and receiving haemodynamic monitoring with ECOM (ECOM group). These patients were compared to a retrospective cohort of 42 adult patients scheduled for elective OPCAB and not receiving haemodynamic monitoring with ECOM (Control group). The primary endpoint was the rate of postoperative admission to the ICU. Secondary outcomes included the time to extubation, the length of stay in ICU, vasopressor requirements, and postoperative measurements of serum troponin and lactate concentration. We also assessed the feasibility of ECOM monitoring in that setting.

Results and discussion: The rate of postoperative admission to the ICU was 38/42 (90%) in the Control group vs. 11/20 (55%) in the ECOM group, $P=0.008$ (Table below). None unexpected admission for haemodynamic instability was observed in the ECOM group vs. 12 in the Control group. All secondary outcomes are reported in Table 1. On a scale ranging from 0 to 5, the easiness of ECOM use and the degree of physicians' satisfaction were 4.30 ± 1.17 and 3.45 ± 0.68 , respectively.

	Control(n=42)	ECOM(n=20)	P value
Admission to the ICU	38 (90%)	11 (55%)	0.008
Expected	24	10	
Unexpected	14	1	
Extubation time (h)	4 [0-696]	2 [0-7]	<0.001
Length of stay in ICU (days)	1.0 [0 - 29]	0.5 [0 - 10]	0.002
Vasopressor requirement	27 (64%)	19 (95%)	0.012
Troponin at admission (ng/ml)	944 [265-1518]	546 [0-275]	0.028
Lactate at H6 (mmol/l)	2.0 ± 0.6	1.5 ± 0.4	0.043

Conclusion: The systematic use of ECOM in OPCAB surgery is associated with a significant reduction in the rate of admission to the ICU and lower time to extubation and length of stay, suggesting an increase in haemodynamic optimization. Further studies are mandatory to confirm the interest of ECOM in improving postoperative outcomes in cardiac surgery.