Observational acute pain study in admitted patients of a tertiary hospital in Spain.

Rovira L.¹, Alcover L.¹, Mazzinari G.¹, Collado B.¹, Madrid E.², Romero E.¹ Hospital de Manises, Dept of Anaesthesiology & Pain Medicine, Manises, Spain.

METHODS

An observational cross-sectional study was performed for 7 consecutive days to all patients admitted to a tertiary hospital.

Acute pain was collected by visual analog scale (VAS) at rest and in motion, every 8 hours.

Other variables of interest as the specific analgesic treatment (locoregional techniques), rescue analgesic medication administered, as well as sex, age, reason for admission, and specialty care was collected.

RESULTS

- 246 patients (149 surgical and 97 medical) were evaluated, the prevalence of episodes of moderate to severe pain (VAS> 4) was 53% (132/246) of all patients admitted.
- The average number of episodes (VAS> 4) occurred 1.45 ± 2.0 times.
- Rescue medication was administered to 48% of the observed patients. Being acetaminophen the most used drug.

CONCLUSION

- Pain assessment should be considered as the fifth leading vital sign and it should be routinely evaluated.
- The percentage of patients with moderate-severe pain is unacceptable and more measures are needed to meet the goal of Pain Free Hospital.

REFERENCES

- Cano JM, de Juan S. Pain evaluation as the fifth vital parameter. Med Clin (Barc). 2007 Feb 3;128(4):159.
- Sommer M, de Rijke JM, et al The prevalence of postoperative pain in a sample of 1490 surgical inpatients. Eur J Anaesthesiol. 2008 Apr;25(4):267-74.
- Joint Commission on Accreditation of Healthcare Organizations. Implementing the new pain managements standards. Oakbrook Terrace: JCAHO;2000.

Prevalence of pain

	Episodes VAS>4 (mean)	[IC95]	N
Surgical patient	1.83	[0-7]	149
Medical patient	0.90	[0-4]	97
Total	1.46	[0-5.4]	246