A CASE OF CLOZAPINE-INDUCED AGRANULOCYTOSIS IN TREATMENT RESISTANT SCHIZOPHRENIA

CATIVO, Catarina; SERRANO, Raquel; GONÇALVES, Patrícia; COSTA-PEDRO, João; DIAS, Manuel; BORJA-SANTOS, Nuno; MAIA, Teresa.

Department of Psychiatry, Hospital Prof. Dr. Fernando Fonseca, Amadora, Portugal.

INTRODUCTION

Clozapine is a multireceptorial atypical antipsychotic, considered to be more effective than other antipsychotics in the management of treatment-resistant schizophrenia (TRS). Nevertheless, it's generally under-used, due to concerns regarding it's adverse effects. Although uncommon (0.8-3%), the most severe adverse effect is clozapine induced neutropenia (CIN), that may lead to clozapine induced agranulocytosis (CIA) which is defined by an absolute neutrophil count below 500 /mm³ and may have fatal consequences.

The authors aim to Report the case of a patient with TRS who developed agranulocytosis after starting treatment with clozapine and discuss the management of CIA.

MATERIALS AND METHODS

Case report and non-systematic review through literature research in PubMed/Medline database, using the key-words: agranulocytosis and clozapine.

RESULTS

CASE REPORT

27-year-old male, completed high school in a professional course program, unemployed for 6 years.

- ✓ **No** medical comorbidities
- ✓ **No** substance abuse
- ! **Mother** diagnosed with **Schizophrenia**

! Patient diagnosed with **Schizophrenia 1 year and 3 months prior**. Since then **2 hospital admissions**.

🧪 During this period, he was successively medicated with:

| **Aripiprazole** 30mg | **Olanzapine** 20mg +, **Zuclopenthixol decanoate** 200mg |
| Zuclopenthixol decanoate 400mg bi-monthly | **Paliperidone** 18mg (later replaced by monthly 100 mg **Paliperidone depot**) + **Clozapine** titulated up to 300mg |
With only **partial remission**. Serial **hemogram monitoring** was always **normal**.

🏠 **2 months** after, he **abandoned all medication** and was hospitalised for the 3rd time, with recrudescence symptoms of social withdrawal, mystical delusions, passivity of volition and auditory verbal hallucinations.

Paliperidone 18mg and **Clozapine** 350mg were gradually re-introduced with some improvement. **Weekly hemogram tests** were **normal**.

LITERATURE REVIEW

CIN - <1500 neutrophils/mm³ - and **CIA** - <500 neutrophils/mm³), are uncommon adverse effects of clozapine treatment, with a cumulative risk of **3%** and **0.8%**, respectively.¹

The risk is **not dose-dependent**, it is **highest in the first 4-20 weeks** of treatment, decreasing sharply after 1 year, but they **can occur at any time** during treatment (idiosyncratic effect).⁴

Regarding **CIA**, there usually is a normal neutrophil count that abruptly falls for 1 week, to a value near 0.00.¹

Haematological monitoring is mandatory to mitigate the CIN/CIA risk.⁴

	USA's REMS Program		European Medicines Agency	
Monitoring*	★ <i>First 6 months:</i> weekly ★ <i>6-12 months:</i> every 2 weeks ★ <i>>1 year:</i> monthly		★ <i>First 18 weeks:</i> weekly ★ <i>>18 weeks:</i> monthly	
	Must continue throughout the duration of treatment and up to 4 weeks after discontinuation.			
ANC	Action	Monitoring*	Action	Monitoring*
1500-2000	<u>Continue treatment</u>	Keep initial schedule	<u>Continue treatment</u>	Twice a week
1000-1500		3 times a week	<u>Stop clozapine</u> Do not reintroduce	Daily
500-1000	<u>Stop clozapine</u> Re-introduce after reaching >1000	Daily until >1000 3 times a week until >1500		
<500 (CIA)	<u>Stop clozapine</u> May be reintroduced if beneficial	Then weekly for 1 month		

Table 1 - Comparison between american and european guidelines for monitoring CIN and CIA. (ANC - Absolute Neutrophil Count, *blood sample with leukocytes count.

In case of CIN/CIA, symptoms and signs of infection should be monitored and infection should be aggressively treated.

Neutrophils usually **return to normal range 1-3 weeks** after stopping clozapine.¹

Drugs like **lithium** as an effort to increase leukocyte counts are **not recommended**.⁴

G-CSF - filgrastim, can potentially **accelerate recovery times** in patients with CIA and secondary infection. It is not yet approved as a co-adjuvant for the reintroduction of clozapine in CIA patients.⁴

CONCLUSION

Despite serious adverse effects like CIN and CIA, clozapine remains the gold standard for managing TRS, with few other efficient treatment alternatives.

Therefore it is of great importance to monitor neutrophil counts closely according to the most recent guidelines, even if it is difficult to do so outside of hospital setting, due to patient non-compliance with repeated blood sampling.

Ultimately this will allow for early identification of these complications, better management and increased safety when prescribing clozapine.

REFERENCES

1 - Wicnski M. et al. (2018) Clozapine-induced agranulocytosis/granulocytopenia: mechanisms and monitoring. Curr Opin Hematol. 2018 Jan;25(1):22-28.
2 - Li K. et al. (2018) Clozapine pharmacogenomics: a review of efficacy, pharmacokinetics, and agranulocytosis. Curr Opin Psychiatry 31(5):403-408.
3 - De Fazio et al. (2015) Rare and very rare adverse effects of clozapine. Neuropsychiatric Disease and Treatment.

4 - Taylor, D. et al. (2018). Maudsley Prescribing Guidelines, 13th Edition. Wiley Blackwell Publishing.
5 - European Medicines Agency (2002) Leponex - clozapine background information. London. In: <https://www.ema.europa.eu/en/medicines/human/referrals/leponex> Accessed 31-07-19
6 - Clozapine REMS. (2014) Clozapine and the Risk of Neutropenia: An Overview for Healthcare Providers. In: <https://www.clozapinerems.com/> Accessed in 31-07-19