Cow's milk protein allergy in a preterm infant, case report

¹Firat Ozkececi, ²Turan Derme, ²Belma Saygili Karagol

¹University of Health Sciences, Gulhane Medical Faculty, Department of Pediatrics, Ankara, Turkey

²University of Health Sciences, Gulhane Medical Faculty, Department of Pediatrics, Division of Neonatology, Ankara, Turkey

Introduction: Cow's milk protein allergy (CMPA) is the most common cause of allergy occurring in the first year of life due to infant formula or breast-milk of mothers who are drinking cow's milk or eating cow's milk products. Most cases are symptomatic in the first months and the most common clinical signs are bloody stool in term and well appearing infant. On the other hand, in a preterm infant with rectal bleeding, necrotizing enterocolitis (NEC) is considered at the first sight. Here we report a preterm infant of CMPA with initially considered as NEC.

Case: A 800 g female preterm infant was born by vaginal delivery at 28 weeks to a 24-year-old gravid 2, para 1 mother. The neonate borned and she did not cry spontaneously. Apgar scores were 2/3/6 at 1/5/10 minutes. The infant was intubated in the delivery room and admitted to the neonatal intensive care unit. Trophic feeds were started on the fifth day of life with breast milk of mother. On the tenth day of life abdominal distension was developed. There is no signs of NEC on X-ray. Since the baby was clinically well we suspected cow's milk-induced colitis and continued feeding with amino acid-based formula (Neocate). After feeding with aminoacid-based formula clinical signs were delayed. The baby was discharged on aminoacidbased formula.

Result: CMPA is one of the causes of colitis in infants. CMPA should be kept in mind especially if the baby is well-appeared and has normal abdominal X-rays. Generally diagnosis is based on response to elimination diet and recurrence with reintroduction of cow's milk protein in the diet.

<https://mypositiveparenting.org/ms/2016/06/26/alergi-protein-susu-lembu/>