Accurate Cultures for Gonorrhoea. How do Community Services and Secondary Care Compare?

Tom Board, Sue Mitchell, Judy Russell

Introduction:

BASHH guidelines emphasise the importance of accurate cultures in the diagnosis and management of gonorrhoea (1). This audit's aim was to establish if there was a measurable difference in the positive culture yield between community and secondary care services in the months following a change in contract which has moved a proportion of walk in patients from secondary care to the community setting.

Each service used a combination of culture and NAATs testing to make the diagnosis. It is known that cultures offer greater specificity and also give information regarding antibiotic sensitivity. This is an important public health parameter.

Therefore accurate culturing is important.

'This is a delicate and fastidious organism which dies rapidly if exposed to desiccating or oxidising conditions and requires a moist carbon dioxide enriched atmosphere and a nutrient medium if it is to be cultured successfully.' (2)

Methods:

Relevant databases were searched for gonorrhoea patients after 1st October 2017 when the service was changed. A retrospective audit of the notes was then carried out to establish the rates of positive NAATS tests and positive culture yield and compared the two services.

Results:

In secondary care 25 patients (20 men & 5 women) were treated for gonorrhoea over two months. In these patients 22 had a positive NAATs test, of these 17 had a positive culture giving a positive culture yield of 77.2%. In comparison the community service identified 10 patients in the first month of the new contract with a positive NAATs test: the NAATs/ culture concordance rate was 11%."

Limitations:

This represents the first two months of data following a significant reorganisation of local services. The community services are making changes to deliver samples to the laboratory in a more timely fashion and preliminary figures show a degree of improvement. Further audits are taking place.

Discussion:

Accurate cultures are vital for the treatment of gonorrhoea, particularly in areas where antibiotic resistance is high. Appropriate storage and prompt processing is important to ensure the viability of these tests.

A contracted responsibility of the specialist GUM service is "clinical leadership and governance" across the providers which should enable the appropriate changes in practice to maximise culture success to be put in place.

Conclusion

This audit raises a question about viability of *Neisseria gonorrhoeae* during transfer from community settings to the central hospital laboratory.

1. Bignell C, Fitzgerald M. UK national guideline for the management of gonorrhoea in adults. Int J STD AIDS. 2011; 22(10): 541-7.
2. Jephcott A. Microbiological diagnosis of gonorrhoea. Genitourin Med. 1997; 73: 245-252