Hypoxic Ischemic Encephalopathy and Heart Rate Variability in Neonates: A Systematic Review

Mads Andersen, MS; Ted C. K. Andelius, MD; Mette V. Pedersen, MS; Kasper J. Kyng, MD, PhD; Tine B. Henriksen, MD, PhD
Department of Pediatrics, Aarhus University Hospital, Denmark

Aim

To systematically review whether the grade of hypoxic ischemic encephalopathy (HIE) is associated with heart rate variability (HRV) in neonates.

Methods

- The review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).
- We searched Pubmed, Embase, Web of Science, Cochrane Database (CENTRAL), and Scopus.
- Two reviewers independently performed selection of articles and the strength of evidence assessment by the Grading of Recommendations Assessment, Development and Evaluation (GRADE).
- Neonates with moderate or severe HIE were compared with neonates with no or mild HIE with respect to HRV measures assessed within seven days after birth.
- HRV measures considered:
 - Time domain measures:** standard deviation of RR-intervals (SDRR) and standard deviation of normal-to-normal intervals (SDNN) (Figure 2).
 - Frequency domain measures:** low frequency power (LF) and high frequency power (HF) in normal units (nu) (Figure 3).

Results

For every measure, the strength of evidence was very low. Three studies found a reduction in time domain measures, while three studies found a mean difference in frequency domain measures.

Figure 1. PRISMA flowchart of the study selection

Figure 2. Time domain measures

Figure 3. Frequency domain measures

Source	Time of measure	HRV measure	Mean difference with 95 % CI
--------	-----------------	-------------	------------------------------

Severe vs. mild HIE

Goulding et al. 2015	12-48 hours	SDNN (ms)	-10.1 (-15.8, -4.4)
Goulding et al. 2017	6-72 hours	SDNN (ms)	-10.3 (-14.8, -5.8)

Moderate vs. mild HIE

Goulding et al. 2015	12-48 hours	SDNN (ms)	-7.4 (-12.8, -2.0)
Goulding et al. 2017	6-72 hours	SDNN (ms)	-5.3 (-11, 0.4)

Severe vs. no HIE

Aliefendioglu et al. 2012	End of first postnatal week	LF (nu)	-45.3 (-53.8, -36.8)
		HF (nu)	28.6 (33.5, 23.7)

Moderate vs. no HIE

Aliefendioglu et al. 2012	End of first postnatal week	LF (nu)	-15.3 (-21.3, -9.3)
		HF (nu)	10.3 (14.8, 5.8)

Severe and moderate vs. mild and no HIE

Vergales et al. 2013	First 24 hours	SDRR (ms)	-9.2 (-16.5, -1.9)
----------------------	----------------	-----------	--------------------

Conclusion

More severe grades of HIE were associated with reduced time domain measures and LF (nu), and increased HF (nu). However, the strength of evidence was very low.