ESR1 hotspot mutations in endometrial stromal sarcomas may confer hormonal resistance

Kimberly Dessources¹, Arnaud Da Cruz Paula¹, Simon Lee², Youran Zou², Robert A Soslow², Martee L Hensley³, Britta Weigelt¹ and Sarah Chiang²

Departments of ¹Surgery, ²Pathology and ³Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

BACKGROUND

- Endometrial stromal sarcomas are a rare subset of uterine mesenchymal tumors, accounting for 0.2% of all uterine malignancies.
- Low-grade endometrial stromal sarcomas (LGESS) may show high-grade morphology in primary or recurrent tumors.
- These lesions are classified as high-grade endometrial stromal sarcomas (HGESS), which in general are more aggressive and have higher rates of resistance to endocrine therapy than LGESS.
- The pathogenesis of hormonal resistance in these tumors has yet to be defined.

AIMS

- Characterize the mutational profiles of primary LGESS and associated recurrences with high-grade morphology following endocrine therapy.

METHODS

- DNA from tumor and normal tissues was extracted from all samples from case 1 and case 2, summarized in Table 1.
- DNA was subjected to our institutional massively parallel sequencing platform targeting 468 cancer genes (MSK-IMPACT).
- RNA from each case was also evaluated for the presence of gene fusions using ARCHER FusionPlex.
- Sequencing data were analyzed using validated bioinformatics algorithms.

RESULTS

- Both patients received at least two lines of hormonal suppressive therapy including letrozole and megestrol.
- ARCHER FusionPlex analysis revealed that cases 1 and 2 harbored *JAZF1-PHF1* and *EPC1-PHF1* fusions, respectively.
- Targeted massively parallel sequencing revealed that these lesions harbored few non-synonymous somatic mutations.
- The primary LGESS and the two HGESS recurrences of case 1 shared a *KDM5C* frame-shift mutation; however in the second HGESS recurrence post endocrine treatment, a clonal *STK40* (R128W) mutation and an *ESR1* (Y537S) hotspot mutation were detected.
- Both LGESS and HGESS recurrences of case 2 harbored an *ESR1* (Y537S) hotspot mutation. In addition, an *HRAS* (Q61R) hotspot mutation restricted to the recurrent LGESS was identified.
- ESR1* hotspot mutations have been shown to play a role in acquired endocrine therapy resistance in breast cancer.

Table 1: Case Summaries			
Year	Sample ID	Pathology	Treatment
Case 1 – JAZF1-PHF1 fusion			
2006	ESR1-1-A	Primary: LG-ESS, ER+ PR+	Megestrol
2012	N/A	Recurrence #1: HG-ESS, ER+ PR+	Arimidex
2015	ESR1-1-B	Recurrence #2: HG-ESS, ER+ PR+	Letrozole
2017	ESR1-1C	Recurrence #3: HG-ESS, PR diffuse+; ER weak +; ESR1 Mutation +	Fulvestrant
Case 2 – EPC1-PHF1 fusion			
2013	N/A	Primary: LG-ESS, ER+ PR+	Letrozole→ Megestrol →Docetaxol/ge mcitabine
2016	ESR1-2	Recurrence #1: LG-ESS, ER+ PR+, ESR1 Mutation +	
2017	ESR1-4	Recurrence #2: HGESS ,ER+ PR+, ESR1 Mutation +	Doxorubicin→ Fulvestrant

CASE 1

Figure 1: Repertoire of somatic mutations in endometrial stromal sarcoma case 1. (Left) Non-synonymous somatic mutations identified in the primary tumor (A), the recurrence 9 years (B) and 11 years (C) after initial diagnosis. (Right) Cancer cell fractions of the mutations identified, color-coded according to the legend. Note the clonal *ESR1* hotspot mutation is present only in last recurrence.

CASE 2

Figure 2: Repertoire of somatic mutations in endometrial stromal sarcoma case 2. (Left) Non-synonymous somatic mutations identified in the recurrences 3 years (2) and 4 years (4) after initial diagnosis. (Right) Cancer cell fractions of the mutations identified, color-coded according to the legend. Note the clonal *ESR1* hotspot mutation was present in all recurrences. Material from the primary tumor was not available for analysis.

CONCLUSION

Our findings suggest that *ESR1* hotspot mutations in LGESS, either pre-existing or acquired, may be associated with endocrine resistance and high-grade transformation in ESS.