Evaluating post-traumatic headache in patients with mTBI in acute and chronic: Resting-State fMRI identifies unique network changes

Jordan Lemme^{1,2,3*}, Scott Holmes^{1,2}, Diana Sibai^{1,2}, Jaymin Upadhyay^{1,2}, David Borsook^{1,2}

¹Center for Pain and the Brain, Department of Anesthesiology, Boston Children's Hospital; ²Harvard Medical School; ³University of New England College of Medicine

Background

- Posttraumatic Headaches (PTH) are a common sequelae from mild traumatic brain injuries (mTBI) with symptoms including headaches, fatigue, dizziness, memory and concentration difficulties.
- PTH typically resolves within 3 months, however a subgroup of individuals will experience PTH >3 months.
- It remains unclear if functional connectivity networks contribute to the lingering symptoms in patient's experiencing persistent PTHs (P-PTH).
- Structural neuroimaging has shown cortical atrophy in brain regions post-mTBI.
- Functional and Structural compensation may be a finite process in which many younger patient's brains can compensate while others continue to have PTHs.

Objective: To evaluate the extent to which resting state functional network (RSN) connectivity is altered between a cohort of patients with resolved (R-PTH) versus P-PTH relative to healthy controls (CTRL).

Hypothesis: Participants with P-PTH would show distinct functional and structural network abnormalities relative to CTRL and to patients with resolved symptoms.

Methods

Participants with PTH (n=28; F=16) were recruited from the Boston Children's Hospital - Sports Medicine Clinic. Controls (n=12; F=5) were recruited from the Boston Area. Concussion participants were divided into Persistent (P-PTH) (n=15) and Resolved (R-PTH) (n=13) Headaches > 3 Months or < 3 Months respectively.

Rivermead QA and Impact Testing were collected in order to verify symptom severity.

Participants underwent Structural and RSN scans. Group comparisons: Structural (FreeSurfer), RSN (FSL Melodic), Seed Based, symptom reporting, age, and gender.

Figure 1. Resting State Networks identified from group ICA: 1-Primary Visual 2-Occipital Visual 3-Executive 4-Sensorimotor 5-Lateral Visual 6-R-Frontal Parietal 7-Auditory 8-Default Mode 9-L-Frontal Parietal 10-Cerebellar (Z=2.3)

Figure 2. Group characteristics presented at the level of Age, Rivermead Post Concussion Symptom Score, and Total Impact Symptom Score. (CTRL=Control, P-PTH= Persistent, R-PTH=Resolved)

Results

Figure 3. Brain regions showing decreased Resting State connectivity in the evaluated networks comparing CTRL to R-PTH(A) and P-PTH (B).

Figure 5. Functional Connectivity Seed based analysis in the same population with networks that have been Identified in previous mTBI literature. A) Left Hippocampus-Precuneus network B) Right Hippocampus-Frontal Pole C) Insula-Middle Frontal Gyrus in the CTRL, R-PTH, and P-PTH.

Figure 6 Precuneus structural changes A) P-PTH>CTRL Left hemisphere B) P-PTH>CTRL R-Hemisphere C) R-PTH>CTRL R-Hemisphere

Discussion

- Impact Headache Symptom scores were significantly elevated in P-PTH relative to R-PTH despite R-PTH demonstrating greater regions of functional RSN hypoconnectivity.
- Functional RSN hypoconnectivity in ACC for R-PTH with increase in structural size may reflect compensatory change and repair contrasting P-PTH.
- Insula- Middle Frontal may show a compensatory response in R-PTH and it could correlate with an increased structural size.
- Findings in the Insula-Middle Frontal FC for P-PTH showed no change from CTRL and a decrease Insular size. This potentially reflects the Insula's inability to adapt.
- Left Hippocampus- Precuneus shows an increase FC and correlates with an increase in structural size for both concussion groups.

Conclusions

- Findings provide support for the use of fMRI in order to identify abnormal brain activity in PTH subjects.
- Regional functional connectivity varies from hyper to hypo and region of interest. Insular-DMN Resting State connectivity could provide insight into the severity of PTH as shown from previous mTBI and migraine classification studies.
- Findings show evidence of FC being linked to structural changes.

Acknowledgments

Acknowledgements: I would like to thank the Center for P.A.I.N in the Brain at Boston Children's Hospital for their time and guidance. **Funding:** NIH Grant #5R01NS095655-04 and the Peter Morgane Research Fellowship*.