DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF ALCOHOL USE DISORDER IN PATIENTS UNDERGOING BARIATRIC SURGERY

M. Oscoz-Irurozqui¹, S. Llebot¹, G. Adam¹, M.D. Moreno¹, L. Yuguero¹.

¹Benito Menni CASM, Centro de Atención y Seguimiento a las drogodependencias, L'Hospitalet de Llobregat, Spain.

moscozi.hbmenni@hospitalarias.es

Background:

- Recent reviews suggest that a significant proportion of patients undergoing bariatric surgery (BS) develop postoperative problems of alcohol (OH) use¹.
- However, few studies have analyzed characteristics, patterns of consumption and possible risk factors of this population.

Objectives:

- To evaluate demographic and clinical variables in patients with a history of bariatric surgery who have an alcohol use disorder (AUD), specifically:
- Consumption severity,
 - Psychiatric comorbidity
 - Type of surgical intervention
 - Temporal relationship between surgery and AUD onset

Materials and methods:

It is a descriptive study of 11 patients (63.64% women) undergoing bariatric surgery (mean age in surgery 43 years), which perform ambulatory monitoring in a drug addiction center (Table 1).

The information is obtained retrospectively through the review of clinical histories.

Percentages are obtained with Excel

Results:

- Most common modalities of surgery are vertical gastropasty and Roux-Y bypass (36.36% both).
- AUD was established after surgery in 45.46% of patients (mean time 3.6 years later).
- Diagnosis of alcohol abuse is the most frequent (54.55%; mean consumption 161.8g alcohol/day).
- Other psychiatric diagnoses also appear in 81.82% of the sample, and the unspecified adjustment disorder is the most prevalent (33.33%).

Table 1. Clinical characteristics of patients of the sample.

Sex	Patient origin	Age BS	BS type	Consume start	g OH/day (máx.)	Other drugs	Diagnosis OH	Other diagnosis
Female	Day hospital	35	Roux-en-Y bypass	20	120	No	Abuse	Adjustment disorder, fibromyalgia, chronic fatigue
Female	Primary care	39	Roux-en-Y bypass	6	150	Cannabis, cocaine	Moderate dependency	No
Female	Primary care	32	Roux-en-Y bypass	15	120	Cannabis, cocaine	Moderate dependency	Cocaine dependence
Female	Primary care	56	Sleeve gastrectomy	"childhood"	10	No	Mild dependency	Adjustment disorder
Female		44	Roux-en-Y bypass	46	50	No	Abuse	Depressive disorder
Female	Mental health center	60	Sleeve gastrectomy	15	280	Tobacco	Severe dependency	Depressive disorder
Female	CSMA (OH)	61	Sleeve gastrectomy	23	300	No	Abuse	No
Male	Primary care	52	Roux-en-Y bypass	"adolescence"	60	No	Abuse	No
Male	Primary care	33	Sleeve gastrectomy	18	60	Cocaine	Abuse	Adjustment disorder
Male	Hospitalization	35	Gastric balloon	13	300	Tobacco, cocaine, heroine, hallucinogens	Abuse	Cocaine dependence, personality disorder
Male	Primary care	26	Sleeve gastrectomy	14	240	Cannabis, cocaine	Moderate dependency	Cocaine abuse

Conclusions:

- ✓ Predominance of **post-surgical diagnosis** of abuse of **alcohol** as compared with dependence, as well as the finding of a relatively **low intake** of this substance suggest us that anatomical and functional changes post-surgery are in relation with alterations in the metabolism of the toxic, that is **worse tolerated**.
- ✓ Results should be taken into account regarding pre and post-surgical **prevention**, and also in the **intervention**. However, further research is required to elucidate risk factors for the development of AUD post-bariatric surgery.

BIBLIOGRAPHY: 1 Quevedo et al. Rev med Chile. 2015 Jan; 143 (1). 2 Blackburn et al. Addict Biol. 2017 Nov;22(6):1540-1553. 3 Sogg. Surg Obes Relat Dis. 2017 Aug;13(8):1402-1404. 4 Smith et al. Obes Surg. 2018 May;28(5):1248-1254.

Disclosure of interest: Authors report no potential conflicts of interest.