Cortical Microinfarcts Detected with 3T MRI

In Patients with Atrial Fibrillation

Y Soo¹, J Abrigo², KT Leung¹, SF Tsang¹, V Ip¹, K Ma¹, SH Ma¹, V Liu¹, B Lam¹, WC Fong³, SH R Li⁵, PW Ng⁶, J Fok⁷, KK Wong⁷, W Chu², L Wong¹, V Mok¹, T Leung¹.

¹ Department of Medicine and Therapeutics, Prince of Wales Hospital, The Chinese University of Hong Kong.
² Department of Imaging and Interventional Radiology, Prince of Wales Hospital, The Chinese University of Hong Kong.
³ Department of Medicine, Queen Elizabeth Hospital.
⁴ Department of Medicine, Northern District Hospital.
⁵ Department of Medicine, Pamela Youde Nethersole Eastern Hospital.
⁶ Department of Medicine and Geriatrics, United Christian Hospital.
⁷ Department of Medicine, Yan Chai Hospital.

Background

- **Cortical microinfarcts (CMI)** detected by MRI brain, is a recently emerged radiological marker, representing microscopic regions of ischaemic-related cell death at brain cortex. (Figure 1)
- CMI is frequently associated with small vessel disease and are commonly identified in patients with Alzheimer's disease and vascular dementia.
- Recent studies have also demonstrated their association with subclinical cardiac disease, suggesting possible role of microembolism contributing to CMI.

Aims & Hypothesis

- This study evaluated if CMI was associated with clinical ischaemic stroke and transient ischemic attack (TIA) in patients with atrial fibrillation (**AF**).
- We hypothesized that CMI would be more common in AF patients with history of ischaemic stroke TIA.

Methods

- In this prospective study, we recruited patients with atrial fibrillation on oral anticoagulants for 3T MRI brain from April 2014 to April 2017.
- Presence of CMI was correlated with whiter score (rated by Age-related White Matter Change) and clinical history of ischaemic stroke or TIA.

Results

- Total **517** AF patients were recruited.
- **CMI** were observed in **58 patients (11.2%)**, including:
 - Asymptomatic chronic CMI in 45 patients
 - Symptomatic acute CMI in 10 patients
 - Both acute and chronic CMI in 3 patients
- The **mean no. of CMI** was **1.03 ± 0.816 (ranged 1 to 5)**.
- Patients with CMI were more likely to have clinical ischaemic stroke or TIA, and a higher CHA2DS2-VASc score. (Table 1).
- There were no difference in mean white matter score between patients with and without CMI. (Table 1)
- **Asymptomatic chronic CMI** were present in **10.9%** of patients with clinical ischaemic stroke or TIA vs **6.0%** without such history ($p = 0.077$).
- In multiple logistic regression, asymptomatic chronic CMI, white matter score, history of dyslipidaemia and congestive heart failure were **independently associated with clinical ischaemic stroke or TIA**. (Table 2)
- **Combining CMI with CHA2DS2-VASc scores improves risk prediction for clinical ischaemic stroke and TIA marginally**. (Table 3)

Conclusion

- This proof-of-concept study suggests that CMI may **help improve stroke risk prediction** in patients with atrial fibrillation.
- CMI may serve as an adjunctive risk-stratification tool for patients with low CHA2DS2-VASc scores when decision for anticoagulation is difficult base on clinical risk factors alone.

Fig 1. Example of CMI detected with 3T MRI. Chronic CMI was defined as lesions measured < 5 mm in diameter, restricted to the cortex, and perpendicular to the cortical surface, which were hyperintense on FLAIR (A) and T2W images (B), and hypointense on T1W images (C). Lesions with restricted diffusion on DWI were defined as acute CMI.

Table 1. Demographic, clinical and radiological characteristics of patients with and without CMI.

	Patients with CMI (n=58)	Patients without CMI (n= 459)	p
Clinical Data			
Age ± SD	74.38 ± 8.31	72.83 ± 9.04	0.190
Male	32 (55.2%)	253 (55.1%)	1.000
Smoker	15 (25.9%)	146 (31.8%)	0.452
Drinker	5 (8.6%)	44 (9.6%)	1.000
Hypertension	50 (86.2%)	383 (83.4%)	0.707
Hyperlipidaemia	31 (53.4%)	269 (58.6%)	0.482
Diabetes mellitus	16 (27.6%)	157 (34.2%)	0.376
Ischaemic heart disease	5 (8.6%)	41 (8.9%)	1.000
Congestive heart failure	9 (15.5%)	69 (15.0%)	0.848
Previous ischaemic stroke or TIA	48 (82.8%)	302 (65.8%)	0.011
Previous intracerebral haemorrhage	1 (1.7%)	8 (1.7%)	1.000
CHA2DS2-VASc score ± SD	5.26 ± 1.86	4.64 ± 1.81	0.020
Warfarin group	29 (50.0%)	232 (50.5%)	1.000
NOAC group	29 (50.9%)	227 (49.5%)	1.000
Radiological features			
White matter score ± SD	6.43 ± 3.73	6.13 ± 3.67	0.559
Cerebral microbleeds*	21.034 (45.3%)	140 (32.6%)	0.090

*CMBs data was only available in in 482 patients.

Table 2. Multiple logistic regression for clinical ischaemic stroke and TIA, adjusted for age, hypertension, and congestive heart failure.

	OR	95% Confidence Interval		p
		Lower	Upper	
Age	1.011	0.987	1.036	0.358
Hypertension	0.451	0.252	0.806	0.007
Congestive heart failure	0.348	0.206	0.588	<0.001
Dyslipidaemia	3.365	2.225	5.091	<0.001
White matter score	1.105	1.039	1.176	0.001
Chronic CMI	2.364	1.099	5.089	0.028

Table 3. Comparative univariate logistic regression of CMI, CHA2DS2-VASc scores for clinical ischaemic stroke and TIA.

	OR	95% Confidence Interval		p
		Lower	Upper	
CMI	2.495	1.229	5.066	0.011
CHA2DS2-VASc score	2.647	2.227	3.211	<0.001
Combined CMI and CHA2DS2-VASc score	2.655	2.214	3.184	<0.001