Incidence Of Pain At 3 Month Follow Up In Elderly Surgical Patients

Casimiro L.G.¹, Azevedo B.¹, Peixoto A.R.¹, Castro J.S.¹, Teles A.R.¹, Abelha F.^{1,2}
¹Centro Hospitalar São João E.P.E., Dept of Anaesthesiology, Porto, Portugal,
²Faculty of Medicine, University of Porto, Dept of Anaesthesiology, Porto, Portugal

BACKGROUND AND GOUL OF STUDY

Postoperative pain is one of the main factors associated with the increase of postoperative complications. Pain control may therefore accelerate recovery and increase quality of life (QoL). The aim of this study was to evaluate the incidence of pain after surgery at 3 months of follow up, its impact in the patients QoL and possible risk factors.

MATERIALS AND METHODS

It was conducted an observational prospective study.

Inclusion criteria	Exclusion criteria
Patients aged >60 years old	Inability to give informed consent
Submitted to elective surgery	ICU admission after surgery
May – July 2017	

Pre-operatively (T0) it was recorded the Quality of Recovery 15 item score (QoR), the Health-related Quality of Life (EQ-5D-5L),

the World Health Organization Disability Assessment Schedule 2.0 (WHODAS), the Lawton Instrumental Activities of Daily Living (IADL) Scale. These scores were then repeated 3 months (T1) after surgery.

We studied the group of patients that had a higher pain score (HPS) recorded at the EQ-5D-5L, at T1 when compared to T0. This group was compared to those patients whose pain score remain the same or decreased (NHPS). The Chi-square, Fisher's exact or Mann-Whitney U tests were used for comparisons.

RESULTS AND DISCUSSION

A total of 226 patients were included. The incidence of PMP was 21%. every dimensions of EQ-5D-5L at T1 ($p < 0.001$). There were no differences in gender, age, ASA physical status, Body Mass Index, type and length of anesthesia, length of hospital stay.

Variable	NHPS	HPS	P value
Female n(%)	97 (78.9)	26 (21.1)	0.890
Male	82 (79.6)	21 (20.4)	
<80 years old	153 (81.4)	35 (18.6)	0.073
>80 years old	26 (68.4)	12 (31.6)	
Previous chemotherapy n(%)			0.102
Yes	8 (61.5)	5 (38.5)	
No	170 (80.6)	41 (19.4)	
Primary neoplasia n(%)			0.850
Yes	57 (78.4)	16 (21.6)	
No	120 (79.5)	31 (20.5)	
Metastised neoplasia n(%)			0.011
Yes	3 (37.5)	5 (62.5)	
No	175 (80.6)	42 (19.4)	
Pulmonary disease n(%)			0.021
Yes	6 (50.0)	6 (50)	
No	172 (80.8)	41 (19.2)	
Tobaco use n(%)			0.052
Yes	19 (65.5)	10 (34.5)	
No	160 (81.2)	37 (18.8)	
Hypothermia n(%)			0.042
Yes	53 (88.3)	7 (11.7)	
No	126 (75.9)	40 (24.1)	

PONV – Postoperative nausea and vomiting

Variable	NHPS	HPS	P value
Prophylaxis PONV n(%)			0.024
Yes	129 (83.2)	26 (16.8)	
No	49 (70.0)	21 (30.0)	
PONV n(%)			0.205
Yes	12 (92.3)	1 (7.7)	
No	167 (78.4)	46 (21.6)	
Lawton - median (p25-p75)	8 (6-8)	6 (3-8)	0.017
Frailty Clinical Scale n(%)			0.001
<5	138 (84.7)	25 (15.3)	
≥5	41 (65.1)	22 (34.9)	
EQ Mobility n(%)			<0.001
0	97 (89.0)	12 (11.0)	
≥1	82 (70.1)	35 (29.9)	
EQ Self Care n(%)			<0.001
0	123 (88.5)	16 (11.5)	
≥1	56 (64.4)	31 (35.6)	
EQ Usual Activities n(%)			<0.001
0	98 (89.9)	11 (10.1)	
≥1	81 (69.2)	36 (30.8)	
EQ Anxiety / Depression n(%)			<0.001
0	90 (94.7)	5 (5.3)	
≥1	89 (67.9)	42 (32.1)	

CONCLUSION

- In this study the patients with HPS had a poor quality of life at 3 months after surgery in all the EQ-5D-5L domains.
- Patients with HPS had higher rates of dependency at activities of daily living evaluated by Lawton scale.
- The clinical conditions associated with worse pain control, were metastatic neoplasm and pulmonary disease.
- Postoperative inadvertent hypothermia and the use of antiemetic drugs given intraoperatively appears to be a protector factor for the increase of pain 3 months after surgery.