Attitudes towards online psychological counseling in persons with disorders and mental illness

„National Science Centre, Poland”
Registration number:
2018/02/X/HS6/01439

Magdalena Chęć; Maria Ligocka; Krystian Konieczny; Agnieszka Samochowiec
Institute of Psychology, University of Szczecin
Email: magda.chec@gmail.com

I. BACKGROUND

One's attitude towards online psychological help is considered to be a "relatively permanent tendency to positively or negatively evaluate this help, regardless of whether this attitude is expressed in the form of beliefs, feelings or behavioral tendencies" (Chęć et al., 2016). World research shows that the attitude towards online psychological help is diverse and depends on many factors, including gender, age, psychological problems and experience associated with the use of this kind of help. People with disorders and mental illness can benefit from psychological help online, and their attitude towards this help may affect the process of their treatment and recovery.

II. METHODOLOGY

The aim was to determine attitudes towards online psychological help circulating among persons suffering from various mental disorders.

Hypotheses: H1/H2:There are differences in attitudes towards online psychological help between patients with different psychiatric diagnoses; **H2:**There are differences in attitudes towards online psychological help between mentally ill patients and patients with mental disorders; **H3:**Patients with previous experience in using online psychological counseling perceive this kind of help in a more positive way; **H4:**Not using online help stems from perceived greater effectiveness of traditional forms of psychological help.

Method: The study was conducted in north-western Poland, using: a self-constructed Attitude Scale for Online Psychological Assistance – 2, which consists of 10 scales that measure attitudes towards online psychological help: (1) in chat; (2) by Skype; (3) by e-mail; (4) in the form of psychotherapy; (5) in the form of psychoeducation; (6) in the form of a crisis intervention; (7) in the form of psychological advice; (8) to mentally ill people; (9) to people with mental disorders; (10) to people experiencing life difficulties; and a socio-demographic Patient Survey.

Sample: 401 patients - 209 women (52,1%) and 192 men (47,9%), recruited at the Department and Clinic of Psychiatry of the Pomeranian Medical University in Szczecin (Poland). There were 234 mental illness people, 82 with disorders, 76 addicted people and 10 with difficulties.

III. RESULTS

Statistical analyzes were performed: Repeated Measures ANOVA (Non-parametric) and independent Samples T-Test.

H1 has't been confirmed. There aren't differences in attitudes towards online psychological help between patients with different psychiatric diagnoses ($p > .05$). All the diagnoses were merged into four larger groups. The results of the analysis showed that mental illness patients have more possitive attitudes than patients with disorders ($W = -3.442$; $p = 0.015$) and with addictions ($W = -2.874$; $p = 0.042$) towards email help; than patients with addictions ($W = -4.495$; $p < 0.001$) to online help in the form of crisis intervention and than patients with disorders ($W = -3.9299$; $p = 0.005$) and addictions ($W = -3.9980$; $p = 0.005$) to help people with hardship (Tab.1).

H2 has been partially confirmed. Patients with mental illness have a more positive attitude than

patients with disorders, but only when online psychological counseling is by e-mail ($t = 2.1175$; $p = 0.035$), in the form of psychoeducation ($t = 2.1917$; $p = 0.029$) and addressed to people with difficulties ($t = 2.5921$; $p = 0.01$) (Tab.2).

Table 1.

Table 2.

H3 has been partially confirmed. Patients with previous experience in using online psychological counseling have a more positive attitudes, but only to the help provided by email ($t = 2.21799$; $p = .027$), in the form of a crisis intervension ($t = 3.40357$; $p < .001$) and for patients with disorders ($t = 2.34357$; $p = .020$) and difficulties ($t = 2.65841$; $p = .008$) (Tab.3).

Only 1/6 of patients used online psychological counseling.

Table 3.

	Group	N	Mean	Median	SD	SE
E-MAIL	Had	44	40.9	42.0	8.46	1.28
	Hadn't	328	38.2	39.0	7.47	0.412
CRISIS	Had	44	33.0	31.0	7.62	1.15
	Hadn't	320	29.2	29.5	6.88	0.385
DISORDER	Had	44	44.5	43.0	9.41	1.42
	Hadn't	322	41.3	42.0	8.31	0.463
HARSHIPS	Had	44	48.5	49.0	9.21	1.39
	Hadn't	320	44.9	45.0	8.15	0.456

H4 has't been confirmed. Only 15.6% of patients did not use online help because they did not believe in its effectiveness, but over 30% preferred direct contact with a psychologist. As many as 32,3 percent of patients have never heard of this form of help (Tab. 4)

Table 4.

Frequencies of Experience (hadn't)			
Levels	Counts	% of Total	Cumulative %
discourage	5	1.4 %	1.4 %
effectiveness	55	15.6 %	17.0 %
availability	27	7.6 %	24.6 %
computer skills	18	5.1 %	29.7 %
knowledge	114	32.3 %	62.0 %
prefer personal	108	30.6 %	92.6 %
qualification	20	5.7 %	98.3 %
other	6	1.7 %	100.0 %

An additional analysis showed that sex does not differentiate attitudes towards online psychological help ($p > .05$).

IV. CONCLUSIONS

1. People with mental illness have the most positive attitudes among psychiatric patients. 2. The non-use of online help by patients is often due to a lack of knowledge about this possibility and a desire to directly contact the therapist. 3. The patients' experience in using online psychological support favors a more positive assessment of such assistance. Our results suggest an important need for social psychoeducation concerning online psychological help, especially among psychiatric patients.

References: (1) Reynolds, D. J., Stiles, W. B., Bailer, A. J., & Hughes, M. R. (2013). Impact of exchanges and client - Therapist alliance in online-text psychotherapy. *Cyberpsychology, Behavior, and Social Networking*, 16(5), 370e377. (2) Murphy, L., Mitchell, D., & Hallett, R. A. (2011). Comparison of client characteristics in cyber and in-person counseling. *Studies in Health Technology and Informatics*, 167, 149e153. (3) Magdalena Chęć, Maria Ligocka, Martyna Maciejewska, Jerzy Samochowiec, Ewa Łodygowska, Agnieszka Samochowiec. On-line psychological support in the evaluation of specialists and future specialists in Poland. *Computers in Human Behavior* 64 (2016) 703e709 (4) Shroeder, J., Moritz, S., Berger, T., Meyer, B., Lutz, W., Hautzinger, M., Spaeth, Ch., Klein, J., P., Eichenberg, Ch. (2017). Attitudes Towards Internet Interventions Among Psychotherapists and Individuals with Mild to Moderate Depression Symptoms. *Cognitive therapy and research* 41(5), 745-756.