
Oregano Essential Oil in Animal Production

Alma Delia Alarcon-Rojo,
Hector Janacua-Vidales and
Ana Renteria-Monterrubio

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66703>

Abstract

There is an increasing interest in the use of natural additives in food production such as the use of phytogetic feed additives especially for use in swine and poultry. Essential oils are a natural alternative that can be used in animal feed due to their potential health benefits, improved performance and meat production. Oregano essential oil minimises meat downgrading due to transport stress or the modification of the ruminal microorganisms. This review suggested a promising development of food natural preservative against spoilage microorganisms in food systems by the use of oregano essential oil. The addition of oregano essential oil is a good way of preserving meat and could replace the synthetic antioxidants. Moreover, oregano oil and modified atmosphere packaging exhibit an additive preservation effect in fresh meat. Oregano essential oil is effective for controlling the growth of microorganisms. However, for meat quality, special attention should be put on the optimal oregano essential oil dose and meat handling to control or improve the physical, chemical and sensory properties of meat.

Keywords: oregano essential oil, animal production, meat, animal health, packaging, pathogens, sensory, meat quality

1. Introduction

Antimicrobial growth promoters (AGP) were used for decades to increase performance in animal production. However, the link between them and the development of antibiotic resistant microorganisms added to consumer pressure caused major changes. In January 2006, the European Union banned AGP in animal production, which in turn emphasised the importance of researching alternative compounds to promote general health and increase performance in animals [1, 2]. Due to their natural origin and reduced side effects, studies have been mainly focused on herbs, spices and their extracts [3]. The effect of the aromatic plants is

primarily based on their essential oils (EO) and other metabolites [4], for example, vitamins, flavonoids, terpenoids, carotenoids, phytoestrogens, minerals, etc. [6]. Among the activities and applications of EO reported in animal and meat production are antioxidant, preservative, antimicrobial and coccidiostatic and they enhance production of digestive enzymes, stimulate blood circulation and improve immune status. [5, 6–9]. Even though there are 3000 estimated EOs, only 300 of them are of commercial use [10]. However, the EO from oregano (OEO) has been one of the most widely studied due to its content of carvacrol and thymol, and to a lesser extent γ -terpinene, p-cymene and myrcene [11]. Some *in vitro* and *in vivo* properties of OEO are antioxidant, antimicrobial, digestive stimulant, etc. [12–14].

2. Oregano essential oil in animal production

2.1. Poultry and egg production

Results of the effect of OEO in poultry and egg production cannot be considered conclusive or consistent, as they can differ even within researches of the same authors [15, 16]. Studies supporting the positive effects of OEO suggest that they increase performance [17], daily and final body weight in broilers [18] and laying hens [15, 19], nutrient utilisation [20, 21], egg weight and production [15, 19, 22] and feed intake [23]. Improvement in feed efficiency and growth performance might be related with changes in the intestinal morphology, like the increment in villus height to crypt depth ratio [18] or enzyme activity as in the protein digestibility due to the chymotrypsin role [21] and the prevention of coccidiosis [24]. It has been implied that antioxidants from the OEO might transfer into the body of the laying hen, which in turn inhibit the chain reaction associated with lipid oxidation and thus reducing the oxidation in egg yolk [11]. In contrast, several authors have suggested that OEO have no effect on animal performance [21, 25, 26]. For example, Arpášová et al. [16] reported that the addition of thyme and OEO did not significantly influence the body weight, feed consumption and conversion, egg production, mass and weight in laying hens.

2.2. Pig production

As in poultry production, results of the effects of the inclusion of OEO in the diets are contradictory in terms of productive performance [27, 28]. However, positive effects link to dietary inclusion of OEO might be clearer in other areas, for example, immunomodulation [28] and changes in blood counts [29]; in sows, it increases their reproductive performance [30] and causes a reduction in the oxidative stress status and enhances the performance of their litters [31]. Oregano essential oil can be used as well to alleviate stress due to transportation as it improved the antioxidant status [32, 33]. Finally, animals fed 1000 ppm of OEO produced meat of good quality with minimum lipid oxidation [34].

2.3. Milk production and ruminant nutrition

The use of EOs in ruminants and milk production is lightly documented and many of these studies are laboratory based [35]. Several essential oils and their components display antimicrobial

properties that may affect rumen metabolism [36] and influence milk production parameters. Many of these compounds (e.g. flavonoids) have distinct flavours and aromas, which if fed to animals, might change the sensory characteristics of the milk; however, this effect might be temporal [37]. The organoleptic quality of the milk can be affected by feeding EO due to a direct transfer of aromatic compounds from the feed (and environment) to the milk [37], formation of aromatic compounds during digestion of the feed [38] and excretion of the EO in the milk. Ruminal microorganisms utilise nutrients to produce volatile fatty acids; however, this process has energy and protein losses, which render the performance inefficient. These losses might be controlled with the inclusion of EO in the diet to limit the growth of Gram-positive and Gram-negative bacteria [39].

3. Oregano essential oil in the meat industry

3.1. Introduction

This section provides an overview of the applications of OEO in raw and processed meat and fish products. It is well known that raw and minimally processed meat is easily targeted by spoilage microorganisms. Moreover, the interest of the industry to replace synthetic chemicals by natural products with bioactive properties is increasing. The need to reduce the use of additives in foods has highlighted the importance of natural antimicrobials such as essential oils. A wide range of antimicrobial agents derived from essential oils have the potential to be used in food processing and preservation since their antimicrobial activity is well recognised.

3.2. Antioxidant activity

The effects of OEO to extend shelf life by controlling lipid oxidation and improving the sensory qualities of meat and meat products are well documented. Kodal Coskun et al. [40] studied the effectiveness of soy-based edible films incorporated with essential oils from oregano or thyme applied on oxidative stability of ground beef patties. The incorporation of OEO or thyme into the edible films reduced the redness value to an acceptable level, but within the appropriate range. The authors concluded that the addition of OEO and thyme EO into edible films retarded the oxidative changes in meats. The potential application of EOs as natural antioxidants has been studied as well in meat products. OEO (alone or in combination) added to fermented meat products as, according to Tunisian [41] and Spanish [42], sausages did not affect proteolysis and rendered a higher unsaturated fatty acid content without affecting the lipolysis. Even more, OEO-added sausages have a better texture due to an increased hardness. Sausages showed a lower number of enterobacteriaceae, coliforms, *Staphylococcus aureus* and moulds. As in beef, the colour of ground poultry meat (breast and thigh) is stabilised due to the effect of OEO [43]. The combination of (200 ppm) and tannic acid (10 ppm) had the highest effect on TBARS, total carbonyl and off-odours volatiles. Hence, OEO might be a proper replacement for synthetic antioxidants in several types of ground meat [44]. The antioxidant effect of OEO included as well cooked meat. Nieto et al. [45] demonstrated that OEO and rosemary EO retarded the loss of thiols under modified atmosphere packaging (MAP, 70% O₂: 20% CO₂:10% N₂) and aerobic conditions. However, this effect was not observed with

garlic essential oil. Even though the antimicrobial activity against pathogens is well reported, flavour changes have been reported. However, these changes are acceptable [46].

3.3. Bioactive films

Antimicrobial-releasing edible films in food packaging are a form of bioactive packaging. A large number of studies have been focused on this topic. Oregano essential oil has been used in active packaging systems to protect foods from microbial contamination. Studies have shown that OEO in milk protein-based is highly effective against pathogen and spoilage bacteria even when compared with other EOs [47]. OEO not only helps control the growth of microorganisms but it can modify positively the characteristics of the bioactive film [48]. The best balance of mechanical, barrier, thermal, antioxidant and antimicrobial properties is achieved when 9% of OEO is incorporated in poly (lactic acid)/polytrimethylene carbonate films. OEO on alginate-based edible films have the potential to limit lipid oxidation, decrease shear forces, colour and water losses. Oregano added bioactive films also modify consumer perception in terms of odour, flavour and overall acceptance [49–51]. OEO not necessarily need to be included in the wrapping film, as it can be spray in meat exudate absorbent pads to extend shelf life for two more days [52]. Studies showed that oregano essential oil-blend film was an effective antimicrobial suitable for the potential food packaging applications. It has also been demonstrated the effectiveness of oregano oil containing whey protein films to increase the shelf life of fresh beef [53]. They incorporated different levels of oregano oil (0.5, 1.0 and 1.5% w/w in the film forming solution) into sorbitol-plasticised whey protein isolate films and evaluated beef quality. Wrapping of beef cuts with the antimicrobial films resulted in smaller changes in colour in chilled storage. The maximum specific growth rate of total flora and pseudomonads were significantly reduced by a factor of 2 with the use of antimicrobial films (1.5% w/w), while the growth of lactic acid bacteria was completely inhibited. However, interesting results were obtained by Emiroğlu et al. [54] who did not find significant effects on total viable counts, lactic acid bacteria and *Staphylococcus* spp. when oregano was applied on ground beef patties. They evaluated the antibacterial activity of soy protein edible films incorporated with 1, 2, 3, 4 and 5% oregano or thyme essential oils against *Escherichia coli*, *E. coli* O157:H7, *S. aureus*, *Pseudomonas aeruginosa* and *Lactobacillus plantarum*. *E. coli*, *E. coli* O157:H7 and *S. aureus* were significantly susceptible to antimicrobial films; meanwhile, *L. plantarum* and *P. aeruginosa* were more resistant. In the study of Seydim and Sarikus [55], antimicrobial properties of whey protein isolate films containing 1.0–4.0% (wt/vol) ratios of oregano, rosemary and garlic essential oils were tested against *E. coli* O157:H7, *S. aureus*, *Salmonella enteritidis*, *Listeria monocytogenes* and *L. plantarum*. As mentioned before, films containing OEO are more effective than those containing other EOs.

3.4. Antimicrobial effects in packaged raw meat

Foodborne pathogens are commonly associated with raw meats. EOs of spices can be used as biopreservatives due to their antimicrobial properties [56]. Authors have demonstrated that OEO might be more effective against pathogens when used in combination with other natural compounds and technologies. Examples of these combinations include OEO and N,O-carboxymethyl chitosan [57], caprylic acid and vacuum packaging (VP) [58], VP [59] and MAP [60–62]. These treatments might extend shelf life of up to 8 days in different types of raw and

cooked meat [59–61]. The treatments also control the growth of Gram-positive and Gram-negative bacteria, and pathogens, such as *L. monocytogenes* and *Salmonella typhimurium* [58, 59, 61, 62]. The minimum inhibitory concentration for OEO against *S. enteritidis* has been reported as 3.90 µl/ml [63]. A longer shelf life (9 days) than the previously reported was observed in the combination MAP and OEO and thyme EOs [64]. Mixed treatment (OEO, orange dietary fibre and MAP) caused a decrease in TBA values and in microbial counts. This combination did not affect the sensory attributes [65]. OEO and sodium lactate not only reduces the number of microbes but render them more susceptible to heat treatments [66]. These studies showed that OEO, by itself or in combination can be used in raw and processed meats to control pathogens and extend shelf life during chill storage.

3.5. Antimicrobial effects in fish products

The demand for natural alternatives to synthetic additives increases also includes raw and minimally processed fish. The effect of VP and OEO in Mediterranean octopus increases as the concentration of EO increased. The highest concentration of EO increases the shelf life to 20 days, meanwhile half the concentration had only an 11-day shelf life or 3 days when VP was by itself [67]. The effect of EOs is also observed with other EOs, as clove. Clove essential oil incorporated in a gelatin-chitosan film decreased the numbers of Gram-negative bacteria, mainly enterobacteriaceae in chilled-stored fish [68]. In rainbow trout fillets, as in other meats, OEO in combination with MAP decreased the numbers of lactic acid bacteria, H₂S-producing bacteria, enterobacteriaceae and *Pseudomonas* spp. This combination avoided lipid oxidation, however, in contrast with other meats, the highest concentration of OEO impact negatively the sensory traits [69]. The concentration of OEO must be considered in order to balance microbial inhibition and sensory characteristics, because in order to obtain the highest inhibition the sensory traits might be negatively affected [70].

4. Conclusions

Even though some results are contradictory, it seems that the use of OEO as a feedstuff in live animals and production has a positive effect. Some of the effects are indirect or might not be evident, as the use of OEO to minimise meat downgrading due to transport stress or the modification of the ruminal microorganisms. The use of EO is still very much under review, as the doses are not fully developed; however, the antimicrobial action of these compounds is fully studied *in vivo* and *in vitro*. It might be of relevance to study not only the inclusion of EO, but also the inclusion of by-products that result from the production of EO. This review suggested a promising development of food natural preservative against spoilage microorganisms in food systems by the use of oregano essential oil. The effectiveness of oregano essential oil in retarding oxidative changes in meats has been widely demonstrated. The addition of oregano essential oil is a good way of preserving meat and could replace the synthetic antioxidants. Also, oregano oil and modified atmosphere packaging exhibit an additive preservation effect in fresh meat. The oregano essential oil is effective by controlling the growth of microorganisms without detrimental changes in sensory and acceptability attributes.

Author details

Alma Delia Alarcon-Rojo^{1*}, Hector Janacua-Vidales² and Ana Renteria-Monterrubio¹

*Address all correspondence to: aalarcon@uach.mx

1 Autonomous University of Chihuahua, Chihuahua, Mexico

2 Autonomous University of Ciudad Juarez, Ciudad Juarez, Mexico

References

- [1] Barug D, De Jong J, Kies AK, Verstegen MWA. Antimicrobial growth promoters: Where do we go from here? 1st ed. Wageningen: Wageningen Academic Publishers; 2006. 196 p. DOI: 10.3920/978-90-8686-570-3
- [2] Yan F, Azizi A, Janke S, Schwarz M, Zeller S, Honermeier B. Antioxidant capacity variation in the oregano (*Origanum vulgare* L.) collection of the German National Genebank. Industrial Crops and Products. 2016;92:19–25.
- [3] Fatima A, Agrawal P, Singh PP. Herbal option for diabetes: An overview. Asian Pacific Journal of Tropical Diseases. 2012;2(suppl 1): S536–S544.
- [4] Brenes A, Roura E. Essential oils in poultry nutrition: Main effects and modes of action. Animal Feed Science and Technology. 2010;158(1–2):1–14. <http://dx.doi.org/etechconri-cyt.idm.oclc.org/10.1016>
- [5] Calucci L, Pinzono C, Zandomenighi M, Capocchi A. Effects of gamma-irradiation on the free radical and antioxidant contents in nine aromatic herbs and spices. Journal of Agriculture and Food Chemistry. 2013;51:927–934.
- [6] Balandrin MF, Klocke JA. Natural plant chemicals: Sources of industrial and medicinal materials. Science. 1985;228:1154–1160.
- [7] Langhout P. New additives for broiler chickens. World Poultry. 2000;16: 22–27.
- [8] Losa R. The use of essential oils in animal nutrition. In: Feed Manufacturing in the Mediterranean Region. Improving Safety: From Feed to Food. In Proceeding 3rd Conf. Feed Manufacture Mediterranean. Reus, Spain. 2001; pp. 39–44.
- [9] Wenk C. Herbs and botanicals as feed additives in monogastric animals. Asian-Australasian Journal of Animal Science. 2003;16:282–289.
- [10] Burt S. Essential oils: Their antibacterial properties and potential applications in foods—a review. International Journal of Food Microbiology. 2004;94:223–253.
- [11] Florou-Paneri P, Dostas D, Mitsopoulos I, Dostas V, Botsoglou E, Nikolakakis I, Botsoglou N. Effect of feeding rosemary and α -tocopheryl acetate on hen performance and egg quality. Poultry Science. 2006;465:143–149.

- [12] Ertas ON, Güler TC, Dalkılıç B, Simsek ÜG. The effect of an essential oil mix derived from oregano, clove and anise on broiler performance. *International Journal of Poultry Science*. 2005;4:879–884.
- [13] Tomaino A, Cimino F, Zimbalatti V, Venuti V, Sulfaro V, De Pasquale A, Saija A. Influence of heating on antioxidant activity and the chemical composition of some spice essential oils. *Food Chemistry*. 2005;89:549–554.
- [14] Ebani VV, Nardoni S, Bertelloni F, Giovanelli S, Rocchigiani G, Pistelli L, Mancianti F. Antibacterial and antifungal activity of essential oils against some pathogenic bacteria and yeasts shed from poultry. *Flavour and Fragrance Journal*. 2016;31:302–309. DOI: 10.1002/ffj.3318
- [15] Arpášová H. 2011. Phytobiotics to replace antibiotic growth promoters and their impact on the usefulness and quality of the eggs of laying hens type. Nitra : SPU, 2011. 101 s. ISBN 978-80-552-0555-7.
- [16] Arpášová H, Branislav G, Hrnčár C, Fik M, Herkel R, Pistová V. The effect of essential oils on performance of laying hens. *Animal Science and Biotechnologies*. 2015;48:8–14.
- [17] Khattak F, Ronchi A, Castelli P, Sparks N. Effects of natural blend of essential oil on growth performance blood biochemistry, cecal morphology, and carcass quality of broiler chickens. *Poultry Science*. 2014;93:132–137.
- [18] Peng QY, Li JD, Li Z, Duanb ZY, Wua YP. Effects of dietary supplementation with oregano essential oil on growth performance, carcass traits and jejunal morphology in broiler chickens. *Animal Feed Science and Technology*. 2016;214:148–153.
- [19] Suchý P, Strakova E, Mas N, Serman V, Vecerek V, Bedrica L, Lukac Z, Horvat Z. The effect of a herbal additive on performance parameters in layers. *Tierärztliche Umschau*. 2010;65:74–78.
- [20] Amrik B, Bilkei G. Influence of farm application of oregano on performances of sows. *Canadian Veterinary Journal*. 2004;45:674–677.
- [21] Basmacıoğlu MH, Baysal S, Misirlioğlu Z, Polat M, Yilmaz H, Turan N. Effects of oregano essential oil with or without feed enzymes on growth performance, digestive enzyme, nutrient digestibility, lipid metabolism and immune response of broilers fed on wheat-soybean meal diets. *British Poultry Science*. 2010;51(1):67–80.
- [22] Abd EL, Motaal AM, Ahmed AMH, Bahakaim ASA, Fathi MM. Productive performance and immunocompetence of commercial laying hens given diets supplemented with eucalyptus. *International Journal of Poultry Science*. 2008;7:445–449.
- [23] Angelovičová M, Kačaniová M, Angelovič M, Lopašovský, L. 2010. Per os use of Thymi aetheroleum for growth performance of the broiler chickens. *Scientific Journal for Food Industry*. Special Issue 2010,4:127–132.
- [24] Mohiti-Asli M, Ghanaatparast-Rashti M. Dietary oregano essential oil alleviates experimentally induced coccidiosis in broilers. *Preventive Veterinary Medicine* 2015;120(2):195–202.

- [25] Jang IS, Ko YH, Kang SY, Lee CY. Effect of a commercial essential oil on growth performance, digestive enzyme activity and intestinal microflora population in broiler chickens. *Animal Feed Science and Technology*. 2007;134:304–315.
- [26] Lee KW, Everts H, Lankhorst AE, Kappert HJ, Beynen AC. Addition of β -ionone to the diet fails to affect growth performance in female broiler chickens *Animal Feed Science and Technology*. 2003;106:219–223.
- [27] Simitzis PE, Symeon GK, Charismiadou MA, Bizelis JA, Deligeorgis SG. The effects of dietary oregano oil supplementation on pig meat characteristics. *Meat Science*. 2010;84(4):670–676. DOI: 10.1016/j.meatsci.2009.11.001
- [28] Zeng Z, Xu X, Zhang Q, Li P, Zhao P, Li Q, Liu J, Piao X. Effects of essential oil supplementation of a low-energy diet on performance, intestinal morphology and microflora, immune properties and antioxidant activities in weaned pigs. *Animal Science Journal*. 2015;86:279–285. DOI: 10.1111/asj.12277
- [29] Stelter K, Frahm J, Paulsen J, Berk A, Kleinwächter M, Selmar D, Dänicke S. Effects of oregano on performance and immunomodulating factors in weaned piglets. *Archives of Animal Nutrition*. 2013;67(6):461–476.
- [30] Allan P, Bilkei G. 2005. Oregano improves reproductive performance of sows. *Theriogenology*. 2005;63(3):716–721. DOI: 10.1016
- [31] Tan Ch, Wei H, Sun H, Ao J, Long G, Jiang S, Peng J. Effects of dietary supplementation of oregano essential oil to sows on oxidative stress status, lactation feed intake of sows, and piglet performance. *Journal of Nutrition*. 2015;1:1–9.
- [32] Zhang T, Zhou YF, Zou Y, Hu XM, Zheng LF, Wei HK, Giannenas I, Jinc LZ, Peng J, Jiang SW. Effects of dietary oregano essential oil supplementation on the stress response, anti-oxidative capacity, and HSPs mRNA expression of transported pigs. *Livestock Science* 2015;180:143–149.
- [33] Zou Y, Xiang Q, Wang J, Wei H, Peng J. Effects of oregano essential oil or quercetin supplementation on body weight loss, carcass characteristics, meat quality and antioxidant status in finishing pigs under transport stress. *Livestock Science*. 2016;192:33–38. DOI: 10.1016/j.livsci.2016.08.005
- [34] Alarcón-Rojo AD, Peña-Gonzalez E, Janacua-Vidales H, Santana V, Ortega JA. Meat quality and lipid oxidation of pork after dietary supplementation with oregano essential oil. *World Applied Sciences Journal*. 2013;21(5):665–673.
- [35] Benchaar C, Calsamiglia S, Chaves AV, Fraser GR, Colombatto D, McAllister TA, Beauchemin KA. A review of plant-derived essential oils in ruminant nutrition and production. *Animal Feed Science and Technology*. 2008;145(1–4):209–228.
- [36] Newbold CJ, McIntosh FM, Williams P, Losa R, Wallace RJ. Effects of a specific blend of essential oil compounds on rumen fermentation. *Animal Feed Science and Technology*. 2004;114(1–4):105–112.

- [37] Lejonklev J, Kidmose U, Jensen S, Petersen MA, Helwing ALF, Mortensen G, Petersen MA, Helwing ALF, Mortensen G, Weisbjerg MR, Larsen MK. Short communication: Effect of oregano and caraway essential oils on the production and flavor of cow milk. *Journal of Dairy Science*. 2016;99:7898–7903.
- [38] Carpino S, Horne J, Melilli C, Licitra G, Barbano DM, Van Soest PJ. Contribution of native pasture to the sensory properties of Ragusano cheese. *Journal of Dairy Science*. 2004;87:308–315.
- [39] Calsamiglia S, Busquet M, Cardozo PW, Castillejos L, Ferret A. Essential Oils as modifiers of rumen microbial fermentation. *Journal of Dairy Science*. 2007;90(6):2580–2595.
- [40] Kodal Coskun B, Çalikoğlu E, Karagöz Emiroğlu Z, Candoğan K. Antioxidant active packaging with soy edible films and oregano or thyme essential oils for oxidative stability of ground beef patties. *Journal of Food Quality*. 2014;37:203–212.
- [41] El Adab S, Hassouna M. Proteolysis, lipolysis and sensory characteristics of a tunisian dry fermented poultry meat sausage with oregano and thyme essential oils. *Journal of Food Safety*. 2016;36:19–32. DOI: 10.1111/jfs.12209
- [42] Martín-Sánchez AM, Chaves-López C, Sendra E, Sayas E, Fenández-López J, Pérez-Álvarez JA. Lipolysis, proteolysis and sensory characteristics of a Spanish fermented dry-cured meat product (salchichón) with oregano essential oil used as surface mold inhibitor. *Meat Science*. 2011;89(1):35–44. DOI: 10.1016/j.meatsci.2011.03.018
- [43] Alhijazeen M. Effect of oregano essential oil and tannic acid on storage stability and quality of ground chicken meat [thesis]. Ames, Iowa. Iowa State University; 2014.
- [44] Fasseas MK, Mountzouris KC, Tarantilis PA, Polissiou M, Zervas G. Antioxidant activity in meat treated with oregano and sage essential oils. *Food Chemistry*. 2008;106(3):1188–1194.
- [45] Nieto G, Jongberg S, Andersen ML, Skibsted LH. Thiol oxidation and protein cross-link formation during chill storage of pork patties added essential oil of oregano, rosemary, or garlic. *Meat Science*. 2013;95(2):177–184. DOI: 10.1016/j.meatsci.2013.05.016
- [46] Pesavento G, Calonico C, Bilia AR, Barnabei M, Calesini F, Addona R, Mencarellia L, Carmagnini L, Di Martino MC, Lo Nostro A. Antibacterial activity of oregano, rosmarinus and thymus essential oils against *Staphylococcus aureus* and *Listeria monocytogenes* in beef meatballs. *Food Control*. 2015;54:188–199. DOI: 10.1016/j.foodcont.2015.01.045
- [47] Oussalah M, Caillet S, Salmiéri S, Saucier L, Lacroix M. Antimicrobial and antioxidant effects of milk protein-based film containing essential oils for the preservation of whole beef muscle. *Journal of Agriculture and Food Chemistry*. 2004;52(18): 5598–5605. DOI: 10.1021/jf049389q
- [48] Dong L, Hongli L, Lin J, Yongming C, Minglong Y, Haiyun C. Characterization of active packaging films made from poly(lactic acid)/poly(trimethylene carbonate) incorporated with oregano essential oil. *Molecules*. 2016;21(6):1–14. DOI:10.3390/molecules21060695

- [49] Vital AC, Guerrero A, Monteschio JdeO, Valero MV, Carvalho CB, de Abreu Filho BA, Madrona GS, do Prado IN. Effect of edible and active coating (with rosemary and oregano essential oils) on beef characteristics and consumer acceptability. *Plos One*. 2016;11(8):1–15. DOI: 10.1371/journal.pone.0160535
- [50] Benavides S, Villalobos-Carvajal R, Reyes JE. Physical, mechanical and antibacterial properties of alginate film: Effect of the crosslinking degree and oregano essential oil concentration. *Journal of Food Engineering*. 2012;110(2):232–239. DOI: 10.1016/j.jfoodeng.2011.05.023
- [51] Kazemi SM, Rezaei M. Antimicrobial effectiveness of gelatin–alginate film containing oregano essential oil for fish preservation. *Journal of Food Safety*. 2015;35:482–490. DOI: 10.1111/jfs.12198
- [52] Oral N, Vatanserver L, Sezer Ç, Aydın B, Güven A, Gülmez M; Baser KHC, Kürkçüoğlu M. Effect of absorbent pads containing oregano essential oil on the shelf life extension of overwrap packed chicken drumsticks stored at four degrees Celsius. *Poultry Science*. 2009;88(7):1459–1465.
- [53] Zinoviadou KG, Koutsoumanis KP, Biliaderis CG. Physico-chemical properties of whey protein isolate films containing oregano oil and their antimicrobial action against spoilage flora of fresh beef. *Meat Science*. 2009;82(3):338–345. DOI: 10.1016/j.meatsci.2009.02.004
- [54] Emiroğlu ZK, Yemis GP, Coskun BK, Candoğan K. Antimicrobial activity of soy edible films incorporated with thyme and oregano essential oils on fresh ground beef patties. *Meat Science*. 2010;86(2):283–288.
- [55] Seydim AC, Sarikus G. Antimicrobial activity of whey protein based edible films incorporated with oregano, rosemary and garlic essential oils. *Food Research International*. 2006;39(5):639–644. DOI: 10.1016/j.foodres.2006.01.013
- [56] Marques JL, Volcão LM, Funck GD, Schneid Kroning I, Padilha da Silva W, Fiorentini ÂM, Ribeiro GA. Antimicrobial activity of essential oils of *Origanum vulgare* L. and *Origanum majorana* L. against *Staphylococcus aureus* isolated from poultry meat. *Industrial Crops and Products*. 2015;77(23):444–450. DOI: 10.1016/j.indcrop.2015.09.013
- [57] Khanjaria A, Karabagias IK, Kontominas MG. Combined effect of N, O-carboxymethyl chitosan and oregano essential oil to extend shelf life and control *Listeria monocytogenes* in raw chicken meat fillets. *LWT—Food Science and Technology*. 2013;53(1):94–99. DOI: 10.1016/j.lwt.2013.02.012
- [58] Hulankova R, Borilova G, Steinhauserova I. Combined antimicrobial effect of oregano essential oil and caprylic acid in minced beef. *Meat Science*. 2013;95(2):190–194. DOI: 10.1016/j.meatsci.2013.05.003
- [59] Ntzimani AG, Gitrakou VI, Savvaidis IN. Combined natural antimicrobial treatments (EDTA, lysozyme, rosemary and oregano oil) on semi cooked coated chicken meat stored in vacuum packages at 4°C: Microbiological and sensory evaluation. *Innovative Food Science & Emerging Technologies*. 2010;11(1):187–196.

- [60] Chouliara E, Karatapanis A, Savvaidis IN, Kontominas MG. Combined effect of oregano essential oil and modified atmosphere packaging on shelf-life extension of fresh chicken breast meat, stored at 4°C. *Food Microbiology*. 2007;24(6):607–617. DOI: 10.1016/j.fm.2006.12.005
- [61] Tsigarida E, Skandamis P, Nychas GJE. Behaviour of *Listeria monocytogenes* and autochthonous flora on meat stored under aerobic, vacuum and modified atmosphere packaging conditions with or without the presence of oregano essential oil at 5°C. *Journal of Applied Microbiology*. 2000;89:901–909. DOI: 10.1046/j.1365-2672.2000.01170.x
- [62] Skandamis P, Tsigarida E, Nychas GJE. The effect of oregano essential oil on survival/death of *Salmonella typhimurium* in meat stored at 5°C under aerobic, VP/MAP conditions. *Food Microbiology*. 2002;19(1): 97–103.
- [63] Coutinho de Oliveira TL, de Araújo SR, Hilsdorf PR. A Weibull model to describe antimicrobial kinetics of oregano and lemongrass essential oils against *Salmonella enteritidis* in ground beef during refrigerated storage. *Meat Science*. 2013;93(3):645–651. DOI: 10.1016/j.meatsci.2012.11.004
- [64] Karabagias I, Badeka A, Kontominas MG. Shelf life extension of lamb meat using thyme or oregano essential oils and modified atmosphere packaging. *Meat Science*. 2011;88(1): 109–116. DOI: 10.1016/j.meatsci.2010.12.010
- [65] Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J, Pérez-Álvarez JA. Effect of orange dietary fibre, oregano essential oil and packaging conditions on shelf-life of bologna sausages. *Food Control*. 2010;21(4):436–443. DOI: 10.1016/j.foodcont.2009.07.004
- [66] Juneja VK, Hwang CA, Friedman M. Thermal inactivation and postthermal treatment growth during storage of multiple *Salmonella* serotypes in ground beef as affected by sodium lactate and oregano oil. *Journal of Food Science*. 2010;75:M1–M6. DOI: 10.1111/j.1750-3841.2009.01395.x
- [67] Atrea I, Papavergou A, Amvrosiadis I, Savvaidis IN. Combined effect of vacuum-packaging and oregano essential oil on the shelf-life of Mediterranean octopus (*Octopus vulgaris*) from the Aegean Sea stored at 4°C. *Food Microbiology*. 2008;26(2):166–172. DOI: 10.1016/j.fm.2008.10.005
- [68] Gómez-Estaca J, López de Lacey A, López-Caballero ME, Gómez-Guillén MC, Montero P. Biodegradable gelatin–chitosan films incorporated with essential oils as antimicrobial agents for fish preservation. *Food Microbiology*. 2010;27(7):889–896. DOI: 10.1016/j.fm.2010.05.012
- [69] Pyrgotou N, Giatrakou V, Ntzimani A, Savvaidis IN. Quality assessment of salted, modified atmosphere packaged rainbow trout under treatment with oregano essential oil. *Journal of Food Science*. 2010;75:M406–M411.
- [70] Van Haute S, Raes K, Van der Meeren P, Sampers I. The effect of cinnamon, oregano and thyme essential oils in marinade on the microbial shelf life of fish and meat products. *Food Control*. 2016;68:30–39. DOI: 10.1016/j.foodcont.2016.03.025

