Persistent deficiency of MAIT cells in dermatomyositis

C.CASSIUS^{1,2}, M.BRANCHTEIN², M.BATTISTELLA^{1,3}, C.LEPELLETIER^{1,2}, A. DE MASSON^{1,2}, A.BENSSUSAN², M.BAGOT^{1,2}, H.LE BUANEC^{2*}, JD.BOUAZIZ^{1,2*}

¹Dermatology Department, ²IRSL-INSERM U976, ³Pathology Department, Saint-Louis hospital, Paris, France. *co-last

Introduction

Dermatomyositis (DM) is a chronic autoimmune disorder affecting skin and muscle.

Mucosal associated invariant T (MAIT) cells are innate like lymphocytes that can be activated either through their semi-invariant T-cell receptors or through unspecific stimulation.

MAIT cells play a role in anti-bacterial responses but they have also been implicated in autoimmune disorders

To assess a potential role of MAIT cells in DM, we analyzed the frequency and activity of MAIT cells in untreated DM patients.

Methodology

Patients: 22 DM patients were enrolled in this study (ENMC criteria). No patient received immunosuppressive drugs before sample collection. 5 moderate to severe atopic dermatitis and 7 severe psoriasis were included as inflammatory controls. 19 healthy controls (HC) were included.

Cell preparation: Peripheral venous blood of patients and healthy donors was diluted with phosphate-buffered saline (PBS), and the peripheral blood mononuclear cells were isolated by density gradient centrifugation on Ficoll-Hypaque and stored liquid frozen until further use.

Staining and flow cytometric analysis: A 18-color immunophenotyping approach was used for the identification of lymphocyte subpopulations. MAIT cells were identified using the 5-OP-RU tetramer which was obtained from the NIH

Results

1 Gating strategy to identify MAIT cells

2 Circulating MAIT-cells were dramatically decreased during DM ((median: 0.25 [0.19-0.6%] vs. 2.32% [1.18-4.45%], p<0.0001)

3 Residual MAIT cells had a peculiar phenotype: increased CD4/CD8 ratio and decreased expression of both CD26 and CD161

Residual MAIT cells expressed higher level of activation (CD25, CD45RA, CD28) and exhaustion (CD39, CTLA4) markers

MAIT cells frequency inversely correlated with CD25 as well as CTLA4 expression on MAIT-cells ($r = -0.7228$, $p = 0.0002$ and $r = -0.6476$, $p = 0.0015$)

Conclusion

This study demonstrates that MAIT cells are deficient in active DM patients with an abnormal phenotype suggesting activation induced cell death.

These abnormalities may contribute to dysregulated cutaneous and systemic immunity in DM patients.