Severe burn in child with hypoplastic left heart syndrome

Anna Ziola, Andrzej M. Bulandra, Tomasz Koszutski
Department of Paediatric Surgery and Urology,
Medical University of Silesia
Uppersilesian Child Health Center, Katowice, Poland

INTRODUCTION

A case of 3-years old girl with extensive burn injury and hypoplastic left heart syndrome is presented.

Hypoplastic left heart syndrome (HLHS) is a complex defect involving the left ventricle hypoplasia, stenosis or occlusion of mitral and aortic valve, as well as hypoplasia of the ascending aorta and the arch.

Presented patient suffered from HLHS.

She was operated by cardiosurgeons in first year of life – the Norwood procedure and hemi-Fontan operation were made. At the time of burn injury she was planned for the second stage of Fontan procedure.

CASE PRESENTATION

The girl was scalded by hot tomato sauce. The burned area covered 30% TBSA - face, anterior surface of chest and abdomen, right arm and forearm as well as right thigh.

Procedures at admission:

- wounds debridement
- biological dressing – amnion – on face and neck
- antibacterial dressings on rest of the burned area

TREATMENT PROBLEMS

low tissue perfusion and low blood oxygenation
(caused by heart defect)

prolonged demarcation
of necrotic tissue

HLHS related
haemodynamic instability
during anaesthesia

no possibility of long term
surgical procedures

solved by multi-stage short
term wound debridement

wounds after 25 days

wounds after 42 days
and 6 surgical debridement

FINAL EFFECT

completely healed wounds
after 133 days

scars after laser therapy

Conclusion

Effective treatment of severe burned child with a complex heart defect is very difficult. It requires the involvement of many specialists and the use of unconventional treatment methods.