Recognition of Early Postoperative Delirium in the Post Anesthesia Care Unit with Confusion Assessment Method by Nurses during Routine Daily Practice

Verda Toprak, Hakan Nuraç, Leyla Çağlayan, Burcu Aydoğan

Başkent University, İstanbul Hospital, Anesthesiology and Reanimation Department-
İstanbul, TURKEY

- **Background and Goal of the Study:**

- There is limited data about emergence delirium in the Post Anesthesia Care Unit (PACU). Recognition of delirium in the PACU is very important in order to avoid further worse outcomes. We aimed to examine the frequency of emergence delirium among ASA I-III patients following anesthesia.

- **Materials and Methods:**

- Ethics committee approval (KA17/227)
- Retrospective cohort
- Statistics: Multivariate logistic regression
- Setting: Postoperative recovery area
- Number of patients included in the study: 3811
- Exclusion criteria: Extubation outside the operating theatre, psychiatric disorders, non-Turkish speaking or deaf people, severe dementia, neuromuscular disorders.
- Inhalation anesthetic used: Sevoflurane
- The Confusion Assessment Method (CAM) was used to evaluate delirium signs during PACU stay. Multivariate logistic regression was used to assess potential, risk factors for delirium during PACU stay including age, ASA status, surgical type, benzodiazepin premedication, long surgery, and surgical type.

- **Results and Discussion:** During the analysed period 5200 patients were operated. With the exclusion of child patients, missed forms and local anesthesia patients 3811 patients were included to the study. Delirium in general anesthesia patients was 43/2976 (1.4 %). Delirium in regional anesthesia with sedation patients was (4/788) (0.5 %). Delirium was observed

Variable	n=3811	
Age	66 (18-102)	
Sex ratio (M/F)	2673/1138	
ASA classification		
1	890	
2	1800	
3	1121	
Surgical operation types and incidence of delirium	General Anesthesia	Regional anesthesia with sedation
General surgery	0/229	1/267 (0.03 %)
Urology	3/323 (0.93 %)	0/105
Orthopedics	0/332	3/88 (0.3 %)
Neurosurgery	0/334	
Vascular Surgery	43 (1.4 %)	
Ear Nose and Throat Surgery	0/229	
Plastic Surgery	3/323 (0.93 %)	
Obstetrics Surgery	0/332	
Total	0/334	0/320
	43 (1.4 %)	4 (0.5 %)
Anesthetic duration	145 (25-180 minutes)	
Benzodiazepine (midazolam)	2 (0-2 miligrams)	
Aldrete score		
PACU admission	8 (8,9)	
PACU discharge	10 (9,10)	

Conclusion(s): Anesthesia nurses and technicians can recognize delirium with CAM-ICU test at PACU. But we recognized that hypoactive delirium was not met by the nurses; and may be because of this, delirium incidence was lower than the general postoperative delirium incidence.

References: Emergence from general anesthesia and evolution of delirium signs in the postanesthesia care unit. Card E. Et al. British Journal of Anaesthesia, 2015;411-17.