Cyber-addiction among medical students

B. Ben Mohamed, Z. Azouz, S. Khedher, A. Abbadi, F. Zaâfrane, L. Gaha
Psychiatry department, the university Hospital of Monastir, Tunisia .

Introduction:

Internet is becoming a basic feature of global civilization. Young people are believed to be the main Internet users. They are also more likely to develop a problematic internet use (PIU) such as unethical, excessive and addictive internet use.

Materials and Methods:

A self-administered questionnaire was filled out by volunteer medical students from the University of Medicine of Monastir (Tunisia) between October and November 2018. The collected data included socio-economic characteristics and Generalized Problematic Internet Use Scale (GPIUS).

Objectives:

The aim of the present study was to assess Internet addiction among medical students and determine its predictive factors.

Results:

- A total of 200 students were included. The mean age was 22 years with a sex-ratio M/F of 1.81.
- In our sample 36.5% were enrolled in the first cycle of medical studies, 44% in the second cycle of medical studies and 19.5% were interns.

Figure 1: Distribution of participants by type of Internet use

Figure 2: Other addictions

Table: Factors associated with PIU

	Problematic Internet Use		No Problematic Internet Use		p
	N	%	N	%	
The daily time spent online >8 hours	11	28.9	27	71.1	0.003
The internet use before bedtime	25	15.2	139	84.8	0.044
The use of internet for playing video games	10	28.6	25	71.4	0.005
Looking for virtual friends	08	25	24	75	0.042

Discussion:

- ✓ In our study, among the 200 participants, 26 (13%) participants had PIU. Various studies have estimated the prevalence of internet addiction from 4 % (1) to 54 % (2) depending on the study population, the diversity of assessment tools applied and the social contexts.
- ✓ The prevalence of PIU was higher among female students. However, different studies suggested that internet addiction appears to have a male preponderance (3). The reason behind the difference may be the increase in female medical students in Tunisia. Thus, females should not be ignored in PIU prevention programs.
- ✓ Consistent with our results, a handful of studies reported that the age of internet users was not associated with the risk of PIU. All age groups may be affected and there is no age threshold (4) . However, lesser age at first Internet use was correlate to PIU.
- ✓ PIU had no significant correlation with other addictions in our study. These findings aren't consistent with other studies reporting drug use (5), and problematic alcohol use (6) in problematic internet users. These comorbidities may be suggestive of similar etiology and increased severity of psychopathology .
- ✓ In our group, PIU was found to be associated with different factors (Table 1). In the study of John and Kavitarati , PIU was significatily associated to similar factors to know: a daily connection time of 2 h/d, the fact of connecting usually at night but also in the morning, and the use for internet for education, entertainment, online shopping and social media (7) .

Conclusion:

Our findings indicate that Problematic Internet Use is common among medical students. We recommend setting a preventive strategy that acts on the main factors that were identified in our study and in the literature.

References:

- (1) Christakis DA, Moreno MM, Jelenchick L. Problematic internet usage in US college students: a pilot study. BMC Med. 2011;9(1):77.
- (2) Mellouli M, Zammit N, Limam M, Zedeni CH. Prevalence and predictors of internet addiction among college students in Sousse, Tunisia. J Res Health Sci. 2018;18(1):e00403.
- (3) Chahoth V, Kodavanji B, Kumar NA, Anupama N, Kini RD, Pai SR. Impact of internet use on lifestyle in undergraduate medical students. Int J Biomed Res.2014;05:190-2.
- (4) Benguigui J .La cyberdependance:Données de la littérature et résultats d'une étude sur 48 cas [Thèse de Doctorat de Médecine].Paris:Université Paris 7 Denis Diderot Faculté de Médecine;2009.
- (5) Gong, J., Chen, X., Zeng, J., Li, F., Zhou, D., & Wang, Z. (2009). Adolescent addictive Internet use and drug abuse in Wuhan, China. Addiction Research and Theory, 17(3), 291–305
- (6) Ko, C.-H., Yen, J.-Y., Yen, C.-F., Chen, C.-S., Weng, C.-C., & Chen, C.-C. (2008a). The association between Internet addiction and problematic alcohol use in adolescents: The problem behavior model. CyberPsychology and Behavior,11(5), 571–576
- (7) John S, Kavitarati D. Prevalence and risk factors of internet addiction in medical students. Int J Med Sci Public Health. 2018;7:595-600