Anti-inflammatory responses in cultured endothelial cells exposed to two conjugated linoleic acids

Valenzuela, C.; Miles E.A.; Calder, P.C.

Human Development & Health, Faculty of Medicine, University of Southampton, Southampton, United Kingdom.
cav1g15@soton.ac.uk

BACKGROUND

- Conjugated linoleic acids (CLAs): positional and geometric isomers of linoleic acid.
- Naturally produced in the rumen of cattle (bacterial fermentation). Non-ruminant animals can endogenously produce the isomer cis-9, trans-11 (CLA9,11).
- CLA9,11 is the most predominant isomer (meat and dairy) (90%). The remaining 10% is mostly trans-10, cis-12 (CLA10,12) CLA isomer.
- Reported beneficial biological properties (animal models and cultured cells) of CLAs: anticarcinogenic, reduction in body fat deposition, reduced development of atherosclerosis, modulation of immune and inflammatory responses.
- Mechanisms of action are still controversial.
- Whether low concentrations of CLAs are able to affect basal, unstimulated endothelial cell (EC) responses is not clear.

Aim: To evaluate the effects of two CLAs on basal inflammatory responses of cultured EA.hy 926 endothelial cells.

METHODS

- EA.hy 926 (HUVEC)
- 32000 cells/cm²

Endothelial Cell model

- Linoleic acid (LA), CLA9,11, CLA10,12
- 1 and 10 µM, 48 hrs incubation

Fatty acids

- MTT
- GC
- Luminex
- RT-PCR
- Adhesion assay

Assays

RESULTS

Fig 1. Viability of EA.hy926 cells after incubations for 48 hrs with supplemented DMEM containing 0.1% of ethanol (CTL) or different concentrations (0, 1, or 10) of FA. Bars are mean ± SEM of 9 samples performed in 3 experiments. ANOVA ONE WAY, Tukey. ****p<0.0001.

Fig 2. Incorporation of LA (A), CLA9,11 (B) and CLA10,12 (C). EA.hy926 cells were incubated for 48 hrs with DMEM containing 0.1% of ethanol (Control) or different concentrations (1 and 10 µM) of FA. Bars are mean ± SEM of 6 to 9 samples performed in 3 experiments.

Figure 3. Production (pg/mL) of MCP-1 (A), VEGF (B), RANTES (C), ICAM-1 (D), IL-8 (E) and IL-6 (F) by EA.hy926 cells incubated for 48 hrs with DMEM containing 0.1% of ethanol (Control) or two concentrations (1, 10 µM) of FA. Bars are mean ± SEM of 9 samples performed in 3 experiments. ANOVA TWO WAY with Tukey as post-hoc test. *p<0.05, **p<0.01, ***p<0.0001 vs. Control. \$\$\$\$p<0.0001, differences between fatty acids at the same concentration.

Figure 4. Relative gene expression of NFkB1 (A), PPAR-α (B), PTGS2 (for COX-2)(C) and IL-6 (D) in EA.hy926 cells incubated for 48 h with 1 and 10 µM of FAs in DMEM containing 0.1% of ethanol (Control). Cq values were normalized by the geometric mean of reference targets YWHAZ and RPL13A genes. Bars are mean ± SEM of 9 samples performed in 3 experiments. ANOVA ONE WAY with Tukey as post-hoc test. *p<0.05; **p<0.01; ****p<0.0001.

Figure 5. Adhesion of THP-1 cells (% of control) to EA.hy926 cells incubated for 48 h with DMEM containing 0.1% of ethanol (Control) or 1 µM (A) and 10 µM (B) of FAs, followed by incubation with DMEM for 24 h and 1 hr co-incubation with THP-1 cells. Bars are mean ± SEM of 9 samples performed in 3 experiments. ANOVA ONE WAY with Tukey as post-hoc test: **p<0.01; ***p<0.001; ****p<0.0001.

Figure 6. Visualisation of adhesion of calcein labelled THP-1 cells to EA.hy926 cells after FA exposure at 10 µM. Adhesion of THP-1 cells to activated EA.hy926 cells without prior incubation with FA (control (A)) or with 48 h prior exposure to linoleic acid (B), CLA9,11 (C), CLA10,12 (D) followed by incubation with DMEM for 6 h and 1 h co-incubation with calcein labelled THP-1 cells. Attached THP-1 cells were visualised by fluorescence microscope Nikon Elipse Ti at a magnification of 100x under transmitted light.

SUMMARY

- Cell viability at 1 and 10 µM is similar in all the experimental conditions used and approaches 100%.
- FA tested are incorporated into the cells in a dose-dependent manner.
- Pre-treatment with CLA9,11 reduced unstimulated basal concentrations of MCP-1, RANTES and IL-8, showing also a trend to diminish all other mediators measured. Pre-treatment with CLA10,12 reduced unstimulated basal concentrations of MCP-1, RANTES and IL-8, although it increased IL-6.
- Both CLAs at 10 µM increased the relative expression of NFkB, while decreasing the relative expression of PPARα, COX-2 and IL-6.
- Only LA significantly decreased THP-1 adhesion when used at 10 µM.

CONCLUSION

These results suggest that both CLAs at a low concentration have a neutral or modest anti-inflammatory effect in basal conditions, which may influence endothelial function and risk of vascular disease. Interestingly, at these low CLA concentrations some pro-inflammatory genes are upregulated while others are down regulated. This suggests complex effects of CLAs on inflammatory pathways.